

Systematic Review

Systematic Review of the Role of BCG in the Treatment of Urothelial Carcinoma of the Prostatic Urethra

Oliver Patschan^{a,*}, Philippe E. Spiess^b, George N. Thalmann^c, Joan Palou Redorta^d and Georgios Gakis^e

^a*Institution of Translational Medicine, Lund University, Malmö, Sweden*

^b*Department of GU Oncology and Department of Tumor Biology, Moffitt Cancer Center, Tampa, FL, USA*

^c*Department of Urology, University Hospital, Inselspital Bern, Switzerland*

^d*Department of Urology, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Fundació Puigvert, Barcelona, Spain*

^e*Department of Urology und Pediatric Urology, University Hospital of Würzburg, Würzburg, Germany*

Received 16 November 2020

Accepted 28 February 2021

Pre-press 16 March 2021

Abstract.

BACKGROUND: In patients with non-invasive urothelial carcinoma of the prostatic urethra (PUC), treatment with Bacillus Calmette-Guárin (BCG) could be beneficial.

OBJECTIVE: To assess the response rates to BCG in the different tumor stages, to describe the clinical impact of transurethral resection of the prostate (TURP) before BCG treatment, and to review the side effects of BCG treatment for PUC.

METHODS: A systematic search was conducted using the PubMed database to identify original studies between 1977 and 2019 reporting on PUC and BCG.

RESULTS: Of a total of 865 studies, ten were considered for evidence synthesis. An indication for BCG treatment was found in non-stromal invasive stages (Tis pu, Tis pd) and in stromal infiltrating cases (T1) of primary and secondary PUC when transitional cell carcinoma was the histology of origin. Studies including patients treated with TURP before BCG showed a better local response in the prostatic urethra with a higher DFS (80–100% vs. 63–89%) and PFS (90–100% vs. 75–94%) than patients in studies in which no TURP was performed. However, this difference in recurrence and progression in the prostate neither affected the total PFS (57–75% vs. 58–93%), nor the disease specific survival (70–100% vs. 66–100%).

CONCLUSIONS: The use of resection loop biopsies of the prostatic urethra in appropriate cases during the primary work-up for suspected PUC, as well as the use of the current TNM classification for PUC, need to be improved. BCG therapy for non-stromal invasive stages of PUC show a good local response. Local response is further improved by a TURP before BCG therapy, although the overall prognosis does not seem to be affected. Further evidence for BCG treatment in the rare cases of stromal invasive PUC is needed. Specific side effects of BCG treatment for PUC are not reported.

*Correspondence to: Oliver Patschan, MD, PhD, Institution of Translational Medicine, Lund University, Malmö, Sweden.
E-mail: oliver.patschan@med.lu.se. <https://orcid.org/0000-0001-5788-794X>

INTRODUCTION

Urethral carcinoma is a rare disease, accounting for less than 1% of all urogenital cancer cases [1]. In the

28 European Union countries, it is estimated that 655 patients are being diagnosed with urethral carcinoma annually, striking patients from the fifth decade of life and with a peak incidence in patients 75 years of age [2, 3]. Primary urethral carcinoma is detected in patients without a previous history of urothelial cancer. Secondary urethral carcinoma occurs during the follow up of a known urothelial cancer of the bladder or the upper urinary tract (UUT).

The histological origin of urethral carcinoma differs between the sexes: In women, adenocarcinoma is the most frequent histology (38–47%) followed by SCC (25–28%), UC (25–28%) and other histological entities (6%) [4, 5]. In women, both surgery and radiation therapy are practical treatment options [6]. In men, approximately 80% of the cases of urethral carcinoma are urothelial cancers, followed by squamous cell carcinoma (15%) and adenocarcinoma (5%), respectively [7]. The histologic features of these cancers vary by anatomical location: Urethral cancers in the penile or bulbar urethra are of squamous cell differentiation in about 90% of the cases. In contrast, in the prostatic urethra, 90% of the cancers are of urothelial origin [8]. Generally, the primary treatment of male urethral carcinoma is surgical excision. In the prostatic urethra, transurethral resection is often the first diagnostic and therapeutic step. In invasive growth of prostatic urethral carcinoma (PUC), radical cystoprostatectomy, either before or after chemotherapy, is indicated.

Bacillus Calmette-Guárin (BCG) is the currently most potent drug for intravesical treatment of high-risk non-muscle invasive bladder cancer (NMIBC). It is known to improve intravesical recurrence-free survival and might even improve progression-free survival in non-muscle invasive bladder cancer [9]. Since PUC originates almost exclusively from the urothelium, BCG is a relevant drug even for the treatment of urethral carcinoma of the prostatic urethra. In patients without stromal invasion (Tis pu, Tis pd) and even with stromal invasion (T1) of urothelial carcinoma of the prostatic urethra, treatment with BCG might be beneficial [6]. However, the evidence for BCG treatment, with or without transurethral resection of the prostate (TURP) before treatment induction, has been sparse. This systematic review aims to elucidate the clinical impact of BCG treatment in the different stages of urothelial carcinoma of the prostate. It also aims to assess the effect of TURP before BCG treatment on the risk for recurrence and progression, and to review the side effects of BCG treatment for PUC.

EVIDENCE ACQUISITION

A systematic review was carried out based on a literature search by PubMed/Medline. Due to the low incidence of the disease, no prospective clinical trials on BCG treatment in PUC were available. This explains why a systematic review according to the PICO description was not possible. All authors participated in the process of literature search and data acquisition process.

This literature search was aimed at identifying all articles that published the results of cohort studies and retrospective clinical studies as full-length articles published in English between 1977 (date of the first publications relative to urethral carcinoma) and March 2019. Case reports and reviews were excluded.

The following keywords were used in the database just cited: Bacillus-Calmette-Guárin OR BCG AND urethral carcinoma OR urothelial cancer or prostatic urethra; carcinoma in situ OR CIS AND urethral carcinoma OR urothelial cancer prostatic urethra; radical cystectomy AND urethral carcinoma OR urothelial cancer prostatic urethra; Bacillus-Calmette-Guárin OR BCG AND urethral carcinoma OR urothelial cancer prostatic urethra AND survival; Bacillus-Calmette-Guárin OR BCG AND urethral carcinoma OR urothelial cancer prostatic urethra AND transurethral resection prostate; Bacillus-Calmette-Guárin OR BCG AND urethral carcinoma OR urothelial cancer prostatic urethra AND radical cystectomy; Bacillus-Calmette-Guárin OR BCG AND urethral carcinoma OR urothelial cancer prostatic urethra AND side effects.

The reference lists of all systematic reviews in the field were screened for additional references. After a first selection, based on the title and abstract of the papers, duplicates were removed. Once selected, the full text of the articles was studied to gather information about study design, inclusion criteria, baseline patient characteristics, TNM-stage, treatment regimen, follow up, disease free survival (DFS), progression free survival (PFS), and disease specific survival (DSS).

EVIDENCE SYNTHESIS

BCG-response in the different tumor stages of prostatic urethral carcinoma

BCG treatment was reported in two clinical scenarios of PUC: Primary and secondary urothelial carcinomas of the prostatic urethra [10–19]. No




Fig. 1. Consolidated Standards of Reporting Trials diagram outlining the selection process of the included studies. Date of search: 30th september 2019.

articles were found describing BCG treatment in patients with primary urethral carcinoma of the anterior urethra. There were also no articles specifically addressing primary cases of PUC without concomitant carcinoma of the urinary bladder or the upper urinary tract. Figure 1 shows the selection process for the included articles. A total of ten retrospective studies focused on patient outcomes in patients with PUC treated with BCG. A total of 162 patients were included in the studies on this topic, of which only 5 patients were shown to have tumor stage T1. The study by Taylor et al. [15] is an update of the study by Schellhammer et al. [18], which in turn is an update of the study by Hillyard et al. [19], and it cannot be ruled out that some of the patients included in these studies were examined twice. BCG was used only in superficial stages of PUC (Tis pu, Tis pd, T1). The TNM classification of these tumor stages is shown in Table 1 [20]. Data from all selected studies are shown in Table 2.

Almost two-thirds of patients who did not have TURP prior to BCG therapy had no recurrence (DFS 63–89%) in the prostatic urethra during follow up, and at least three out of four of these patients did not

Table 1

TNM classification (eighth edition) for urethral carcinoma. Non-muscle invasive stages of urothelial carcinoma of the prostatic urethra only [20]

Tis pu	Carcinoma in situ, involvement of prostatic urethra
Tis pd	Carcinoma in situ, involvement of prostatic ducts
T1	Tumor invades subepithelial connective tissue (fortumors involving prostatic urethra only)

Urothelial (transitional cell) carcinoma of the prostate.

experience tumor progression to the prostate during follow up (PFS 75–91%). In four out of ten studies, only patients with primary urethral carcinoma were included. Only three out of ten studies differentiated between tumor stages, i.e. Tis pu versus Tis pd versus T1. The results of these studies were recorded and presented in different ways, which made it impossible to assess the different response rates of BCG therapy in the different tumor stages of PUC. In addition, the number of cases in the studies would have been too small to obtain reliable prognostic information. A study by Ovesen et al. [11] examined the Danish BCG strain and not the commonly used Pasteur or Connaught BCG strains. In the study by Taylor et al. [15] the strain used was not outlined.

136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159

160
161
162
163
164
165
166
167
168
169
170
171
172
173
174

Table 2

Follow-up studies on patients with different stages of non-invasive urothelial carcinoma of the prostatic urethra treated with or without TURP before BCG treatment. Tumor progression was defined by the development of muscle infiltration, metastatic disease or the occurrence of superficial disease refractory to transurethral resection and intravesical BCG necessitating a change in therapy. Survival data in relation to the median follow-up time of the cited study. Abbreviations: TURP = transurethral resection of the prostate; TURP = transurethral resection of the prostate; BCG = Bacille Calmette Guérin; DFS = disease free survival; DSS = disease specific survival (i.e. disease specific cause was including urothelial cancer in the bladder or the upper urinary tract); PFS = progression free survival; N.A. = not available/not applicable

Reference	+/- TURP before BCG	Total number of prostatic urethral cancer Patients treated with BCG	Primary prostatic urethral cancer cases n/N (%)	Mucosal prostatic urethral cancer (Tis pu) n/N (%)	Ductal prostatic urethral cancer (Tis pd) n/N (%)	Stromal prostatic urethral cancer (T1) n/N (%)	BCG regimen	BCG strain	Median follow up in months (range)	Bladder cancer DFS n/N (%)	Bladder cancer PFS n/N (%)	Prostatic urethral cancer DFS n/N (%)	Prostatic urethral cancer PFS n/N (%)	Prostatic urethral+bladder cancer DFS n/N (%)	Prostatic urethral+bladder cancer PFS n/N (%)	DSS n/N (%)
Gofrit et al. (2008) [10]	+	20	16/20 (80%)	N.A.	N.A.	N.A.	1/week for 6 weeks	Connaught	52.5 (N.A.)	7/20 (35%)	17/20 (85%)	18/20 (90%)	20/20 (100%)	6/20 (30%)	15/20 (75%)	14/20 (70%)
Ovesen et al. (1993) [11]	+	10	10/10 (100%)	N.A.	N.A.	N.A.	1/week for 6 weeks in 8/10, for 12 weeks in 2/10	Danish	26 (3–68)	N.A.	N.A.	8/10 (80%)	9/10 (90%)	N.A.	N.A.	10/10 (100%)
Bretton et al. (1989) [12]	+	23	23/23 (100%)	19/23 (83%)	4/23 (17%)	0/23 (0%)	1/week for 6 weeks	Pasteur	51.6 (6–105)	N.A.	13/23 (57%)	23/23 (100%)	23/23 (100%)	13/23 (57%)	13/23 (57%)	19/22 (91%)
Orihuela et al. (1989) [13]	-8/15 +7/15	15	N.A.	N.A.	N.A.	N.A.	1/week for 6 weeks	Pasteur	N.A. (18–51)	14/15 (93%)	15/15 (100%)	13/15 (86%)	14/15 (93%)	13/15 (86%)	14/15 (93%)	15/15 (100%)
Palou et al. (1996) [14]	-17/18 +1/18	18	15/18 (79%)	N.A.	N.A.	N.A.	1/week for 6 weeks	Connaught	31.1 (7–57)	17/18 (94%)	17/18 (94%)	13/18 (68%)	17/18 (94%)	15/18 (79%)	15/18 (79%)	17/18 (94%)
Taylor et al. (2007) [15]		28	28/28 (100%)	N.A.	N.A.	N.A.	1/week for 6 weeks in most cases	N.A.	90 (N.A.)	N.A.	20/28 (71%)	25/28 (89%)	25/28 (89%)	N.A.	19/28 (68%)	25/28 (89%)
Palou et al. (2006) [16]		11	6/11 (54%)	11/11 (100%)	0/11 (0%)	0/11 (0%)	1/week for 6 weeks in 10/11, for 12 weeks in 1/11	Connaught	40 (8–157)	7/11 (63%)	10/11 (90%)	9/11 (81%)	9/11 (81%)	6/11 (54%)	8/11 (72%)	11/11 (100%)
Canda et al. (2004) [17]		12	12/12 (100%)	7/12 (58%)	N.A.	5/12 (42%)	1/week for 6 weeks in 10/12, for 12 weeks in 2/12	Connaught	62.5 (24–110)	6/12 (50%)	8/12 (66%)	10/12 (83%)	11/12 (91%)	5/12 (42%)	7/12 (58%)	8/12 (66%)
Schellhammer et al. (1995) [18]	-	17	8/17 (47%)	N.A.	N.A.	N.A.	1/week for 6 weeks	Pasteur	64.0 (29–90)	9/17 (52%)	14/17 (82%)	11/17 (65%)	14/17 (82%)	8/17 (47%)	10/17 (59%)	16/17 (94%)
Hillyard et al. (1988) [19]	-	8	8/8 (100%)	N.A.	N.A.	N.A.	1/week for 6 weeks	Pasteur	22.3 (15–52)	6/8 (75%)	7/8 (86%)	5/8 (63%)	6/8 (75%)	N.A.	5/8 (63%)	8/8 (100%)

175 Only in one study did 5 of 12 patients have stage T1
 176 urethral carcinoma of the prostate [17]. In this study,
 177 the DFS (83%) and the PFS (91%) in the prostate
 178 were within the range of the remaining nine studies.
 179 However, the overall DSS was only 66% compared
 180 to an overall DSS of 70–100% in the other studies. It
 181 could not be completely excluded that these stromal
 182 invasive cases reduced the survival of the patients
 183 in this study compared to the other studies. Without
 184 access to the full data, we could not perform statistical
 185 analyzes.

186 *The role of TURP before BCG in prostatic* 187 *urethral carcinoma*

188 Prospective randomized clinical trials or even
 189 directly applicable clinical studies of good quality
 190 addressing the question if performing TURP prior
 191 to BCG improves DSS or lowers morbidity with-
 192 out compromising DSS in PUC were not available.
 193 This issue was exclusively addressed by retrospec-
 194 tive cohort studies. Unfortunately, the authors of the
 195 selected studies did not use the same format in pre-
 196 senting their data. Some authors only described the
 197 results in the text, others used tables with limited
 198 information on e.g. follow-up, DFS, PFS or DSS.

199 A total of ten articles found that examined the
 200 effectiveness of BCG in urethral carcinoma in the
 201 prostatic urethra with or without TURP prior to
 202 BCG treatment (Table 2). The studies by Gofrit [10],
 203 Ovesen [11] and Bretton [12] only included patients
 204 who had been treated with TURP before BCG. Ori-
 205 huela [13] and Palou [14] included both patients with
 206 and without TURP prior to BCG treatment.

207 Although an appropriate comparative statistical
 208 analysis is not suitable for these heterogeneously
 209 designed studies, there appears to be a pattern that
 210 suggests slightly better local response with higher
 211 DFS (80–100% vs. 63–89%) and PFS (90–100%
 212 vs. 75–94%) in the prostatic urethra in the studies
 213 in which patients were treated with TURP prior to
 214 BCG treatment compared to the studies in which no
 215 TURP was performed. However, this difference in
 216 recurrence and progression in the prostate affected
 217 neither the total PFS (57–75% vs. 58–93%) nor the
 218 DSS (70–100% vs. 66–100%).

219 **DISCUSSION**

220 Intravesical BCG treatment with or without TURP
 221 prior BCG induction therapy was retrospectively
 222 evaluated in non-invasive stages of PUC. Upon

223 reviewing the published literature, it becomes clear
 224 that the incidence of primary PUC is likely to be
 225 underestimated: Almost all cases of primary PUC
 226 were detected during the workup of primary blad-
 227 der cancer or UC of the UUT. This impression is
 228 supported by the study by Giannarini et al, which
 229 indicated that an undiagnosed PUC could often be the
 230 cause of BCG failure. [21]. In this study, bladder CIS
 231 was an independent risk factor for BCG failure due to
 232 recurrence in the prostatic urethra or UUT. Since not
 233 every patient in this study had a TUR biopsy before
 234 starting BCG therapy, it was discussed that the actual
 235 incidence of PUC is difficult to estimate and is likely
 236 to be underestimated.

237 The most accurate method of detecting PUC during
 238 the primary workup of suspected urothelial cancer in
 239 the prostate is to perform a TUR biopsy, which has
 240 been shown to be superior to prostate needle biopsy
 241 and fine needle aspiration of the prostate [22]. Donat
 242 and Herr suggested a resectoscope loop biopsy of
 243 the prostatic urethra between the 5 and 7 o'clock
 244 positions from the bladder neck and distally around
 245 the verumontanum in patients with suspected PUC
 246 [23]. However, not only should suspicious lesions in
 247 the prostatic urethra be biopsied: Palou showed that
 248 almost 12% of patients with T1G3 bladder cancer
 249 had carcinoma in situ of the prostatic urethra on a
 250 resection biopsy [24]. In several series for the eval-
 251 uation of cystectomy specimens, involvement of the
 252 prostatic urethra was found in 15–48% [25–27]. Both
 253 CIS in the bladder and multifocal tumor growth were
 254 associated with a higher risk of involvement of carci-
 255 noma of the prostatic urethra. With the data available,
 256 it seems reasonable to recommend resection biop-
 257 sies in patients with positive urinary cytology of
 258 unknown origin, if CIS in the bladder or upper uri-
 259 nary tract is suspected, if bladder cancer is multifocal
 260 or occurs around the bladder neck, and when radical
 261 cystectomy is pending especially if chemotherapy is
 262 planned before surgery [6]. The AUA/SUO guide-
 263 lines so far only recommend prostate loop biopsies
 264 in cases with a suspicious lesion in the prostate and
 265 in patients with a history of NMIBC with normal
 266 cystoscopy and positive cytology [5].

267 It was discussed that cases of PUC with stage T1
 268 and Tis pd might have a worse prognosis than tumors
 269 with stage Tis pu [28]. There are no prospective
 270 studies on the conservative treatment of non-invasive
 271 PUC in its various tumor stages, which is why this
 272 hypothesis has not yet been confirmed. In seven of
 273 the ten available studies in this review, no TNM stag-
 274 ing of PUC was carried out; and only one of the

ten studies [17] included patients with stage T1 urethral carcinoma ($n=5$). Accurate staging of tumors is the basic requirement for evaluating the prognostic effects of the extent of TURP prior to BCG treatment in malignant diseases of the urothelium of the prostatic urethra and ducts. This underscores the importance of improving the use of the current TNM staging system for PUC in the future.

According to the EAU-guidelines, a urethra-sparing approach with TURP and intravesical-BCG is indicated in patients with non-invasive UC or carcinoma in situ of the prostatic urethra and ducts (level of evidence 3, grade of recommendation C) [6]. In the available studies, of all BCG-treated patients who were not treated with TURP, almost 2/3 of the patients had no tumor recurrence in the prostate and at least 3/4 had no tumor progression in the prostate. In the available studies examining the use of TURP before BCG therapy, about 4/5 of the patients had no recurrence and 9/10 no tumor progression in the prostate. However, these improved local response rates after TURP had no influence on the overall response (PFS, DSS) in these cohorts. This finding supports the rationale for performing TURP prior to BCG treatment for non-invasive PUC, namely improving effectiveness of BCG treatment, at least with regard to the local reaction in the prostate, without necessarily improving the overall prognosis. Prospective studies need to confirm these results. In particular, the treatment of the rare cases of stage T1 PUC should be carefully studied.

Systemic side effects are seen in approximately 30% of patients treated with BCG for bladder cancer. Local symptoms of BCG instillations are cystitis, epididymitis or granulomatous prostatitis with dysuria, urination problems and hematuria. These symptoms occur in about 60% of BCG treated bladder cancer patients [29]. Approximately 70% of bladder cancer patients treated with BCG have both systemic and local side effects. In 16–22% of treated patients BCG treatment is discontinued because of toxicity [30]. Since almost all patients with PUC also have concomitant bladder cancer, specific side effects of BCG in the treatment of PUC could not be determined and cannot be found in the literature. It was therefore impossible to derive specific side effects of BCG therapy for PUC, which is a limitation of the study.

Apart from that, several other limitations of the current study have to be mentioned: Different definitions of progression were used in the cited studies. A comparative statistical analysis was not possible. No

data on the natural history of the various non-invasive stages of PUC were available for comparison. Most studies only offered a BCG induction course. It cannot be ruled out that maintenance therapy for non-invasive PUC could have an improved prognosis. In addition, different BCG strains were used (Connaught and Pasteur). In one study [11] the rare Danish BCG strain was used. It is uncertain how this affected the results. Another problem is inconsistent follow-up of tumor status in the prostatic urethra: In some studies, cold cup biopsies were used more or less routinely, and in other studies, TUR biopsies of the prostate were taken only when there were visible or suspicious lesions. These differences in follow up schemes and methods might have affected differences of the outcomes after BCG therapy significantly.

CONCLUSIONS

Taken together, the use of TUR biopsies of the prostatic urethra in appropriate cases during the primary work-up for suspected urothelial cancer, as well as the use of the current TNM classification for PUC, need to be improved. BCG therapy for non-stromal invasive stages of PUC show a good local response. Local response rates are further improved by a TURP, although the overall prognosis does not seem to be affected. Further evidence for BCG treatment in the rare cases of stromal invasive PUC is needed. Specific side effects of BCG treatment for PUC, either with or without TURP before treatment, are not reported.

ACKNOWLEDGMENTS

The authors have no acknowledgments

FUNDING

The authors report no funding.

AUTHORS CONTRIBUTIONS

All authors contributed to the conception, sampling and interpretation of data, and writing of the article.

ETHICAL CONSIDERATIONS

This study, as a literature review, is exempt from any requirements for Institutional Review Board approval. No human or animal research was involved in the elaboration of this manuscript.

CONFLICT OF INTEREST

Georgios Gakis receives honoraria for advisory board activities: MSD, Astellas, Ferring, BMS, Bayer, Leo Pharma, medac, Merck, and he receives speaker honoraria/travel grants: Roche, MSD, IPSEN, Erbe, medac. Philippe E Spiess is the vice chair of the NCCN bladder and penile cancer panel; he is as well the president of the Global Society of Rare GU Tumors. Joan Palou Redorta reports trial participation: Cepheid, Presurgery, Ipsen, Combat, Storz, Archer, IDL Biotech, Palex and receives honoraria or consultation fees from Combat BRS, Olympus, Sanofi, Cepheid, Ipsen, Fidia Farmaceutici, and fees for Advisory/Scientific Board activities from: Pfizer, Astra Zeneca, Merck, Janssen, Ferring. Oliver Patschan and George N Thalmann have no conflict of interest to declare.

REFERENCES

- [1] Gatta G, van der Zwan JM, Casali PG, et al. Rare cancers are not so rare: the rare cancer burden in Europe. *Eur J Cancer*. 2011;47(17):2493-511. doi:10.1016/j.ejca.2011.08.008
- [2] Visser O, Adolffson J, Rossi S, et al. Incidence and survival of rare urogenital cancers in Europe. *Eur J Cancer*. 2012;48(4):456-64. doi:10.1016/j.ejca.2011.10.031
- [3] Swartz MA, Porter MP, Lin DW, Weiss NS. Incidence of primary urethral carcinoma in the United States. *Urology*. 2006;68(6):1164-8. doi:10.1016/j.urology.2006.08.1057
- [4] Aleksic I, Rais-Bahrami S, Daugherty M, Agarwal PK, Vourganti S, Bratslavsky G. Primary urethral carcinoma: A Surveillance, Epidemiology, and End Results data analysis identifying predictors of cancer-specific survival. *Urol Ann*. 2018;10(2):170-4. doi:10.4103/UA.UA.136.17
- [5] Sui W, RoyChoudhury A, Wenske S, Decastro GJ, McKiernan JM, Anderson CB. Outcomes and Prognostic Factors of Primary Urethral Cancer. *Urology*. 2017;100:180-6. doi:10.1016/j.urology.2016.09.042
- [6] Gakis G, Bruins HM, Cathomas R, et al. European Association of Urology Guidelines on Primary Urethral Carcinoma-2020 Update. *Eur Urol Oncol*. 2020;3(4):424-32. doi:10.1016/j.euo.2020.06.003
- [7] Rabbani F. Prognostic factors in male urethral cancer. *Cancer*. 2011;117(11):2426-34. doi:10.1002/cncr.25787
- [8] Dalbagni G, Zhang ZF, Lacombe L, Herr HW. Male urethral carcinoma: analysis of treatment outcome. *Urology*. 1999;53(6):1126-32. doi:10.1016/s0090-4295(98)00659-1
- [9] Sylvester RJ, van der Meijden AP, Lamm DL. Intravesical bacillus Calmette-Guerin reduces the risk of progression in patients with superficial bladder cancer: a meta-analysis of the published results of randomized clinical trials. *J Urol*. 2002;168(5):1964-70. doi:10.1097/01.ju.0000034450.80198.1c
- [10] Gofrit ON, Pode D, Pizov G, Zorn KC, Katz R, Shapiro A. Prostatic urothelial carcinoma: is transurethral prostatectomy necessary before bacillus Calmette-Guérin immunotherapy? *BJU Int*. 2009;103(7):905-8. doi:10.1111/j.1464-410X.2008.08210.x
- [11] Ovesen H, Poulsen AL, Steven K. Intravesical Bacillus Calmette-Guérin with the Danish strain for treatment of carcinoma in situ of the bladder. *Br J Urol*. 1993;72(5 Pt 2):744-8. doi:10.1111/j.1464-410x.1993.tb16260.x
- [12] Bretton PR, Herr HW, Whitmore WF Jr, et al. Intravesical bacillus Calmette-Guerin therapy for in situ transitional cell carcinoma involving the prostatic urethra. *J Urol*. 1989;141(4):853-6. doi:10.1016/s0022-5347(17)41031-7
- [13] Orihuela E, Herr HW, Whitmore WF Jr. Conservative treatment of superficial transitional cell carcinoma of prostatic urethra with intravesical BCG. *Urology*. 1989;34(5):231-7. doi:10.1016/0090-4295(89)90314-2
- [14] Palou J, Xavier B, Laguna P, Montleó M, Vicente J. In situ transitional cell carcinoma involvement of prostatic urethra: bacillus Calmette-Guérin therapy without previous transurethral resection of the prostate. *Urology*. 1996;47(4):482-4. doi:10.1016/S0090-4295(99)80481-6
- [15] Taylor JH, Davis J, Schellhammer P. Long-term follow-up of intravesical bacillus Calmette-Guérin treatment for superficial transitional-cell carcinoma of the bladder involving the prostatic urethra. *Clin Genitourin Cancer*. 2007;5(6):386-9. doi:10.3816/CGC.2007.n.021
- [16] Palou Redorta J, Schatteman P, Huguet Pérez J, et al. Intravesical instillations with bacillus calmette-guérin for the treatment of carcinoma in situ involving prostatic ducts. *Eur Urol*. 2006;49(5):834-8. doi:10.1016/j.eururo.2005.12.019
- [17] Canda AE, Tuzel E, Mungan MU, Yorukoglu K, Kirkali Z. Conservative management of mucosal prostatic urethral involvement in patients with superficial transitional cell carcinoma of the bladder. *Eur Urol*. 2004;45(4):465-70. doi:10.1016/j.eururo.2003.12.014
- [18] Schellhammer PF, Ladaga LE, Moriarty RP. Intravesical bacillus Calmette-Guerin for the treatment of superficial transitional cell carcinoma of the prostatic urethra in association with carcinoma of the bladder. *J Urol*. 1995;153(1):53-6. doi:10.1097/00005392-199501000-00020
- [19] Hillyard RW Jr, Ladaga L, Schellhammer PF. Superficial transitional cell carcinoma of the bladder associated with mucosal involvement of the prostatic urethra: results of treatment with intravesical bacillus Calmette-Guerin. *J Urol*. 1988;139(2):290-3. doi:10.1016/s0022-5347(17)42389-5
- [20] Brierley JD, et al. TNM classification of malignant tumors. UICC International Union Against Cancer. Wiley/Blackwell; 2017. pp. 208-9.
- [21] Giannarini G, Birkhäuser FD, Recker F, Thalmann GN, Studer UE. Bacillus Calmette-Guérin failure in patients with non-muscle-invasive urothelial carcinoma of the bladder may be due to the urologist's failure to detect urothelial carcinoma of the upper urinary tract and urethra. *Eur Urol*. 2014;65(4):825-31. doi:10.1016/j.eururo.2013.09.049
- [22] Wood DP Jr, Montie JE, Pontes JE, Levin HS. Identification of transitional cell carcinoma of the prostate in bladder cancer patients: a prospective study. *J Urol*. 1989;142(1):83-5. doi:10.1016/s0022-5347(17)38667-6
- [23] Donat SM, Wei DC, McGuire MS, Herr HW. The efficacy of transurethral biopsy for predicting the long-term clinical impact of prostatic invasive bladder cancer. *J Urol*. 2001;165(5):1580-4.
- [24] Palou J, Sylvester RJ, Faba OR, et al. Female gender and carcinoma in situ in the prostatic urethra are prognostic factors for recurrence, progression, and disease-specific mortality in T1G3 bladder cancer patients treated with bacillus Calmette-Guérin. *Eur Urol*. 2012;62(1):118-25. doi:10.1016/j.eururo.2011.10.029

- 488 [25] Revelo MP, Cookson MS, Chang SS, Shook MF, Smith
489 JA Jr, Shappell SB. Incidence and location of prostate
490 and urothelial carcinoma in prostates from cystoprostate-
491 ctomies: implications for possible apical sparing surgery.
492 J Urol. 2004;171(2 Pt 1):646-51. doi:10.1097/01.ju.0000
493 107380.40481.bc
- 494 [26] Esrig D, Freeman JA, Elmajian DA, et al. Transitional
495 cell carcinoma involving the prostate with a proposed
496 staging classification for stromal invasion. J Urol. 1996;
497 156(3):1071-6.
- 498 [27] Nixon RG, Chang SS, Lafleur BJ, Smith JA JA, Cook-
499 son MS. Carcinoma in situ and tumor multifocality predict
500 the risk of prostatic urethral involvement at radical cystec-
501 tomy in men with transitional cell carcinoma of the bladder.
502 J Urol. 2002;167(2 Pt 1):502-5. doi:10.1097/00005392-
503 200202000-00012
- 504 [28] Njinou Ngninkeu B, Lorge F, Moulin P, Jamart J, Van Cangh
505 PJ. Transitional cell carcinoma involving the prostate:
506 a clinicopathological retrospective study of 76 cases. J
507 Urol. 2003;169(1):149-52. doi:10.1097/01.ju.0000042810.
508 43380.36
- [29] Brausi M, Oddens J, Sylvester R, et al. Side effects
509 of Bacillus Calmette-Guárin (BCG) in the treatment of
510 intermediate- and high-risk Ta, T1 papillary carcinoma of
511 the bladder: results of the EORTC genito-urinary cancers
512 group randomised phase 3 study comparing one-third dose
513 with full dose and 1 year with 3 years of maintenance BCG.
514 Eur Urol. 2014;65(1):69-76. doi:10.1016/j.eururo.2013.
515 07.021
- [30] Oddens JR, Sylvester RJ, Brausi MA, et al. Increasing age
517 is not associated with toxicity leading to discontinuation of
518 treatment in patients with urothelial non-muscle-invasive
519 bladder cancer randomised to receive 3 years of main-
520 tenance bacille Calmette-Guárin: results from European
521 Organisation for Research and Treatment of Cancer Genito-
522 Urinary Group study 30911. BJU Int. 2016;118(3):423-8.
523 doi:10.1111/bju.13474
524