

Erratum

Ex vivo neutralization of HIV-1 quasi-species by a broadly reactive humanized monoclonal antibody KD-247

Shuzo Matsushita^{a,*}, Soichiro Takahama^a, Junji Shibata^a, Tetsuya Kimura^a, Koichi Shiozaki^b, Yasuyuki Eda^b, Atsushi Koito^a, Toshio Murakami^b and Kazuhisa Yoshimura^a

[*Human Antibodies* 14 (3,4), 2005, pp. 81–88]

Figure 4 from this article has been omitted. Instead, Fig. 2 has been published twice.

Figure 4 on page 86 should be replaced with the correct Fig. 4 as shown below:




Fig. 4. Neutralization *ex vivo* of HIV-1 quasi-species by MAb KD-247. The culture supernatants were collected at day 11, 15 and 7 (CT) for patient 1 (A) and at day 10, 20 and 9 (CT) for patient 2 (B). The CD8-depleted PBMC samples from HIV-infected patients were cultured in the presence of KD-247 (60 $\mu\text{g}/\text{mL}$; gray, 120 $\mu\text{g}/\text{mL}$; diagonal up hatch, 240 $\mu\text{g}/\text{mL}$; open) or control MAb (240 $\mu\text{g}/\text{mL}$ of C β 1; diagonal down hatch), or in the absence of MAb (closed). Neutralization was determined by detecting the concentration of p24 antigen in the supernatant of each culture following the assay protocol described in Fig. 3.