

Hypothesis

Alzheimer's Disease as the Product of a Progressive Energy Deficiency Syndrome in the Central Nervous System: The Neuroenergetic Hypothesis

Edward R. Blonz*

Department of Clinical Pharmacy, University of California, San Francisco, CA, USA

Accepted 10 August 2017

Abstract. The decreased availability of metabolizable energy resources in the central nervous system is hypothesized to be a key factor in the pathogenesis of Alzheimer's disease. More specifically, the age-related decline in the ability of glucose to cross the blood-brain barrier creates a metabolic stress that shifts the normal, benign processing of amyloid- β protein precursor toward pathways associated with the production of amyloid- β plaques and tau-containing neurofibrillary tangles that are characteristic of the disease. The neuroenergetic hypothesis provides insight into the etiology of Alzheimer's disease and illuminates new approaches for diagnosis, monitoring, and treatment.

Keywords: Alzheimer's disease, alpha-secretase, amyloid- β , amyloid- β protein precursor, beta-hydroxy butyrate, beta-secretase, blood-brain barrier, glucose, gamma-secretase, GLUT1, neuroenergetic hypothesis, tau, type-3 diabetes

INTRODUCTION

Alzheimer's disease (AD) is the most common form of dementia in developed countries. It is characterized by progressive dysfunction of memory and cognition with a long prodromal period. Pathologic accumulations of amyloid- β (A β) plaques and tau-containing neurofibrillary tangles (tau) are found, but an understanding of their cause remains a work in progress. AD represents an ever-growing public health problem that places unique burdens on families, care providers, and the healthcare system in general. The 2017 report from the Alzheimer's Association notes that more than five million people are living with AD in the United States. Given the absence of effective interventions to slow progress, it

is projected to affect 106 million people by 2050 [1]. It is the only disease in the top ten causes of death that cannot be prevented, cured, or even slowed [2].

Hypotheses have attempted to explain aspects of AD, with the most basic being the amyloid cascade hypothesis which associates AD with its characteristic extracellular accumulation of A β and tau [3]. The mitochondrial cascade hypothesis expanded the discussion to include effects of inherited and environmental factors on mitochondrial function, correlating these with A β and tau formation and the trajectory of AD pathogenesis [4, 5]. Hypotheses have suggested roles of various metals, including copper, iron, zinc, aluminum, and calcium [6–10]. One hypothesis has advanced the relationship between AD and neuroinflammation [11], and there has been a proposal that AD be considered as “type-3 diabetes” [12]. While research has advanced and expanded our knowledge of the pathology of AD, an understanding of its

*Correspondence to: Edward R. Blonz, PhD, 139 Purdue Avenue, Kensington, CA 94708, USA. E-mail: er@blonz.com.

etiology has yet to be fully elucidated, and prevention and control measures remain beyond our grasp.

This paper reviews the evidence and suggests that the status of available metabolizable energy resources in the central nervous system (CNS) is a central determinant of AD pathogenesis. The neuroenergetic hypothesis posits that the chronic, progressive decline in the ability of glucose to cross the blood-brain barrier (BBB)—a normal event of aging—contributes to an energy-deficiency stress in the CNS. In reaction to this glucose/energy-deficiency stress, there is a shift from pathways associated with the normally benign metabolism of amyloid- β protein precursor (A β PP), to a pathway associated with A β /tau production and AD progression. Considering AD from this perspective provides insight into its pathophysiology while embracing and clarifying the specific biochemical roles of previously-identified risk factors and correlates, and can thus inform regarding potential approaches for diagnosis, monitoring, treatment, and prevention.

NEUROENERGETICS AND THE ALZHEIMER'S DISEASE PROCESS

Glucose is the brain's primary source of energy. To enter the CNS from the general circulation, however, glucose must traverse the protective BBB. The primary conduit through the BBB is GLUT1 (SLC2A1), a membrane-bound glucose transporter [13]. Even in the absence of identified pathologies, the level of glucose available to the CNS deteriorates with age [14]. It then follows that available metabolizable energy from glucose would be further reduced by factors affecting its net ability to traverse the BBB and be utilized by the CNS.

To illustrate, the $\epsilon 4$ variant of the APOE gene is a known risk factor for AD. Transgenic mice carrying the APOE $\epsilon 4$ allele have a 29% reduction in glucose transport through the BBB compared to those having the protective APOE $\epsilon 2$ allele [15]. The higher risk and lower age of AD onset in human APOE $\epsilon 4$ carriers may, therefore, be the result of their lower baseline of glucose availability in the CNS, this being present for decades before any perceived or reported impacts on cognition [16].

Consistent with the neuroenergetic hypothesis, reduced GLUT1 activity has been reported to correlate with an acceleration of AD symptomatology [17]. Diseases involving glucose dysregulation, such as diabetes mellitus, are associated with an increased

AD risk [18]. Reduced availability of glucose in the brain may contribute to obesity, this condition representing a risk factor for AD even if independent of diabetes [19].

Disease processes affecting vascularity and endothelial function, including hypertension, have been linked with increased AD risk [20]. Increased risk of AD has also been associated with viral conditions such as infections with members of the herpes family, including Epstein-Barr virus, human herpes virus 6 and cytomegalovirus; all reported to alter GLUT1 activity through a negative effect on its integral enzyme, protein kinase C (PKC) [21]. Conversely, factors reported to increase PKC activity have been associated with cognitive improvements in memory-impaired individuals [22], such evidence having led to PKCs being referred to as a "cognitive kinase" [23].

Events affecting systemic blood sugar levels can play a role if they contribute to a decreased ability of adequate glucose to cross the BBB. In particular, hyperglycemia has been linked to the creation of advanced glycation end products that can impact BBB functionality and contribute to the risk of AD [24]. Dietary fats, given their role in BBB structure, must also be considered. The Chicago Health and Aging Project (CHAP) reported an increased risk of AD with higher intakes of saturated fats and trans fatty acids, while monounsaturated fats were associated with a decreased risk [25].

Metal toxicity from chronic aluminum exposure is linked with AD [9, 26] and is related to decreases in PKC activity [27]. Ethyl alcohol, which readily crosses the BBB, can affect glycolysis the main glucose utilization pathway, and alcohol, when consumed excessively, is associated with an increased risk of AD [28]. Repeated or severe episodes of systemic hypoglycemia have also been connected with dementia [29]. More directly, cerebral hypoglycemia can induce hyperphosphorylation of the tau protein, possibly contributing to the increased risk of neurofibrillary tangles associated with AD [30].

AMYLOID- β AND ITS PROTEIN PRECURSOR

The initial step in the metabolism of A β PP is critical in AD pathogenesis as it determines whether or not A β will be produced as a byproduct [31]. When A β PP is first acted upon by alpha-secretase (ADAM10) [32], the intermediate product is a large




Fig. 1. The generation of products from the amyloid- β protein precursor (A β PP). During periods of central nervous system (CNS) energy adequacy, A β PP will get cleaved along a non-amyloidogenic pathway by the action of α -secretase to produce sAPP α , which, after the action of γ -secretase, produces p3 peptides which have an ability to exit the CNS through the blood-brain barrier (BBB). During periods of CNS energy inadequacy, A β PP gets cleaved along an amyloidogenic pathway by the action of β -secretase to product sAPP β , which, after the action of γ -secretase, produces intact amyloid- β (A β) peptides that cannot effectively exit the CNS through the BBB, and are associated with the pathogenesis of Alzheimer's disease. Adapted from [32] (CC BY 4.0).

ectodomain referred to as sAPP α , a product in which A β is no longer intact. When sAPP α is subsequently acted upon by γ -secretase, a soluble, non-AD pathogenic p3 peptide is produced that can exit the CNS through the BBB. If, however, A β PP is first acted upon by β -secretase (BACE1) [32], the intermediate product is the ectodomain referred to as sAPP β that contains the intact A β . When sAPP β is subsequently acted upon by γ -secretase, A β gets liberated, this being a substance that cannot effectively exit the CNS through the BBB and one associated with AD's pathology as well as its pathophysiology [33]. Factors, therefore, that promote A β PP metabolism by ADAM10 should be considered as AD benign, while those factors promoting metabolism via BACE1 should be considered as causing pathology characteristic of AD. Neuroenergetic status is hypothesized to be a dynamic determinant of the A β PP degradative pathway; inadequate CNS glucose favoring the A β PP \rightarrow A β pathway (see Fig. 1).

An understanding of a physiological basis for significant production of A β in normal CNS functionality has been elusive. Evidence now suggests the possibility of an antimicrobial role [34, 35]. It should be considered that one hallmark of viral and

microbial success is an ability to outcompete their host for essential resources. These entities commonly rely on glucose for energy, thus potentially contributing to the risk of hypoglycemic stress if affecting glucose adequacy in the CNS. An element of the response to a microbially-induced CNS hypoglycemia could be the shunting of A β PP processing toward BACE1 to produce antimicrobial A β , which could ironically be neuroprotective, at least in an acute setting. Consistent with this premise, chronic viral activity, such as that from the herpes simplex virus, has indeed been associated with an upregulation of BACE1 and an increased risk of AD [36]. Inadequate CNS glucose, thus, can act as a cellular stressor contributing to increased BACE1 activity [37].

PKC status plays an indirect role through its association with GLUT1, and the ability of glucose to traverse the BBB. Decreases in PKC activity are linked with a shift from the AD-benign ADAM10 to the amyloidogenic BACE1 [38]. Conversely increased PKC activity could help reduce the risk of hypoglycemic stress in the CNS and help maintain, or even shift, A β PP degradation in favor of ADAM10. This supports a logic behind the present use of PKC

activation as a treatment for AD [22]. An existing energy deficiency in the CNS can also drive utilization of myelin lipids for conversion into ketones as an alternate source of metabolizable energy [39], such a breakdown also reportedly associated with the AD process [40]. Factors known or suspected to be associated with the AD process would thus merit study for potential impacts on neuroenergetic status.

It is also noteworthy that a hypoglycemic environment in the CNS has been reported to create oxidative stress with an increased presence of reactive oxygen species [41]. This could help explain key aspects of hypotheses related to metal dyshomeostasis and AD pathogenesis [42], shifting that putative role from casual to opportunistic, that is, the product of events in the developing energy-deficient CNS environment [43, 44].

Consideration of AD as a “type-3” diabetes is of interest, given that bulk delivery of glucose to the CNS through the BBB is not an insulin-dependent phenomenon [45]. Systemic hyperinsulinemia and insulin resistance reportedly correlate with various factors consistent with AD pathogenesis [46, 47], yet improvements in cognition have been reported with the administration of intranasal insulin [48], which bypasses the BBB and gains direct access to the CNS through the olfactory bulb. This phenomenon may be the result of an effect on local neurotransmitter release and synaptic plasticity rather than an increase in net glucose transport through the BBB via the hypothesized impact of neuroenergetic status on the AD process [49, 50].

The inflammatory hypothesis suggests that CNS inflammation helps cause, or at least promote, AD. The concept is supported, in part, by reports of AD-sparing effects observed in individuals consuming anti-inflammatory drugs [51]. Inflammation is an integral part of the body’s first line of defense against invasive pathogens. mRNA levels for peroxisome proliferator-activator receptor gamma (PPAR γ), a hormone-regulating receptor that plays a role in the body’s anti-inflammatory response, tend to be elevated in AD [52]. The chronic, progressive decline in the ability of glucose to cross the BBB could provoke an inflammatory response if responded to as suspected pathogenic activity in the CNS, given the sufficiency of energy elsewhere in the body. Although a local phenomenon, neuroinflammation must also be acknowledged as a glucose-consuming event with the potential to further drain this energy resource in the CNS [53]. The reduction of energy-consuming local inflammation could, in theory, represent a mechanism

behind AD-protective effects from anti-inflammatory activity, but it is unclear whether, or to what degree, it could function as a block to overall neuroenergetic inadequacy and AD progression.

ALTERNATE SOURCES OF METABOLIZABLE ENERGY IN THE CNS

Another source of metabolizable energy in the CNS are ketone bodies that include D- β -hydroxybutyrate (β -HB) and acetoacetate, these substances being able to traverse the BBB independent of GLUT1 functionality. β -HB and acetoacetate are made and utilized in the human body during periods of starvation, diabetic ketoacidosis, or when there is a prolonged lack of digestible carbohydrates in the diet [54]. β -HB is also produced when unabsorbed food components, often polysaccharide in structure, are metabolized by microorganisms in the human microbiome to produce butyric acid, a short-chain fatty acid that appears in the blood as β -HB after passage through the liver [55]. By way of its contribution to available β -HB, a high-fiber diet has been suggested as an asset to brain health [56, 57]. Butyric acid is also present in milk fat, representing about 3% of the lipids in butter [58]. Nutrients or agents contributing to increased circulating levels of β -HB could, in effect, serve as alternative substrates for CNS energy metabolism, helping to explain reports of cognitive improvements in AD populations given β -HB directly from external sources [59–61]. Caprylic acid, an eight-carbon fatty acid comprising approximately 8% of the lipids in coconut and 4% of the lipids in palm kernel, also gets metabolized to β -HB in the liver and has been reported to produce ketonemia in a case of AD, with improvements in cognition tracking with the plasma ketone level [62]. Triheptanoin, already in use clinically to treat inherited metabolic disorders, is another source of metabolic energy for the CNS not reliant on GLUT1 that would merit study for use as an AD intervention [62]. In animal models, the induction of ketosis in dogs reportedly reduced the accumulation of A β /tau [63]. Using the mouse model of AD, ketones were reported to reduce amyloid in the brain and improve cognition [64].

EPIDEMIOLOGICAL CONSIDERATIONS

A theme emerges when epidemiological studies examining lifestyle factors are viewed through the lens of the neuroenergetic hypothesis. Lower risks of

AD have been reported in cultures with a traditional high-fiber cuisine, such as those in Sub-Saharan Africa, as well individuals adhering to the Mediterranean, or Mediterranean-DASH Diet Intervention for Neurodegenerative Delay (MIND) diets [65–68]. Exercise contributes to vascular health, which has a positive impact on BBB in aging [69]. Lactate, created during exercise, is another fuel that can traverse the BBB via monocarboxylate transporters that are independent of GLUT1 and the BBB's age-related changes [70, 71]. Consistent with this premise, pathologies associated with negative effects on vascular health and BBB performance are associated with an increased AD risk, including obesity [72], oxidative stress [73], hypertension [74], hyperinsulinemia [75], metabolic syndrome [73], and insulin resistance and glycemic disorders [76].

CONCLUSION

The neuroenergetic hypothesis seeks to explain the process of AD as resulting from the chronic decline in available metabolizable energy resources, mainly glucose, in the CNS. The range of metabolic consequences of this age-related, progressive decline is consistent with existing hypotheses and epidemiological observations. Viewing AD from this perspective can provide new insights and approaches to diagnosis, treatment, and risk mitigation. In order to fully test this hypothesis, CNS energy resources consistent with normal health will need to be assessed at various ages and developmental stages. There will also need to be a better understanding of the role of GLUT1, PKC, BBB, and any other functional element known to affect metabolizable energy resources in the CNS, and AD progression. Assessments can then be made of the impact of genetic factors, and volitional elements such as diet, disease, co-existing drug and dietary supplement regimens, and lifestyle issues. The goal would be to grasp the range of AD's neuroenergetic footings to develop an effective method of integrated, personalized AD risk assessments. Interventions could then be investigated for their potential effects on the net availability of energy resources in the CNS. Additionally, non-GLUT1-dependent energy resources can be studied for their ability to affect AD's characteristic A β /tau proclivity directly, or indirectly through increasing available glucose or alternate energy resources in the CNS. Research efforts could then focus on examining various methods to maintain adequate CNS energy substrates.

The key will be an identification and mainstream application of factors associated with a non-amyloidogenic CNS energy profile.

ACKNOWLEDGMENTS

Sincere thanks to Drs. William Pereira, Richard Bordow, Steven Jacobsohn, and John Swartzberg for critical reading and valuable guidance in the preparation of this manuscript.

The author's disclosure is available online (<http://j-alz.com/manuscript-disclosures/17-0549r1>).

REFERENCES

- [1] Alzheimer's Association (2017) 2017 Alzheimer's disease facts and figures. Available from: http://www.alz.org/documents_custom/2017-facts-and-figures.pdf
- [2] Deb A, Thornton JD, Sambamoorthi U, Innes K (2017) Direct and indirect cost of managing Alzheimer's disease and related dementias in the United States. *Expert Rev Pharmacoecon Outcomes Res* **17**, 189-202.
- [3] Hardy JA, Higgins GA (1992) Alzheimer's disease: The amyloid cascade hypothesis. *Science* **256**, 184-185.
- [4] Swerdlow RH, Khan SM (2004) A "mitochondrial cascade hypothesis" for sporadic Alzheimer's disease. *Med Hypotheses* **63**, 8-20.
- [5] Swerdlow RH, Burns JM, Khan SM (2014) The Alzheimer's disease mitochondrial cascade hypothesis: Progress and perspectives. *Biochim Biophys Acta* **1842**, 129-131.
- [6] Brewer GJ (2017) Copper-2 hypothesis for causation of the current Alzheimer's disease epidemic together with dietary changes that enhance the epidemic. *Chem Res Toxicol* **30**, 763-768.
- [7] Bush AI, Pettingell WH, Multhaup G, Paradis M, Vonsattel JP, Gusella JF, Beyreuther K, Masters CL, Tanzi RE (1994) Rapid induction of Alzheimer A β amyloid formation by zinc. *Science* **265**, 464-467.
- [8] Lovell MA, Robertson JD, Teesdale WJ, Campbell JL, Markesbery WR (1994) Copper, iron and zinc in Alzheimer's disease senile plaques. *J Neurol Sci* **158**, 47-52.
- [9] Mirza A, King A, Troakes C, Exley C (2017) Aluminium in brain tissue in familial Alzheimer's disease. *J Trace Elem Med Biol* **40**, 30-36.
- [10] Alzheimer's Association Calcium Hypothesis Workgroup (2017) Calcium hypothesis of Alzheimer's disease and brain aging: A framework for integrating new evidence into a comprehensive theory of pathogenesis. *Alzheimers Dement* **13**, 178-182.e17.
- [11] McGeer PL, Rogers J, McGeer EG (2016) Inflammation, anti-inflammatory agents, and Alzheimer's disease: The last 22 years. *J Alzheimers Dis* **54**, 853-857.
- [12] de la Monte SM, Wands JR (2008) Alzheimer's disease is type 3 diabetes-evidence reviewed. *J Diabetes Sci Technol* **2**, 1101-1113.
- [13] Erickson MA, Banks WA (2013) Blood-brain barrier dysfunction as a cause and consequence of Alzheimer's disease. *J Cereb Blood Flow Metab* **33**, 1500-1513.
- [14] Daulatzai MA (2016) Cerebral hypoperfusion and glucose hypometabolism: Key pathophysiological modulators

- promote neurodegeneration, cognitive impairment, and Alzheimer's disease. *J Neurosci Res* **95**, 943-972.
- [15] Alata W, Ye Y, St-Amour I, Vandal M, Calon F (2015) Human apolipoprotein E ϵ 4 expression impairs cerebral vascularization and blood-brain barrier function in mice. *J Cereb Blood Flow Metab* **35**, 86-94.
- [16] Verghese PB, Castellano JM, Holtzman DM (2011) Apolipoprotein E in Alzheimer's disease and other neurological disorders. *Lancet Neurol* **10**, 241-252.
- [17] Winkler EA, Nishida Y, Sagare AP, Rege SV, Bell RD, Perlmutter D, Sengillo JD, Hillman S, Kong P, Nelson AR, Sullivan JS, Zhao Z, Meiselman HJ, Wenby RB, Soto J, Abel ED, Makshanoff J, Zuniga E, De Vivo DC, Zlokovic BV (2015) GLUT1 reductions exacerbate Alzheimer's disease vasculo-neuronal dysfunction and degeneration. *Nat Neurosci* **18**, 521-530.
- [18] Prasad S, Sajja RK, Naik P, Cucullo L (2014) Diabetes mellitus and blood-brain barrier dysfunction: An overview. *J Pharmacovigil* **2**, 125.
- [19] Heneka MT, Nicotera P (2016) Thoughts on obesity and brain glucose. *Cell* **165**, 773-775.
- [20] Lyros E, Bakogiannis C, Liu Y, Fassbender K (2014) Molecular links between endothelial dysfunction and neurodegeneration in Alzheimer's disease. *Curr Alzheimer Res* **11**, 18-26.
- [21] Carbone I, Lazzarotto T, Ianni M, Porcellini E, Forti P, Masliah E, Gabrielli L, Licastro F (2014) Herpes virus in Alzheimer's disease: Relation to progression of the disease. *Neurobiol Aging* **35**, 122-129.
- [22] Talman V, Pascale A, Jäntti M, Amadio M, Tuominen RK (2016) Protein kinase C activation as a potential therapeutic strategy in Alzheimer's disease: Is there a role for embryonic lethal abnormal vision-like proteins? *Basic Clin Pharmacol Toxicol* **119**, 149-160.
- [23] Govoni S, Amadio M, Battaini F, Pascale A (2010) Senescence of the brain: Focus on cognitive kinases. *Curr Pharm Des* **16**, 660-671.
- [24] Sasaki N, Fukatsu R, Tsuzuki K, Hayashi Y, Yoshida T, Fujii N, Koike T, Wakayama I, Yanagihara R, Garruto R, Amano N, Makita Z (1998) Advanced glycation end products in Alzheimer's disease and other neurodegenerative diseases. *Am J Pathol* **153**, 1149-1155.
- [25] Morris MC, Evans DA, Bienias JL, Tangney CC, Bennett DA, Aggarwal N, Schneider J, Wilson RS (2003) Dietary fats and the risk of incident Alzheimer disease. *Arch Neurol* **60**, 194-200. Erratum in: *Arch Neurol* **60**, 1072.
- [26] Mirza A, King A, Troakes C, Exley C (2016) The identification of aluminum in human brain tissue using lumogallion and fluorescence microscopy. *J Alzheimers Dis* **54**, 1333-1338.
- [27] Wang B, Xing W, Zhao Y, Deng X (2010) Effects of chronic aluminum exposure on memory through multiple signal transduction pathways. *Environ Toxicol Pharmacol* **29**, 308-313.
- [28] Tyas SL (2001) Alcohol use and the risk of developing Alzheimer's disease. *Alcohol Res Health* **25**, 299-306.
- [29] Meneilly GS, Tessier DM (2016) Diabetes, dementia and hypoglycemia. *Can J Diabetes* **40**, 73-76.
- [30] Xie S, Jin N, Gu J, Shi J, Sun J, Chu D, Zhang L, Dai CL, Gu JH, Gong CX, Iqbal K, Liu F (2016) O-GlcNAcylation of protein kinase A catalytic subunits enhances its activity: A mechanism linked to learning and memory deficits in Alzheimer's disease. *Aging Cell* **15**, 455-464.
- [31] Kummer MP, Heneka MT (2014) Truncated and modified amyloid-beta species. *Alzheimers Res Ther* **6**, 28.
- [32] Chow VW, Mattson MP, Wong PC, Gleichmann M (2010) An overview of APP Processing enzymes and products. *Neuromolecular Med* **12**, 1-12.
- [33] Merlini G, Bellotti V (2003) Molecular mechanisms of amyloidosis. *N Engl J Med* **349**, 583-596.
- [34] Bourgade K, Le Page A, Bocti C, Witkowski JM, Dupuis G, Frost EH, Fülöp T (2016) Protective effect of amyloid- β peptides against herpes simplex virus-1 infection in a neuronal cell culture model. *J Alzheimers Dis* **50**, 1227-1241.
- [35] Bourgade K, Dupuis G, Frost EH, Fülöp T (2016) Antiviral properties of amyloid- β peptides. *J Alzheimers Dis* **54**, 859-878.
- [36] Wozniak MA, Itzhaki RF, Shipley SJ, Dobson CB (2007) Herpes simplex virus infection causes cellular beta amyloid accumulation and secretase upregulation. *Neurosci Lett* **429**, 95-100.
- [37] Velliquette RA, O'Connor T, Vassar R (2005) Energy inhibition elevates beta-secretase levels and activity and is potentially amyloidogenic in APP transgenic mice: Possible early events in Alzheimer's disease pathogenesis. *J Neurosci* **25**, 10874-10883.
- [38] Skovronsky DM, Moore DB, Milla ME, Doms RW, Lee VM (2000) Protein kinase C-dependent alpha-secretase competes with beta-secretase for cleavage of amyloid-beta precursor protein in the trans-golgi network. *J Biol Chem* **275**, 2568-2575.
- [39] Klosinski LP, Yao J, Yin F, Fonteh AN, Harrington MG, Christensen TA, Trushina E, Brinton RD (2015) White matter lipids as a ketogenic fuel supply in aging female brain: Implications for Alzheimer's disease. *EBioMedicine* **2**, 1888-1904.
- [40] Bartzokis G (2011) Alzheimer's disease as homeostatic responses to age-related myelin breakdown. *Neurobiol Aging* **32**, 1341-1371.
- [41] Singh P, Jain A, Kaur G (2004) Impact of hypoglycemia and diabetes on CNS: Correlation of mitochondrial oxidative stress with DNA damage. *Mol Cell Biochem* **260**, 153-159.
- [42] Bush AI, Tanzi RE (2008) Therapeutics for Alzheimer's disease based on the metal hypothesis. *Neurotherapeutics* **5**, 421-432.
- [43] Duce JA, Bush AI (2010) Biological metals and Alzheimer's disease: Implications for therapeutics and diagnostics. *Prog Neurobiol* **92**, 1-18.
- [44] Greenough MA, Camakaris J, Bush AI (2013) Metal dyshomeostasis and oxidative stress in Alzheimer's disease. *Neurochem Int* **62**, 540-555.
- [45] Hasselbalch SG, Knudsen GM, Videbaek C, Pinborg LH, Schmidt JF, Holm S, Paulson OB (1999) No effect of insulin on glucose blood-brain barrier transport and cerebral metabolism in humans. *Diabetes* **48**, 1915-1921.
- [46] Freude S, Plum L, Schnitker J, Leeser U, Udelhoven M, Krone W, Bruning JC, Schubert M (2005) Peripheral hyperinsulinemia promotes tau phosphorylation *in vivo*. *Diabetes* **54**, 3343-3348.
- [47] Ribe EM, Lovestone S (2016) Insulin signalling in Alzheimer's disease and diabetes: From epidemiology to molecular links. *J Intern Med* **280**, 430-442.
- [48] Craft S, Claxton A, Baker LD, Hanson AJ, Cholerton B, Trittschuh EH, Dahl D, Caulder E, Neth B, Montine TJ, Jung Y, Maldjian J, Whitlow C, Friedman S (2017) Effects of regular and long-acting insulin on cognition and Alzheimer's

- disease biomarkers: A pilot clinical trial. *J Alzheimers Dis* **57**, 1325-1334.
- [49] Plum L, Schubert M, Brüning JC (2005) The role of insulin receptor signaling in the brain. *Trends Endocrinol Metab* **16**, 59-65.
- [50] Marks DR, Tucker K, Cavallin MA, Mast TG, Fadool DA (2009) Awake intranasal insulin delivery modifies protein complexes and alters memory, anxiety, and olfactory behaviors. *J Neurosci* **29**, 6734-6751.
- [51] Vlad SC, Miller DR, Kowall NW, Felson DT (2008) Protective effects of NSAIDs on the development of Alzheimer disease. *Neurology* **70**, 1672-1677.
- [52] Kummer MP, Heneka MT (2008) PPARs in Alzheimer's disease. *PPAR Res* **2008**, 403896.
- [53] Backes H, Walberer M, Ladwig A, Rueger MA, Neumaier B, Endepols H, Hoehn M, Fink GR, Schroeter M, Graf R (2016) Glucose consumption of inflammatory cells masks metabolic deficits in the brain. *Neuroimage* **128**, 54-62.
- [54] Cahill GF Jr, Veech RL (2003) Ketoacids? Good medicine? *Trans Am Clin Climatol Assoc* **114**, 149-161.
- [55] Bergman EN (1990) Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiol Rev* **70**, 567-590.
- [56] Bourassa MW, Alim I, Bultman SJ, Ratan RR (2016) Butyrate, neuroepigenetics and the gut microbiome: Can a high fiber diet improve brain health? *Neurosci Lett* **625**, 56-63.
- [57] Smith AP, Sutherland D, Hewlett P (2015) An investigation of the acute effects of oligofructose-enriched inulin on subjective wellbeing, mood and cognitive performance. *Nutrients* **7**, 8887-8896.
- [58] Stilling RM, van de Wouw M, Clarke G, Stanton C, Dinan TG, Cryan JF (2016) The neuropharmacology of butyrate: The bread and butter of the microbiota-gut-brain axis? *Neurochem Int* **99**, 110-132.
- [59] Henderson ST, Vogel JL, Barr LJ, Garvin F, Jones JJ, Costantini LC (2009) Study of the ketogenic agent AC-1202 in mild to moderate Alzheimer's disease: A randomized, double-blind, placebo-controlled, multicenter trial. *Nutr Metab (Lond)* **6**, 31.
- [60] Newport MT, VanItallie TB, Kashiwaya Y, King MT, Veech RL (2015) A new way to produce hyperketonemia: Use of ketone ester in a case of Alzheimer's disease. *Alzheimers Dement* **11**, 99-103.
- [61] Cunnane SC, Courchesne-Loyer A, St-Pierre V, Vandenberghe C, Pierotti T, Fortier M, Croteau E, Castellano CA (2016) Can ketones compensate for deteriorating brain glucose uptake during aging? Implications for the risk and treatment of Alzheimer's disease. *Ann N Y Acad Sci* **1367**, 12-20.
- [62] Mochel F, Hainque E, Gras D, Adanyeguh IM, Caillet S, Héron B, Roubertie A, Kaphan E, Valabregue R, Rinaldi D, Vuillaumier S, Schiffmann R, Ottolenghi C, Hogrel JY, Servais L, Roze E (2016) Triheptanoin dramatically reduces paroxysmal motor disorder in patients with GLUT1 deficiency. *J Neurol Neurosurg Psychiatry* **87**, 550-553.
- [63] Studzinski CM, MacKay WA, Beckett TL, Henderson ST, Murphy MP, Sullivan PG, Burnham WM (2008) Induction of ketosis may improve mitochondrial function and decrease steady-state amyloid-beta precursor protein (APP) levels in the aged dog. *Brain Res* **1226**, 209-217.
- [64] Yin JX, Maalouf M, Han P, Zhao M, Gao M, Dharshaun T, Ryan C, Whitelegge J, Wu J, Eisenberg D, Reiman EM, Schweizer FE, Shi J (2016) Ketones block amyloid entry and improve cognition in an Alzheimer's model. *Neurobiol Aging* **39**, 25-37.
- [65] Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, Ganguli M, Hall K, Hasegawa K, Hendrie H, Huang Y, Jorm A, Mathers C, Menezes PR, Rimmer E, Sczufca M, Alzheimer's Disease International (2005) Global prevalence of dementia: A Delphi consensus study. *Lancet* **366**, 2112-2117.
- [66] Petersson SD, Philippou E (2016) Mediterranean diet, cognitive function, and dementia: A systematic review of the evidence. *Adv Nutr* **7**, 889-904.
- [67] Yusuf M, Weyand LL, Piryatinsky I (2017) Alzheimer's disease and diet: A systematic review. *Int J Neurosci* **127**, 161-175.
- [68] Morris MC, Tangney CC, Wang Y, Sacks FM, Bennett DA, Aggarwal NT (2015) MIND diet associated with reduced incidence of Alzheimer's disease. *Alzheimers Dement* **11**, 1007-1014.
- [69] Maliszewska-Cyna E, Lynch M, Oore JJ, Nagy PM, Aubert I (2017) The benefits of exercise and metabolic interventions for the prevention and early treatment of Alzheimer's disease. *Curr Alzheimer Res* **14**, 47-60.
- [70] Quistorff B, Secher NH, Van Lieshout JJ (2008) Lactate fuels the human brain during exercise. *FASEB J* **22**, 3443-3449.
- [71] Simpson IA, Carruthers A, Vannucci SJ (2007) Supply and demand in cerebral energy metabolism: The role of nutrient transporters. *J Cereb Blood Flow Metab* **27**, 1766-1791.
- [72] Tucsek Z, Toth P, Sosnowska D, Gautam T, Mitschelen M, Koller A, Szalai G, Sonntag WE, Ungvari Z, Csiszar A (2014) Obesity in aging exacerbates blood-brain barrier disruption, neuroinflammation, and oxidative stress in the mouse hippocampus: Effects on expression of genes involved in beta-amyloid generation and Alzheimer's disease. *J Gerontol A Biol Sci Med Sci* **69**, 1212-1226.
- [73] Hishikawa N, Fukui Y, Sato K, Kono S, Yamashita T, Ohta Y, Deguchi K, Abe K (2016) Cognitive and affective functions in Alzheimer's disease patients with metabolic syndrome. *Eur J Neurol* **23**, 339-345.
- [74] Nelson L, Tabet N, Richardson C, Gard P (2013) Antihypertensives, angiotensin, glucose and Alzheimer's disease. *Expert Rev Neurother* **13**, 477-482.
- [75] Fishel MA, Watson GS, Montine TJ, Wang Q, Green PS, Kulstad JJ, Cook DG, Peskind ER, Baker LD, Goldgaber D, Nie W, Asthana S, Plymate SR, Schwartz MW, Craft S (2005) Hyperinsulinemia provokes synchronous increases in central inflammation and beta-amyloid in normal adults. *Arch Neurol* **62**, 1539-1544.
- [76] Arrieta-Cruz I, Gutiérrez-Juárez R (2016) The role of insulin resistance and glucose metabolism dysregulation in the development of Alzheimer's disease. *Rev Invest Clin* **68**, 53-58.