

## Research Report

# Fixation Duration and Pupil Size as Diagnostic Tools in Parkinson's Disease

Panagiota Tsitsi<sup>a,b,\*</sup>, Mattias Nilsson Benfatto<sup>c</sup>, Gustaf Öqvist Seimyr<sup>c</sup>, Olof Larsson<sup>c</sup>, Prof Per Svenningsson<sup>a,b,d</sup> and Ioanna Markaki<sup>a,b</sup>

<sup>a</sup>Department of Clinical Neuroscience, Neuro, Karolinska Institutet, Stockholm, Sweden

<sup>b</sup>Center of Neurology, Academic Specialist Center, Region Stockholm, Sweden

<sup>c</sup>Department of Clinical Neuroscience, Eye and Vision, Karolinska Institutet, Stockholm, Sweden

<sup>d</sup>Neurology Department, Karolinska University Hospital, Stockholm, Sweden

Accepted 28 January 2021  
Pre-press 16 February 2021

### Abstract.

**Background:** Visual and oculomotor problems are very common in Parkinson's disease (PD) and by using eye-tracking such problems could be characterized in more detail. However, eye-tracking is not part of the routine clinical investigation of parkinsonism.

**Objective:** To evaluate gaze stability and pupil size in stable light conditions, as well as eye movements during sustained fixation in a population of PD patients and healthy controls (HC).

**Methods:** In total, 50 PD patients (66% males) with unilateral to mild-to-moderate disease (Hoehn & Yahr 1–3, Schwab and England 70–90%) and 43 HC (37% males) were included in the study. Eye movements were recorded with Tobii Pro Spectrum, a screen-based eye tracker with a sampling rate of 1200 Hz. Logistic regression analysis was applied to investigate the strength of association of eye-movement measures with diagnosis.

**Results:** Median pupil size (OR 0.811; 95% CI 0.666–0.987;  $p = 0.037$ ) and longest fixation period (OR 0.798; 95% CI 0.691–0.921;  $p = 0.002$ ), were the eye-movement parameters that were independently associated with diagnosis, after adjustment for sex (OR 4.35; 95% CI 1.516–12.483;  $p = 0.006$ ) and visuospatial/executive score in Montreal Cognitive Assessment (OR 0.422; 95% CI 0.233–0.764;  $p = 0.004$ ). The area under the ROC curve was determined to 0.817; 95% (CI) 0.732–0.901.

**Conclusion:** Eye-tracking based measurements of gaze fixation and pupil reaction may be useful biomarkers of PD diagnosis. However, larger studies of eye-tracking parameters integrated into the screening of patients with suspected PD are necessary, to further investigate and confirm their diagnostic value.

Keywords: Parkinson's disease, eye-tracking, fixation, pupil size, eye movements

## INTRODUCTION

Visual and oculomotor problems in Parkinson's disease (PD) are among the most common non-motor symptoms reported by patients. Diplopia (double

vision), alterations in visuoperception, impairment in contrast, and color vision, as well as visual hallucinations, are some of the usual complaints [1]. Common findings during clinical evaluation are hypometric saccades, hyperreflexivity, increased latency of voluntary saccades, and saccadic intrusions during smooth pursuit eye movements [2]. These symptoms are not specific for PD although the majority appear early in the course of the disease. So far, the use of

\*Correspondence to: Panagiota Tsitsi, MD, Center for Neurology, Academic Specialist Center, Solnavägen 1E, 113 65 Stockholm, Sweden. Tel.: +46 765678180; E-mail: panagiota.tsitsi@ki.se.

eye-tracking to assess eye movements has not been part of the routine clinical workup in parkinsonian disorders mainly because, to date, most high-end eye-tracking systems suitable for clinical research have been rather complicated to use by non-experts, making them impractical as an everyday assessment tool in typical clinical settings.

Research on eye movements and oculomotor control in PD has traditionally focused on saccades and saccadic performance using test paradigms like the pro- and anti-saccade task. Small eye movements made during attempted fixation have received less attention although this has started to change in recent years due to the renewed interest in fixational eye movements and their relation to visual perception, attention, and cognition. The role of fixation is to maintain the image of the object of interest in the fovea which is the area of the retina where the visual acuity is best. To keep the target near the fovea, multiple brain structures are recruited. Apart from excitatory activity that stabilizes the eyes, inhibitory processes during fixation prevent saccadic eye movements that would break it. Both cortical (frontal and parietal areas) and subcortical domains (such as the nucleus raphe interpositus and superior colliculus) contribute to this equilibrium [3]. Nevertheless, if the eye was completely stable during the visual fixation of a target, sensor adaptation on the retina would lead to blurring and eventually fading of the image. Therefore, corrective microsaccades, tremor, and conjugational drifts are necessary, not only to improve the function of fixation but also to correct errors [4]. Apart from these mini regulatory movements, saccadic intrusions may interrupt fixation. In the case of PD, saccadic intrusions are common, and they appear in the form of square wave jerks (SWJ) that move the eye away from the target during sustained fixation or smooth pursuit. A return saccade back to the target follows the SWJ after approximately 200 ms. The role of the superior colliculus and the fastigial oculomotor region of the cerebellum has been discussed in the genesis of SWJ [5].

Another interesting factor, apart from gaze stability, that can be studied during sustained fixation, while a person is required to maintain focus on a stable target, is the size of the pupil that is affected both by light and cognitive processes that include target detection and attention [6]. Changes in pupil reactivity have already been described in PD and the role of the parasympathetic autonomic nervous system in impaired pupil reflex has previously been discussed [7]. More specifically, it has been shown

that cholinergic deficits in PD patients with cognitive impairment may affect the pupil light reflex [8]. Studies on pupil's reaction to light have yielded rather controversial results due to the role of psychiatric features as well as cognitive impairment but despite the extended literature on light reflex in PD, studies on pupil size in stable light conditions during fixation are scarce.

The purpose of this study was to evaluate gaze stability and pupil size in stable light conditions during sustained fixation in a population of PD patients and healthy controls (HC). Eye-tracking is an objective, non-invasive, and cost-effective analysis method that enables accurate and detailed gaze and eye-movement examination, where processing with advanced algorithms allows for accurate calculation of eye position and movement parameters. It is easy to use in different environments and does not require any verbal, written, or other active actions by the test person.

## MATERIALS AND METHODS

### *Participants*

Fluent Swedish-speaking PD patients with Hoehn and Yahr (H&Y)  $\leq 3$  and age-matched healthy volunteers were included in the study. The diagnosis was based on the United Kingdom Brain Bank Criteria [9]. Individuals suffering from eye conditions such as macular degeneration or non-operable/non-corrected cataract were excluded and vision was normal or corrected to normal (visual acuity of a logMAR score  $\leq 0.00$ ), regarding both, refraction, and presbyopia. The study was approved by the Stockholm Ethical Committee (DNR: 2018/437-31/2) and participants provided written and oral informed consent, according to the declaration of Helsinki. In total, 50 PD patients (66% males) with unilateral to mild-to-moderate disease (H&Y 1–3, Schwab and England 70–90%) and 43 HC (37% males) were recruited at the Center of Neurology, Academic Specialist Center in Stockholm, Sweden. The participants were clinically assessed with the Unified Parkinson's Disease Rating Scale (UPDRS) [10] as well as the Montreal Clinical Assessment (MoCA), Mini-Mental State Examination (MMSE), and Frontal Assessment Battery (FAB).

### *Apparatus and stimuli presentation*

Eye movements were recorded with Tobii Pro Spectrum, a screen-based eye tracker with a sampling

143 rate of 1200 Hz. Data were recorded binocularly, i.e.,  
144 from left and right eye simultaneously. Stimuli were  
145 presented on the native 23.8" Tobii Pro Spectrum  
146 screen (EIZO FlexScan EV2451) with pixel reso-  
147 lution 1920×1080 (52.8×29.7 cm). The screen was  
148 located approximately 65 cm in front of the partic-  
149 ipant who was sitting on a steady and comfortable  
150 chair in a dimly lit room. Visual acuity was tested with  
151 the Landolt C Chart and participants were included  
152 only if they received a logMAR score of 0.00 or less,  
153 after correcting for refraction and/or presbyopia. The  
154 participant was given clear and simple instructions  
155 by the examiner to keep his or her eyes focused on  
156 the black "dot" (6 mm diameter fixation target sub-  
157 tending approximately 0.5 degrees of visual angle)  
158 in the center of a bright white screen. The fixa-  
159 tion target was on display for 15 seconds in each  
160 trial, after which it briefly disappeared for 5 sec-  
161 onds. Each participant performed eight trials with a  
162 total duration of 3 minutes, providing 2 minutes of  
163 fixation data for analysis. A 5-point calibration fol-  
164 lowed by four points of validation was performed  
165 on each participant prior to the first trial. Recal-  
166 ibrations were made if deemed necessary by the  
167 examiner.

### 168 *Data processing*

169 The recorded data were filtered through the Tobii I-  
170 VT (Identification by Velocity Threshold) algorithm  
171 available in the analysis software Tobii Pro Lab. The  
172 primary purpose of the filtering algorithm is to iden-  
173 tify fixations and saccades in the raw gaze data. Given  
174 a sampling frequency of 1200 Hz, the eye tracker pro-  
175 duces a gaze sample approximately every 0.83 ms.  
176 The I-VT filter uses a velocity criterion to calculate  
177 whether a sequence of raw gaze samples belongs to  
178 the same fixation or whether they are part of a saccade  
179 in progress.

180 Since eye-movement patterns may vary substan-  
181 tially between different types of tasks and stimuli, it  
182 is often necessary to adapt the fixation filter's param-  
183 eter settings to the type and quality of data at hand.  
184 We used the algorithm's default settings in the fil-  
185 tering, with two exceptions: (1) a 5-point running  
186 median was used to reduce the level of noise (default:  
187 3-point running median); (2) gap fill-in interpola-  
188 tion was used to fill in short periods of data loss  
189 (default: gap fill-in interpolation is inactive). We used  
190 the default value for the velocity threshold, which  
191 is 30°/s, to allow for relatively short and fast move-  
192 ments to be detected as saccades. It is worth noting

193 that all recordings were subjected to the same filter-  
194 ing, which is standard practice, i.e., the parameters  
195 of the I-VT filter were not adjusted to fit each partic-  
196 ipant or recording individually. After data filtering,  
197 a set of eye-movement parameters were computed  
198 over the fixation target stimulus for each trial. In line  
199 with Castet and Crossland [11], the first 1.5 seconds  
200 of each trial were removed before the calculation in  
201 order to discard data when the participant was first  
202 locating the fixation target in a trial. All computed  
203 gaze parameters were based on left and right eye aver-  
204 ages. However, when only one eye was found for a  
205 data sample, that eye was used in the computation.

206 Two categories of parameters were computed:  
207 sample- and event-based. Sample-based parameters  
208 were based on the stream of eye-tracker samples  
209 rather than on fixation and saccade events detected  
210 by the I-VT fixation algorithm. These measures pro-  
211 vide information on the horizontal and vertical gaze  
212 position (degrees of visual angle), and bivariate con-  
213 tour ellipse area (BCEA) (square degrees of visual  
214 angle). The BCEA measures dispersion as the area  
215 of an ellipse encompassing a given proportion  $P$   
216 of gaze points. Here we used  $P=0.682$ , thus giving the  
217 area of the ellipse over which gaze positions were  
218 found 68.2% of the time (i.e., dispersion of the gaze  
219 about its mean position with  $\pm 1$  standard deviation).  
220 Lower BCEA-values indicate higher/better fixation  
221 stability. Pupil size (mm) was also computed from  
222 the stream of eye-tracker samples. By contrast, event-  
223 based parameters, such as fixation duration (sec)  
224 and saccade rate (number of detected saccades per  
225 second), during fixation, were computed over the  
226 detected events, either fixations or saccades and they  
227 depend on the I-VT algorithm's ability to identify  
228 them. The mean and median values over the trials  
229 were then computed for each participant. Since the  
230 median is generally more tolerant to outliers than the  
231 mean, eye-movement measures that were aggregated  
232 over the trials are given as the medians across the  
233 trials. For example, the "mean pupil size" parameter  
234 reflects the mean of medians of diameters of the pupil  
235 measured during the fixation task.

236 The gaze parameters primarily attempt to quan-  
237 tify the fixational eye stability during the task. A  
238 possible hypothesis is that PD participants are less  
239 stable in their fixation than the control group and less  
240 able to keep their eyes fixated over time. If true, we  
241 would expect, for example, higher dispersion around  
242 the mean fixation position, as well as shorter periods  
243 of fixation uninterrupted by saccades in the PD group  
244 compared to the control group.

## Statistical analysis

Statistical analysis was done with IBM SPSS 25  
Statistic Data Editor. Non-parametric tests were used,  
and the significance level was defined at 0.05. Logistic  
regression analysis was applied to investigate the  
strength of association of the eye-movement mea-  
sures, alone or in combination, as predictors of dia-  
gnosis (HC vs PD), in separate multivariate models,  
including also sex, age, and cognitive scores. Re-  
ceiver Operating Characteristic (ROC) curve was  
plotted to visualize the final model's separation  
potential between diagnoses. Spearman correlation  
was used to investigate the correlation between eye-  
movement measures and cognitive scores.

## RESULTS

### Participants' characteristics

In total, 50 PD patients and 43 HC were included in  
the analysis. Sex distribution differed with more men  
included in the PD than the HC group ( $p=0.006$ ),  
whereas age and years of education did not differ  
between groups (Table 1). Median disease duration  
in the PD group was 2 years, corresponding to early-  
stage disease, as indicated also by the median LEDD  
(545) and UPDRS part 3 (21 points).

With regard to cognition, MoCA, MMSE, and FAB  
scores did not differ significantly between PD patients  
and HC. However, PD patients had lower score than  
HC in the visuospatial/executive domain in MoCA  
(median 4 vs 5;  $p=0.005$ ; Table 2).

Overall, regarding comorbidities and concomi-  
tant treatments (Table 1), we only found statistically  
significant differences regarding prostate hyperplasia  
and/or urinary incontinence (1 HC vs 9 PD,  
 $p=0.018$ ), and B12 supplementation that was more  
common in the PD group (3 HC vs 16 PD,  $p=0.04$ ).  
Only one HC had a diagnosis of diabetes mellitus.  
In the PD group, four patients were receiving  
benzodiazepines (three patients received low dose  
clonazepam for the treatment of REM sleep behav-  
ior disorder and one was treated with oxazepam as  
required against anxiety). Additionally, three patients  
were treated with anticholinergic medication; two  
were treated with solifenacin for urinary inconti-  
nence and the other one under trihexyphenidyl for  
the treatment of tremor. Depression was more common  
in the PD group (1 HC vs 7 PD participants). One HC  
was treated for glaucoma versus three participants in  
the PD group.

Table 1

Clinical and demographical characteristics of PD patients and HC

	HC, N=43	PD, N=50	<i>p</i>
Sex, Male/Female	16/27	33/17	<b>0.006</b>
Age	63 (16)	64 (10.5)	0.728
Education, years	15 (5)	16 (4)	0.895
Age at onset,		61 (12)	NA
Age at diagnosis,		62 (11.5)	NA
Years since onset		4 (4.25)	NA
Years since diagnosis		2 (2.5)	NA
LEDD		545 (523.75)	NA
UPDRS part 1		1 (2)	NA
UPDRS part 2		10 (6)	NA
UPDRS part 3		21 (15.5)	NA
UPDRS part 4		2 (3.25)	NA
UPDRS total		36.5 (21.75)	NA
Schwab & England		90 (10)	NA
Comorbidities and treatments, % (n)			
Diabetes	2.3 (1)	0	0.5
Atrial fibrillation	2.3 (1)	8 (4)	0.4
Hypertension	23.3 (10)	30 (15)	0.5
Depression	2.3 (1)	14 (7)	0.07
Glaucoma	2.3 (1)	6 (3)	0.6
Prostate hyperplasia/ Urinary incontinence	2.3 (1)	18 (9)	<b>0.018</b>
B12 supplementation	7 (3)	32 (16)	<b>0.04</b>
Use of Benzodiazepines	0	8 (4)	0.12
Use of Anticholinergics	0	4 (2)	0.5

Values are reported as medians and interquartile ranges (IQR),  
apart from the gender ratio that is reported in absolute values.  
LEDD, Levodopa Equivalent Daily Dose; UPDRS, Unified Parkin-  
son's Disease Rating Scale; NA, non-applicable.

Table 2

Cognitive scores of PD patients and healthy controls HC

	HC, N=43	PD, N=50	<i>p</i>
MoCA -Visuospatial/Executive	5 (1)	4 (2)	<b>0.005</b>
MoCA - Naming	3 (0)	3 (0)	0.2
MoCA - Attention	6 (0)	6 (1)	0.5
MoCA - Language	3 (1)	3 (1)	0.3
MoCA - Abstraction	2 (0)	2 (0)	0.4
MoCA - Delayed Recall	3 (2)	3 (2)	0.9
MoCA - Orientation	6 (0)	6 (0)	0.2
MoCA Total score	27 (3)	27 (3)	0.2
MMSE Total score	29 (2)	28 (2)	0.4
FAB Total score	18 (2)	17 (3)	0.1

Values are reported as medians and interquartile ranges (IQR).  
FAB, Frontal Assessment Battery; MMSE, Mini-Mental State  
Examination; MoCA, Montreal Cognitive Assessment.

### Fixation task results

Tables 3 and 4 summarize the fixation task results  
in PD patients and HC. Regarding the sample-  
based parameters, comparison of horizontal and  
vertical gaze position showed significant differences  
in the median absolute deviation (MAD) and the

Table 3

Sample-based eye-movement parameters in PD patients and HC	HC, N=43	PD, N=50	<i>p</i>
SD gaze point horizontal	0.2 (0.37)	0.32 (0.59)	0.1
SD gaze point vertical	0.34 (0.42)	0.33 (0.57)	0.2
MAD gaze point horizontal	0.09 (0.05)	0.12 (0.2)	<b>0.023</b>
MAD gaze point vertical	0.08 (0.06)	0.11 (0.12)	0.1
QD gaze point horizontal	0.1 (0.05)	0.13 (0.27)	<b>0.028</b>
QD gaze point vertical	0.09 (0.07)	0.11 (0.19)	0.1
BCEA	0.5 (1.73)	0.8 (3.02)	0.2
Mean pupil size	2.5 (0.3)	2.36 (0.31)	<b>0.002</b>
Median pupil size	2.5(0.29)	2.36 (0.31)	<b>0.003</b>

Values are reported as medians and interquartile ranges (IQR). BCEA, Bivariate contour ellipse area (square degrees of visual angle); MAD gaze point, Median absolute deviation of the horizontal/vertical gaze position (degrees of visual angle); QD gaze point, Quartile deviation of the horizontal/gaze position (degrees of visual angle); SD gaze point, Standard deviation of the horizontal/vertical gaze position (degrees of visual angle). Mean and median pupil size computed in mm.

Table 4

Event-based eye-movement parameters in PD patients and HC	HC, N=43	PD, N=50	<i>p</i>
Mean fixation duration	3.02 (4.25)	1.3 (3.93)	<b>0.007</b>
Median fixation duration	2.55 (4.48)	0.73 (5.09)	<b>0.016</b>
Longest fixation period	6.1 (5.46)	4.35 (5.34)	<b>0.008</b>
Saccade rate	0.44 (0.97)	1.11 (2.49)	<b>0.015</b>

Values are reported as medians and interquartile ranges (IQR). Longest fixation period: The maximum period of fixation uninterrupted by saccades, blinks or noise (sec); Mean and median fixation duration computed in sec; Saccade rate: The number of detected saccades per second.

quartile deviation (QD) of the horizontal gaze position ( $p=0.023$  and  $p=0.028$ , respectively; Table 3). It is worth to notice that both standard deviation (SD) and BCEA use the mean in their formula, while MAD and QD are non-parametric estimates, and this could possibly explain the non-significant results in the SD and BCEA comparisons. Also, the horizontal component seems primarily affected, as there are no significant differences in the vertical components of the sample-based gaze measures. Moreover, both mean and median pupil diameter were larger in the HC participants than in PD ( $p=0.002$  and  $0.003$ , respectively).

Table 4 displays comparisons between event-based parameters computed between the PD and HC populations. Our results showed that during fixation, mean ( $p=0.007$ ) and median fixation duration ( $p=0.016$ ), and longest fixation period ( $p=0.008$ ) were significantly longer in the HC group than the PD group. In accordance with the above, the number of detected

saccades during the task was higher in the PD group ( $p=0.015$ ).

In order to address the fatigue effect, we examined each eye-movement parameter on a trial-by-trial basis for the two groups and compared them schematically. Parameters shown in Fig. 1 are measures of gaze variability/instability, which would be expected to increase in variability, in case fatigue-effect was present, however, this is not observed. Fixation parameters (Fig. 2) would be expected to show a decreasing trend with shorter fixations over time as participants become increasingly tired and find it harder to maintain fixation. This is observed to some extent, but the effect is of similar magnitude in the PD and HC group. Conversely, an increase in the rate of saccades over time would be expected due to fatigue. This is observed to some extent and somewhat more pronounced in the PD compared to the HC group. Yet, the saccade rate in the PD group is consistently higher than in HC, from the first to the last trial.



In the PD group, MMSE scores correlated moderately with the median fixation duration ( $\rho=0.361$ ;  $p=0.01$ ), longest fixation period ( $\rho=0.324$ ;  $p=0.022$ ) and saccade rate ( $\rho=-0.313$ ;  $p=0.028$ ) as well as gaze parameters: SD of the horizontal gaze position ( $\rho=-0.439$ ;  $p=0.001$ ), MAD of the horizontal ( $\rho=-0.482$ ;  $p<0.001$ ) and vertical ( $\rho=-0.473$ ;  $p=0.001$ ) gaze position, QD of horizontal ( $\rho=-0.457$ ;  $p=0.001$ ) and vertical ( $\rho=-0.441$ ;  $p=0.001$ ) gaze position, and BCEA ( $\rho=0.316$ ;  $p=0.026$ ). No correlation was found between the rest of the cognitive tests performed and eye-movement parameters.

Looking further for differences between male and female participants in the cohort, both mean and median pupil size difference was statistically significant ( $p=0.01$  and  $p=0.008$  respectively), with women presenting with larger pupil size (mm) than men during the task (median (IQR): 2.51 (0.28) vs 2.35 (0.28) and 2.59 (0.28) vs 2.46 (0.29) respectively).



#### Logistic regression analysis

Significant predictors of diagnosis in univariate models were: longest fixation period (OR 0.874; 95% CI 0.779–0.98;  $p=0.021$ ), mean and median pupil size (OR 0.798; 95% CI 0.664–0.939;  $p=0.007$  and OR 0.794; 95% CI 0.669–0.943;  $p=0.008$  respectively), saccade rate (OR 1.476; 95% CI 1.042–2.09;  $p=0.028$ ), visuospatial/executive score (OR 0.514;



Gaze point: Standard deviation (SD)



Gaze point: Median absolute deviation (MAD)



Gaze point: Quartile deviation (QD)



Gaze point: Bivariate contour ellipse area (BCEA)








Fig. 1. Gaze point parameters on a trial-by-trial basis in PD and HC groups.


Pupil size



Fixation duration



Longest fixation period



Saccade rate




Fig. 2. Mean and median pupil size, mean and median fixation duration, longest fixation period and saccade rate, on a trial-by-trial basis in PD and HC groups.

367 95% CI 0.32–0.825;  $p=0.006$ ), sex (OR 3.276;  
 368 95% CI 1.398–7.674;  $p=0.006$ ). The final model  
 369 included the median pupil size (OR 0.811; 95% CI  
 370 0.666–0.987;  $p=0.037$ ), longest fixation period (OR  
 371 0.798; 95% CI 0.691–0.921;  $p=0.002$ ), sex (OR  
 372 4.35; 95% CI 1.516–12.483;  $p=0.006$ ), visuospatial/executive score in MoCA (OR 0.422; 95% CI  
 373 0.233–0.764;  $p=0.004$ ). In all univariate and multi-  
 374 variate models we transformed the mean and median  
 375 pupil size by multiplying by ten; the models, there-  
 376 fore, refer to mean and median pupil size in mm/ 10.  
 377

ROC analysis was used to visualize the multivariate  
 model's potential to separate between HC and PD.  
 The area under the ROC was determined to 0.817;  
 95% CI 0.732–0.901 (Fig. 1).

To assess residual confounders, conditional multi-  
 variate logistic regression analysis was repeated in  
 a subgroup of 28 PD patients and 28 HC matched  
 for age ( $\pm 1$  year) and sex, and the results remained  
 unchanged with OR for median pupil size 0.57, 95%  
 CI 0.33–0.98 ( $p=0.04$ ), and OR for longest fixation  
 period 0.67, 95% CI 0.46–0.99 ( $p=0.04$ ).

378  
 379  
 380  
 381  
 382  
 383  
 384  
 385  
 386  
 387




Fig. 3. Receiver Operating Characteristic (ROC) curve of the multivariate model that includes longest fixation period, median pupil size, sex, and visuospatial/executive subscore of the Montreal Cognitive assessment test. AUC, Area under the curve; CI, Confidence interval.

## DISCUSSION

In our study, we report the results of eye-movement measurements, and more specifically parameters of the fixation task examined with eye-tracker, as potential diagnostic biomarkers in early-stage PD. We found that HC kept their eyes more stable during a longer period on a stable target, whereas fixation was interrupted more easily in PD patients. Additionally, during effort and in stable luminance conditions, PD patients' pupils were smaller than those of HC.

The fixation task required participants to keep their gaze on a dark target in the middle of a white screen for approximately three minutes, with short pauses when the target disappeared. Since the contrast between the background and the target, a simple dot, was sharp, and given that vision was corrected to normal, identifying the target, itself, was not challenging. Our results showed that PD patients were easily distracted during fixation; a higher median saccade rate and a shorter median longest fixation duration point to the same conclusion. This is in accordance with previous studies that have also reported higher distractibility and frequent saccadic intrusions during fixation in PD patients compared to HC, independently of cognitive status and PD stage [2, 12, 13]. Overall, PD patients disengage their attention easily compared to healthy adults. This phenomenon

has widely been studied in order to investigate its relationship with anatomical and functional structures. Inhibition of unnecessary eye movements requires the involvement of cortical (dorsolateral prefrontal cortex) and subcortical (superior colliculus) brain structures [14] as well as an intact connection between the basal ganglia and these structures. These connective loops have been described to be impaired in PD, thus generating abnormal saccades, some of which are SWJ that are commonly seen in PD, even in early phases of the disease [2, 5]. Moreover, it has previously been suggested that increased inhibition upon the SC leads to a compensatory increase of the frontal eye field activity resulting in SWJ in PD [5]. In another study [15], it was shown that the frequency of saccadic intrusions during the oculomotor examination was negatively correlated with brain volume in PD patients. Finally, in a study on PD patients that had undergone unilateral pallidotomy [16], increased number and amplitude of SWJ was reported during sustained fixation, possibly attributed to alterations in the function of frontal and prefrontal cortical areas following the disruption of pallidal influence on the thalamocortical loops. Such studies on fixation analysis highlight the importance of investigating ocular fixation in PD.

Although the participants needed to maintain attention during the task, fixation per se is not cognitively demanding to the same extent as other common eye movement tasks (e.g., antisaccades), thus, large fatigue effect was not expected. Our analysis has confirmed this assumption. If fatigue was driving the results, we would expect to see an increasing or decreasing trend over the trials in different parameters, but this was not observed.

Although an increase in saccadic intrusions and a decrease in fixation periods with age has been described in healthy populations with an age range between 21 and 81 years [17], we were not able to find such a correlation neither in the HC nor in the PD group. This might be attributed to the relatively narrower age range in our sample (42–77 years in the HC, 42–76 years in the PD group).

Regarding cognition, fixation and gaze parameters correlated weakly with MMSE but not with MoCA score in the PD group only, which may be attributed to the fact that all participants had good cognitive performance (i.e., high scores in both tests) combined with the ceiling effect of MMSE score [18]. MoCA has widely been recognized as a superior screening tool in PD as it includes executive function testing, while MMSE lacks this part [18].



Another interesting finding in our analysis was that both mean and median pupil size were significantly larger in the HC group than in the PD group. This poses significant questions on the role of the autonomous nervous system on the modulation of pupil size during cognitively demanding tasks in health and disease. The dilator and sphincter muscles of the pupil are controlled by the sympathetic and parasympathetic branches of the nervous system via neurotransmission that is done with catecholamines and acetylcholine, respectively. Sympathetic fibers innervate the dilator muscle evoking mydriasis, whereas inhibition of parasympathetic activity has the same effect by reducing constriction of the sphincter muscle [19]. However, the difference between the parasympathetic and sympathetic pathways is remarkable. While the parasympathetic pathway is mainly responsible for the luminance-based pupil constriction, and the circuit is short, in contrast, the sympathetic pathway driving mydriasis involves multiple cortical and subcortical domains such as the frontal cortex, the hypothalamus, the locus coeruleus as well as the spinal cord (C8-T2) [19]. Modulation of the norepinephrine and acetylcholine systems is done by cortical and subcortical areas that are involved in cognitive control and attention. Here lies the role of the locus-coeruleus norepinephrine (LC-NE) system that modulates mydriasis with respect to task demands [19–21] and affects the pupil both directly, increasing the sympathetic activity, and indirectly by sending inhibitory projections to the parasympathetic Edinger-Westphal nucleus that normally causes miosis. The complexity of the aforementioned pathway, as well as the functional role of the LC-NE in cognitive processes [21] explains how pupil dilation is related not only with luminance but also to cognitive factors such as attention, memory and cognitive load, prediction, decision, as well as autonomic activity. During the fixation paradigm, in stable light conditions, trying to focus on the target for almost three minutes with only short intervals of rest was demanding on an attentional level. Based on our knowledge of the physiological mechanisms that underlie pupil size regulation, pathology of the LC-NE in PD [22] is one possible mechanism that explains why PD patients' pupils were smaller than HC's during sustained fixation. However, apart from the role of pre-ganglionic structures of the autonomous nervous system, speculations can also be made on the role of the post-ganglionic innervation of the pupil based on previous descriptions of the post-ganglionic deficiency of the sympathetic

and parasympathetic system that affect pupil size modulation in PD [23].

It is important to consider the possible effect of concomitant medication and comorbidities that may have played an important role in our findings. Depression was, not surprisingly, more common in the PD group, although the difference did not reach statistical significance ( $p=0.07$ ). Treatment with antidepressants is expected to cause mydriasis as a side effect; our PD group, however, presented with smaller pupil size. Additionally, the use of benzodiazepines and anticholinergic medication that could have an effect on the pupil size was more common in the PD group, yet, the differences lacked statistical significance. Diabetes may affect the pupil size but was only present in one HC. Finally, regarding the differences in urinary incontinence and/or prostate hyperplasia, the finding, itself, can be explained by the fact that urinary incontinence can be part of the non-motor symptoms in PD and by the unequal sex distribution in the two groups. Six PD patients were treated with medication that could possibly affect pupil size (solifenacine, alfuzosin, doxazosin, finasteride) vs only one in the HC group, a difference, however, the difference was not statistically significant.

Based on previous literature that used pupillometry to investigate sex differences during cognitive effort for similar behavioral performance [24, 25], we included some additional comparisons between male and female participants with respect to pupil size during the fixation task. Our results are in accordance with previous studies that show that women present with larger pupils while putting cognitive effort [24] while other studies show no difference between sexes [25]. Results should therefore be interpreted with caution and might be attributed to multiple factors.

### Limitations

Our study has several limitations. Sex distribution differed significantly between HC and PD group; however, sex was included in the multivariate model in our analysis in order to adjust for this parameter. Also, the clinical evaluation of PD patients was performed only in the ON-medication state, thus preventing the investigation of the effect of levodopa treatment on saccade rate and ocular fixation characteristics. However, as an indirect indicator, we didn't find any significant correlations between the eye-movement parameters and the amount of levodopa equivalent intake. Finally, we were not able to precisely characterize the nature of the saccadic

intrusions that interrupted fixation. Although we assume, based on the literature, that they were SWJ, they could as well be reflexive saccades or microsaccades. However, in our study we aimed to study sustained fixation in a natural and ecologically valid setting where the participant is free to move. Studies that have investigated such movements have used a very rigid eye tracker set-up with chin- and forehead rest and sometimes even a bite-bar. It is unclear whether SWJ and microsaccades can be reliably detected in a set-up without head-restraint due to the potential confounding with small head movements.

## CONCLUSION

Eye-tracking is a quick and easy method to investigate brain function, with only slight, if any, discomfort to the participant. In this case-control study, eye-tracking based measurements of gaze fixation and pupil reaction could be used to discriminate PD patients with short disease duration from healthy, age-matched participants. Our study suggests that PD diagnosis at an early stage can be aided by using oculomotor tests such as fixation and pupil reaction as physiological biomarkers, although larger studies are needed in order to evaluate the generalizability of the results.

## ACKNOWLEDGMENTS

The study was supported by Vinnova, Sweden's innovation agency (Grant number: 2017-02317; Early detection of Alzheimer's and Parkinson's disease based on eye-tracking and AI).

We would like to express thanks to Vinnova and to the Sigvard and Marianne Bernadotte Research Foundation for Children's Eye Care.

## CONFLICT OF INTEREST

Benfatto, M. N. and Seimyr, G. own equity and receive salary from Optolexia, a company whose aim is to offer new technologies for the assessment of based on eye tracking and artificial intelligence. The venture is a result of projects funded by Sweden's innovation agency – VINNOVA – (2014-03459; 2017-02317) and Karolinska Institutet Innovations.

PS receives funding from Vinnova, the Stockholm County Council, and Wallenberg Clinical Scholarship, as well as honoraria from Abbvie.

IM receives funding from Stockholm County Council (20180200), Neuro Fund Stockholm and Parkinson Research Foundation, Stockholm

## REFERENCES

- [1] Ekker MS, Janssen S, Seppi K, Poewe W, de Vries NM, Theelen T, Nonnekes J, Bloem BR (2017) Ocular and visual disorders in Parkinson's disease: Common but frequently overlooked. *Parkinsonism Relat Disord* **40**, 1-10.
- [2] Gorges M, Pinkhardt EH, Kassubek J (2014) Alterations of eye movement control in neurodegenerative movement disorders. *J Ophthalmol* **2014**, 658243.
- [3] Krauzlis RJ, Goffart L, Hafed ZM (2017) Neuronal control of fixation and fixational eye movements. *Philos Trans R Soc Lond B Biol Sci* **372**.
- [4] Martinez-Conde S, Macknik SL, Hubel DH (2004) The role of fixational eye movements in visual perception. *Nat Rev Neurosci* **5**, 229-240.
- [5] Shaikh AG, Xu-Wilson M, Grill S, Zee DS (2011) 'Staircase' square-wave jerks in early Parkinson's disease. *Br J Ophthalmol* **95**, 705-709.
- [6] Wang CA, Munoz DP (2015) A circuit for pupil orienting responses: Implications for cognitive modulation of pupil size. *Curr Opin Neurobiol* **33**, 134-140.
- [7] Micieli G, Tassorelli C, Martignoni E, Pacchetti C, Bruggi P, Magri M, Nappi G (1991) Disordered pupil reactivity in Parkinson's disease. *Clin Auton Res* **1**, 55-58.
- [8] Granholm E, Morris S, Galasko D, Shults C, Rogers E, Vukob B (2003) Tropicamide effects on pupil size and pupillary light reflexes in Alzheimer's and Parkinson's disease. *Int J Psychophysiol* **47**, 95-115.
- [9] Hughes AJ, Daniel SE, Kilford L, Lees AJ (1992) Accuracy of clinical diagnosis of idiopathic Parkinson's disease: A clinico-pathological study of 100 cases. *J Neurol Neurosurg Psychiatry* **55**, 181-184.
- [10] Fahn S, Elton RL, UPDRS program members (1987) Unified Parkinsons Disease Rating Scale. In *Recent developments in Parkinsons disease*, vol 2, Fahn S, Marsden CD, Goldstein M, Calne DB, eds. Macmillan Healthcare Information, Florham Park, NJ, pp. 153-163.
- [11] Castet E, Crossland M (2012) Quantifying eye stability during a fixation task: A review of definitions and methods. *Seeing Perceiving* **25**, 449-469.
- [12] Gorges M, Muller HP, Lule D, Pinkhardt EH, Ludolph AC, Kassubek J (2016) The association between alterations of eye movement control and cerebral intrinsic functional connectivity in Parkinson's disease. *Brain Imaging Behav* **10**, 79-91.
- [13] Pinnock RA, McGivern RC, Forbes R, Gibson JM (2010) An exploration of ocular fixation in Parkinson's disease, multiple system atrophy and progressive supranuclear palsy. *J Neurol* **257**, 533-539.
- [14] Pretegianni E, Optican LM (2017) Eye movements in Parkinson's disease and inherited parkinsonian syndromes. *Front Neurol* **8**, 592.
- [15] Vintonyak O, Gorges M, Muller HP, Pinkhardt EH, Ludolph AC, Huppertz HJ, Kassubek J (2017) Patterns of eye movement impairment correlate with regional brain atrophy in neurodegenerative parkinsonism. *Neurodegener Dis* **17**, 117-126.
- [16] O'Sullivan JD, Maruff P, Tyler P, Peppard RF, McNeill P, Currie J (2003) Unilateral pallidotomy for Parkinson's

- 674 disease disrupts ocular fixation. *J Clin Neurosci* **10**,  
675 181-185.
- 676 [17] Canice McGivern R, Mark Gibson J (2006) Characterisation  
677 of ocular fixation in humans by analysis of saccadic intru-  
678 sions and fixation periods: A pragmatic approach. *Vision*  
679 *Res* **46**, 3741-3747.
- 680 [18] Chou KL, Lenhart A, Koeppel RA, Bohnen NI (2014) Abnor-  
681 mal MoCA and normal range MMSE scores in Parkinson  
682 disease without dementia: Cognitive and neurochemical  
683 correlates. *Parkinsonism Relat Disord* **20**, 1076-1080.
- 684 [19] Ajasse S, Benosman RB, Lorenceau J (2018) Effects of  
685 pupillary responses to luminance and attention on visual  
686 spatial discrimination. *J Vis* **18**, 6.
- 687 [20] van der Wel P, van Steenbergen H (2018) Pupil dilation as an  
688 index of effort in cognitive control tasks: A review. *Psychon*  
689 *Bull Rev* **25**, 2005-2015.
- 690 [21] Eckstein MK, Guerra-Carrillo B, Miller Singley AT, Bunge  
691 SA (2017) Beyond eye gaze: What else can eyetracking  
692 reveal about cognition and cognitive development? *Dev*  
693 *Cogn Neurosci* **25**, 69-91.
- [22] Bari BA, Chokshi V, Schmidt K (2020) Locus coeruleus-  
norepinephrine: Basic functions and insights into Parkin-  
son's disease. *Neural Regen Res* **15**, 1006-1013.
- [23] Hori N, Takamori M, Hirayama M, Watanabe H, Nakamura  
T, Yamashita F, Ito H, Mabuchi N, Sobue G (2008) Pupillary  
supersensitivity and visual disturbance in Parkinson's  
disease. *Clin Auton Res* **18**, 20-27.
- [24] Campbell MJ, Toth AJ, Brady N (2018) Illuminating sex dif-  
ferences in mental rotation using pupillometry. *Biol Psychol*  
**138**, 19-26.
- [25] Toth AJ, Campbell MJ (2019) Investigating sex differences,  
cognitive effort, strategy, and performance on a comput-  
erised version of the mental rotations test via eye tracking.  
*Sci Rep* **9**, 19430.

694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707

Uncorrected Author Proof