

Erratum

Otolith-ocular responses in patients with surgically confirmed unilateral peripheral vestibular loss

by Joseph M. Furman, Li-Chi Hsu, Susan L. Whitney and Mark S. Redfern

[*Journal of Vestibular Research* **13** (2,3) (2003), pp. 143–151]

Table 2B in this article is incorrect. The correct and reorganized Table 2B is given below:

Table 2
Semicircular canal-ocular response parameters in patients with unilateral peripheral vestibular loss

B. Responses to EVAR Sinusoids									
	0.02 Hz /50°/sec EVAR			0.05 Hz /50°/sec EVAR			0.1 Hz /25°/sec EVAR		
	Gain	Phase (deg)	DC Offset (deg/sec)	Gain	Phase (deg)	DC Offset (deg/sec)	Gain	Phase (deg)	DC Offset (deg/sec)
Patients	0.24 ± 0.09	49.2 ± 9.0	2.2 ± 2.0	0.35 ± 0.12	27.3 ± 9.0	2.7 ± 2.3	0.37 ± 0.10	19.4 ± 7.0	1.9 ± 1.4
Controls	0.44 ± 0.14	25.4 ± 9.7	0.6 ± 1.3	0.49 ± 0.21	12.0 ± 7.9	0.9 ± 1.2	0.54 ± 0.14	9.1 ± 8.1	-0.9 ± 1.6
	0.1 Hz /50°/sec EVAR			0.1 Hz /100°/sec EVAR			0.1 Hz /150°/sec EVAR		
	Gain	Phase (deg)	DC Offset (deg/sec)	Gain	Phase (deg)	DC Offset (deg/sec)	Gain	Phase (deg)	DC Offset (deg/sec)
Patients	0.40 ± 0.14	18.5 ± 7.2	2.7 ± 2.3	0.36 ± 0.10	15.3 ± 8.1	4.0 ± 3.1	0.33 ± 0.13	16.2 ± 5.3	7.4 ± 4.2
Controls	0.51 ± 0.17	7.0 ± 5.0	0.1 ± 1.7	0.50 ± 0.18	6.4 ± 4.7	0.3 ± 3.1	0.51 ± 0.16	8.3 ± 5.3	0.1 ± 3.0
	0.5 Hz /50°/sec EVAR								
	Gain	Phase (deg)	DC Offset (deg/sec)						
Patients	0.54 ± 0.14	8.5 ± 6.2	3.5 ± 2.1						
Controls	0.63 ± 0.23	-0.3 ± 5.0	-0.1 ± 2.3						