

## Review

# Double Immune Checkpoint Blockade in Renal Cell Carcinoma

Luyo M<sup>a</sup>, Carril-Ajuria L<sup>a</sup>, Fabio Schutz F<sup>b</sup>, Castellano D<sup>a</sup> and De Velasco G<sup>a,\*</sup>

<sup>a</sup>*Department of Medical Oncology, University Hospital 12 de Octubre, Madrid, Spain*

<sup>b</sup>*Department of Medical Oncology, Hospital São José, Beneficiencia Portuguesa de São Paulo, Brazil*

**Abstract.** Long considered an immunogenic tumour, immunotherapy has been the cornerstone of systemic treatment in renal cell carcinoma (RCC) for decades. Since the introduction of interleukin 2 and interferon-alfa in the 80s to the more recently approved nivolumab (anti-PD-1) in second line setting. Moreover, on the basis that anti-CTLA-4 and anti-PD-1/PD-L1 intrinsic mechanisms are different, double checkpoint inhibition was proposed to further improve anti-tumor immune response. The first trial to assess double checkpoint inhibition was the Checkmate 016 (nivolumab and ipilimumab), showing acceptable safety and promising antitumor activity that led to the first phase III trial with combination immunotherapy in RCC, the Checkmate 214. This trial showed superior overall survival and response rate of the combination immunotherapy (nivolumab and ipilimumab) versus sunitinib in intermediate- and poor- risk advanced RCC, leading to its approval in this setting. Despite these advances, there is still room for improvement. In this sense, cytokines and T-cell costimulatory molecules are currently under investigation. This review summarizes the principles of immunotherapy and its role in RCC, provides an update on double checkpoint blockade and finally discusses the major challenges with double checkpoint blockade.

**Keywords:** Immune checkpoint inhibitors, double immune checkpoint blockade, renal cell carcinoma, review, immunotherapy, clinical trials

## RATIONAL FOR IMMUNOTHERAPY

Three steps need to be achieved in order to yield antitumor immunity: 1) immunization which implies antigen presentation and dendritic cell maturation, 2) T-cell response which requires interaction between stimulatory and inhibitory molecules and surface proteins, and 3) immunosuppression that implicates the tumor microenvironment strategies that alter T cell effects through suppressive ligands (such as PD-L1,PD-L2) or molecules (indoleamine 2,3-dyoxigenase-IDO) among others [1, 2].

T-cell response regulation has revolutionized the targets in cancer. By removing known inhibitory pathways, T-cell response can be let freely to act. One of the effectors of this carefully delineated balance are the cytotoxic T-cells surface proteins that allow for immune response restrain, known as checkpoint proteins [3]. On nature they display a wide variety, but so far only two have been well described in literature. Cytotoxic T lymphocyte-associated antigen-4 (CTLA-4) and Programmed cell death protein (PD-1) are surface proteins that undergo upregulation after T-cells are activated [4, 5]. These inhibitory checkpoint proteins exemplify different mechanisms and pathways but ultimately own the same objective: lighten and improve antitumor immune response [6, 7].

T-cell activation requires interaction between CD28 on the T-cell surface and B7 on the

\*Correspondence to: Guillermo de Velasco MD, PhD University Hospital 12 de Octubre, Department of Medical Oncology, Av Cordoba s/n 28041, Madrid, Spain. E-mail: Guillermoantonio.velasco@salud.madrid.org.

antigen-presenting cell surface. CTLA-4 displaces CD28 from B7 which ensues T-cell suppression and at the same time enhances T regulatory cells activation and suppresses helper T-cells [4, 8]. *Mellman et al.* suggests that the explanation for therapy at this point needs previous tumor-protective T-cells that would become activated after CTLA-4 blocking [1]. In contrast, PD-1 and its ligands (PD-L1 and PD-L2), members of the B7 gene family, operate through a different inhibitory pathway [9]. These transmembrane proteins, found in T- and B-cells, are responsible for activation of T regulatory cells through de novo conversion of CD4+ T-cells and inhibition of interferon- $\gamma$ , tumor necrosis factor- $\alpha$  and IL-2 production allowing for immunosuppressive function. Furthermore, if PD-1 is expressed persistently on T-cells an exhaustive phenotype emerges depicting diminished immune function [10]. PD-1 ligands (PD-L1 and PD-L2) are not only expressed by antigen presenting cells but many non-immune cells and even tumor cells [5, 11]. Thus, they present a diversity of ligands different in nature, function and implications on the therapeutic approach.

The well-balanced immune structure can be hijacked by the tumor microenvironment so it can develop, progress and prevail in the organism. This understanding introduced the concepts of immune-evasion and tumor-promoting inflammation among cancer hallmarks [12, 13]. Immune checkpoint blockade with an anti-CTLA-4 and anti-PD-1/PD-L1 represent part of these advances in several immunogenic cancer types such as melanoma, non-small-cell lung cancer (NSCLC) and renal cell carcinoma (RCC), among others [14].

Immunotherapy has been used in kidney cancer for decades [15, 16]. RCC has been considered as an immunogenic tumor because of the increased infiltration of immune cells including T-cells, dendritic cells, natural killer T-cells, macrophages and memory cells, and also increased cytokine secretion. In the 80s and 90s, advanced RCC immunotherapy was based on high-dose interleukin-2 (IL-2) and interferon- $\alpha$  (IFN $\alpha$ ) which elicited a high burden of toxicities and yield low efficacy. More recently, Nivolumab, an anti-PD-1 monoclonal antibody, was the first immune checkpoint inhibitor approved by the FDA in 2015 based on results of the CheckMate 025 study [16–18].

In addition, based on the core concept that anti-CTLA-4 and anti-PD-1/PD-L1 intrinsic mechanisms are different, double checkpoint inhibition was proposed to further improve anti-tumor immune

response. Anti-CTLA-4 acts in lymphoid organs in early steps of immune response. In contrast, anti-PD-1/PD-L1 acts in the tumor microenvironment later in the response. In this review we will summarize the current evidence of double immune checkpoint blockade in renal-cell carcinoma (RCC).

#### *Double immune checkpoint blockade*

Simultaneous pathway blockade and sequential blockade are the backbone of combination therapy in kidney cancer. The association of both immune checkpoint inhibitors in a preclinical setting is more effective disrupting the tumor microenvironment [19]. In RCC, the concept has been proven to be clinically effective within the Checkmate-016 and Checkmate-214 trials.

### **IPILIMUMAB-NIVOLUMAB COMBINATION IN ADVANCED RENAL CELL CARCINOMA**

#### *Checkmate-016*

The first study to assess the combination of immune checkpoint inhibitors (ICI) in mRCC was the Checkmate-016 study (NCT01472081) [20]. This phase I trial was based on previous results on patients with melanoma and lung cancer where double immune checkpoint blockade was more effective than monotherapy and also on efficacy evidence of anti-CTLA-4 antibodies in mRCC [14, 21]. Its primary endpoint was the overall safety and tolerability of nivolumab plus ipilimumab. Eligible patients were divided in 3 groups that received nivolumab 3mg/kg plus ipilimumab 1mg/kg (N3I1), nivolumab 1mg/kg plus ipilimumab 3mg/kg (N1I3) or nivolumab 3mg/kg plus ipilimumab 3mg/kg (N3I3) every 3 weeks for four cycles and then only nivolumab 3mg/kg every 2 weeks until progression or toxicity. Accrual on N3I3 group was discontinued due to the observed high-grade 3 or 4 adverse effects (AEs) (83.3%). On the other hand, high-grade AEs in N3I1 and N1I3 treatment groups were 38.3% and 61.7%, respectively. N3I1 treatment group had also lower treatment-related AEs, lower treatment discontinuation rates and less patients requiring immune modulating medications (61.7% vs 83.0%). After a median follow-up of 22.3 months, the objective response rate (ORR) for both treatment-naive arms was 40% and the progressive disease (PD) rate as best

Table 1  
Efficacy outcomes in the Checkmate-016 and the Checkmate-214

Efficacy	Checkmate 016 (N3I1)	Checkmate 214
ORR	40% (95% CI 26.4 to 55.7)	42% (95% CI, 37 to 47)
Overall Survival	not reached (95% CI, 26.7 months to not reached)	not reached (95% CI, 28.2 months to not estimable)
12 -month OS	81%	80% (95CI, 76 to 84)
PFS	7 months (95% CI, 3.7 to 14.3 months)	11.6 months (95% CI, 8.7 to 15.5)

response was 17%. Complete response (CR) and partial response (PR) rates in N3I1 treated patients were 10.6% and 29.8%, respectively. In comparison, there were no CR in N1I3 treated patients and the PR rate was 40.4%. Safety profile favoured N3I1 treatment group, therefore it was selected for further development in a phase III trial.

#### Checkmate-214

Ipilimumab plus nivolumab combination has not been compared with either nivolumab or ipilimumab alone in mRCC. However, results from Checkmate-016 led to the conduct of a phase III trial with the combination. In the Checkmate 214, 1096 patients with previously untreated mRCC were randomized 1:1 to receive nivolumab 3mg/kg plus ipilimumab 1mg/kg every 3 weeks for four doses, then only nivolumab 3 mg/kg every 2 weeks or sunitinib 50mg orally daily for 4 weeks on followed by 2 weeks off. Primary endpoints were overall survival (OS), ORR and progression-free survival (PFS) in the intermediate and poor IMDC risk-groups. After a median of 25.2 months, there was a significant improvement in 18-month OS (75% with the combination versus 60% with sunitinib) and ORR (42% vs 27%, respectively,  $P < 0.001$ ), favouring nivolumab plus ipilimumab combination [22]. The median OS was not reached (NR) and 26 months for immunotherapy combination and sunitinib, respectively. Of note, the CR rate was 9% and 1%, respectively, and the median PFS was 11.6 and 8.4 months, respectively (HR 0.82,  $p = 0.03$ ). Similar benefit was seen when evaluating the whole cohort including all IMDC risk-groups (good, intermediate and poor) (secondary endpoint). The safety profile showed any-grade and high-grade AEs of 93% and 46%, respectively, in the immunotherapy combination arm, and 97% and 63%, respectively, in the sunitinib arm. Interestingly, although the AE-related treatment discontinuation rate was superior in the ICI combination arm (22%) versus the sunitinib arm (12%), patients who discontinued in the experimental arm due to drug-related AEs experienced long-term

benefit, with durable responses after the last dose of treatment. Moreover, the recently reported PROs (patient reported outcomes) as an exploratory endpoint, showed an improve of the health-related quality of life (HRQOL) across all instruments (Assessment of Cancer Therapy Kidney Symptom Index-19, Functional Assessment of Cancer Therapy-General, and EuroQol five dimensional three level) for the immunotherapy combination arm in patients with intermediate or poor risk mRCC [23]. Efficacy in the good IMDC risk-group only was evaluated as an exploratory outcome. In this good-risk group, sunitinib treated patients had higher 18-month OS, ORR and PF. However, it should be mentioned that the CR rate in good-risk patients treated was higher with the immunotherapy combination (11% versus 5%). Also as an exploratory analysis, among intermediate and high risk patients who had quantifiable PD-L1 expression ( $< \text{or} \geq 1\%$ ) the combination immunotherapy arm showed superiority in both OS and ORR irrespective of PD-L1 expression versus the control arm. However, mPFS among patients with  $< 1\%$  PD-L1 expression was similar in both arms of treatment (HR 1.00; 95% CI, 0.80 to 1.26), whereas among patients with  $\geq 1\%$  PD-L1 expression mPFS was superior with the immunotherapy combination (HR 0.46; 95% CI, 0.31–0.67) [22]. On April 16, 2018, the FDA approved the double immune checkpoint blockade, nivolumab plus ipilimumab, for the treatment of intermediate or poor risk, previously untreated advanced RCC patients.

#### OTHER COMBINATIONS

Despite the remarkable progress that has been achieved with ipilimumab and nivolumab combination, there is still room for improvement in efficacy and safety profile. It remains unclear whether the enhanced efficacy of combined anti PD-1 and anti-CTLA4 therapy is mediated by additive engagement or through independent mechanisms from each drug [24]. T-cell costimulatory molecules represent a vast number of proteins under investigations

Table 2  
Safety outcomes in the Checkmate-016 and the Checkmate-214

Treatment related adverse effects.	Checkmate 016	Checkmate 214
N	47	547
Median Age (range)	54(26–68)	62(26–85)
MSKCC/ IMDC prognosis risk intermediate-poor	55.30%	77.70%
Treatment Arm	Nivo 3mg/kg+Ipi 1mg/kg	Nivo 3mg/kg+Ipi 1mg/kg
Any grade	92%	93%
Grade 3-4 toxicity	38.30%	46%
Discontinuation rate for toxicity	10.60%	22%

such as LAG3, TIM3 or TIGIT, VISTA from the immunoglobulin superfamily (IgSF) or OX40, GITR, 4-1BB, CD40 from the tumor necrosis factor receptor superfamily (TNFRSF) and others. Better understanding of each specific protein and its stimulatory role are required for an enhanced development of novel immune checkpoint blockade combinations [25]. Table 3 shows selected combinations in early clinical development phase.

Cytokines were the main treatment for mRCC during decades; however the rate of response was low. A novel C122-biased cytokine (NTRK-214) is currently under development in combination with nivolumab and ipilimumab in the phase I/II PIVOT-02 trial. It is expected that NKTR-214 may unleash T cell proliferation and disrupt the tumor microenvironment. The combination of NKTR-214 and nivolumab assures a potentially non-overlapping effective synergistic mechanism. This study was conceived as a 4 part study (NCT02983045). Study part 1 determined doses and scheduling according to safety and efficacy profile of the combination. The selected dose and schedule was NKTR-214 0.006 mg/kg and nivolumab 360 mg every 3 weeks per intravenous infusion. In study Part 2, the combination was given to patients with metastatic melanoma, RCC, urothelial carcinoma, non-small cell lung cancer or triple negative breast cancer. The RCC cohort enrolled 48 patients of which PD-L1 status was unknown for 4 patients, 30 had a negative status, and 14 had a positive status. Preliminary results showed in previously untreated RCC patients an overall ORR of 46%. The median time on study was 5.6 months. The PD-L1 negative patients ORR was 53% (9/17), while it was 29% (2/7) for PD-L1 positive patients [26]. In the study part 3, a triple combination will be attempted in order to assess safety, tolerability and efficacy (ORR): NKTR-214 0.006 mg/kg plus nivolumab 360mg, every 3 weeks, and ipilimumab 1mg/kg every 6 weeks. Study part 4 will plan to enrol more patients to further evaluate the triplet combination.

## KEY POINTS AND CHALLENGES WITH DOUBLE IMMUNE CHECKPOINT BLOCKADE

### *Why double and not single-agent?*

Double immune checkpoint blockade has been shown to be more powerful than single-agent immunotherapy across different tumour types, but also more toxic [27]. However, the superiority of immunotherapy doublets versus single agents has not yet been demonstrated for mRCC. It is known that nivolumab alone benefits previously treated metastatic renal cell carcinoma patients [22, 28], but we do not know the magnitude of the benefit with the addition of ipilimumab. Recent data with pembrolizumab monotherapy in untreated mRCC patients has shown encouraging activity [29], with a confirmed ORR of 38.2% ( $n=42$ ; 95%CI 29.1–47.9), with 3 complete responses (2.7%) and 39 (35.5%) partial responses. The durable clinical response rate was 59.1 with a favourable safety profile. It seems clear that the addition of ipilimumab is more toxic and some patients may not need the combinations [14]. Further research to better understand which patients do really benefit the combination is required to avoid toxicities (financial and non-financial). Clinical trials with adaptive design such as TITAN (NCT02917772) or OMNIVORE (NCT03203473) will help to understand this issue.

### *What is the optimal duration of treatment?*

There is evidence showing that patients who respond to immunotherapy and combination immunotherapy achieve durable responses and long-term survival [30]. Although immune checkpoint blockade is already a standard of care across different tumour types, there is no clear evidence about the duration of treatment on responders. In recent data from the Keynote-006 4-year survival and outcomes after cessation of pembrolizumab

Table 3  
Summary of Selected Studies of Double Immune checkpoint blockade on going

Phase	Trial	Year	Treatment	Title
Phase IIIb/IV	NCT02982954 Checkmate 920	January 2023	Nivolumab plus Ipilimumab	Safety Trial of Nivolumab Combined With Ipilimumab in Subjects With Previously Untreated, Advanced or Metastatic RCC
Phase III	NCT03138512 Checkmate 914	July 2023	Nivolumab plus Ipilimumab	A Phase 3 Randomized Study Comparing Nivolumab and Ipilimumab Combination vs Placebo in Participants with Localized Renal Cell Carcinoma Who Underwent Radical or Partial Nephrectomy and Who Are at High Risk of Relapse
	NCT03288532 RAMPART	December 2037	Active Monitoring or Durvalumab or Durvalumab plus Tremelimumab	An International Investigator-led Phase III Multi Arm Multi Stage Multi-centre Randomised Controlled Platform Trial of Adjuvant Therapy in Patients With Resected Primary Renal Cell Carcinoma (RCC) at High or Intermediate Risk of Relapse
	NCT03260894 Keynote-679/Echo-302	June 2020	Pembrolizumab plus Epacadostat or Sunitinib or Pazopanib	A Randomized, Open-Label, Phase 3 Study to Evaluate Efficacy and Safety of Pembrolizumab (MK-3475) Plus Epacadostat vs Standard of Care (Sunitinib or Pazopanib) as First-Line Treatment for Locally Advanced or Metastatic Renal Cell Carcinoma (mRCC)
Phase II	NCT03203473 OMNIVORE Study	November 2014	Nivolumab or Nivolumab plus Ipilimumab	Phase II Study of Optimized Management of Nivolumab Based on REsponse in Patients With Advanced Renal Cell Carcinoma
Phase I/II	NCT02335918	January 2020	Nivolumab plus Varlilumab	A Phase I/II Dose Escalation and Cohort Expansion Study of the Safety, Tolerability and Efficacy of Anti-CD27 Antibody (Varlilumab) Administered in Combination With Anti-PD-1 (Nivolumab) in Advanced Refractory Solid Tumors
	NCT02089685 KEYNOTE-029	June 2020	Pembrolizumab plus Pegylated Interferon alfa-2b or Pembrolizumab plus Ipilimumab	Pembrolizumab Plus Pegylated Interferon alfa- 2b or Ipilimumab for Advanced Melanoma or Renal Cell Carcinoma: Dose-Finding Results from the Phase Ib Study
	NCT02983045 PIVOT-02	June 2021	Phase I: NKTR-214 + nivolumab Phase II: NKTR-214 + nivolumab+ipilimumab	A Dose Escalation and Cohort Expansion Study of CD122-Biased Cytokine (NKTR-214) in Combination With Anti-PD-1 Antibody (Nivolumab) or in Combination With Nivolumab and Anti-CTLA4 Antibody (Ipilimumab) in Patients With Select Advanced or Metastatic Solid Tumors

(Continued)

Table 3  
(Continued)

Phase	Trial	Year	Treatment	Title
Phase I	NCT02496208	September 2019	Cabozantinib plus Nivolumab or Cabozantinib plus Nivolumab plus Ipilimumab	A Phase 1 Study of Cabozantinib Plus Nivolumab (CaboNivo) Alone or in Combination With Ipilimumab (CaboNivoIpi) in Patients With Advanced/Metastatic Urothelial Carcinoma and Other Genitourinary Tumors
	NCT01975831	December 2019	Durvalumab plus Tremelimumab	Phase 1 study to evaluate the safety and tolerability of MEDI4736 (durvalumab, DUR)+tremelimumab (TRE) in patients with advanced solid tumors.
	NCT02210117	November 2020	Arm A: Nivolumab or Arm B: Nivolumab plus Bevacizumab or Arm C: Nivolumab plus Ipilimumab	A Pilot Randomized Tissue-Based Study Evaluating Anti-PD1 Antibody or Anti-PD1 + Bevacizumab or Anti-PD1 + Anti-CTLA-4 in Patients With Metastatic Renal Cell Carcinoma Who Are Eligible for Cytoreductive Nephrectomy, Metastasectomy or Post-Treatment Biopsy
	NCT02614456	December 2021	Nivolumab plus Interferon-gamma	Combination Immunotherapy With Interferon- gamma and Nivolumab for Patients With Advanced Solid Tumors: A Phase 1 Study

showed that 86% of the patients who completed pembrolizumab 2 years remained progression-free 20 months after the last dose of treatment [31]. Also, it has been shown that nearly 70% of the melanoma patients who discontinued single-agent nivolumab due to toxicity and that obtained a complete or partial response maintained their responses despite the stop of treatment [30, 32]. Moreover, this may be even more complex when it comes to patients who achieve a complete response. The clinical trial Keynote-001 showed that a high proportion of patients who achieved a complete response are able to maintain the response for a long time after the end of treatment, with an estimated 24 month disease-free survival of 89% [31]. Altogether, these results suggest stopping immune checkpoint blockade after a certain period of time on treatment could be a reasonable approach. However, further research on this topic is warranted.

#### *How should we assess atypical responses?*

Although there is lack of evidence, we are now familiar with the so known “pseudoprogression” which usually appears as a radiological progres-

sion with target lesion increase but without clinical deterioration. There are data suggesting that treatment beyond progression may be an option for some patients [33, 34]. However, there is also evidence supporting that some patients may have hyperprogressive disease, but little is known about the mechanisms involved [35]. It has been speculated to be a result of immunotherapy but it has also been suggested to be the consequence of aggressive natural tumour behaviour. Nonetheless, knowledge on this field is scarce and further studies are needed to better understand and define which patients are more likely to rapidly progress and which ones will instead experience pseudoprogression and therefore benefit of treatment beyond radiological progression [36].

## CONCLUSIONS

Several trials will be conducted in the next years that will elucidate future combination’s impact on RCC. In a near future, the patient’s signature tumor biology should be matched with the best corresponding first line option among a handful of options: either monotherapy or combinations. Double immune checkpoint blockade has shown favourable

353 results but adverse effects should be taken into con-  
 354 sideration. The most definitive evidence of double  
 355 checkpoint inhibitor therapy on intermediate-poor  
 356 risk patients with previously untreated advanced RCC  
 357 are the results from Checkmate trial 214 with a  
 358 remarkable 9% complete response rate.

359 Although we are now familiar with immune check-  
 360 point blockade therapies, there are still some open  
 361 questions such as optimal duration of treatment or  
 362 unexpected radiological responses, including pri-  
 363 mary progressions, that require further study.

## 364 REFERENCES

- 365 [1] Mellman I, Coukos G, Dranoff G. Cancer immunotherapy  
 366 comes of age. *Nature*. 2011 Dec;480(7378):480-9.  
 367 [2] Chen DS, Mellman I. Elements of cancer immunity and the  
 368 cancer-immune set point. *Nature*. 2017 Jan;541(7637):321-  
 369 30.  
 370 [3] Chikuma S, Kanamori M, Mise-Omata S, Yoshimura A.  
 371 SOCS: Potential immune checkpoint molecules for cancer  
 372 immunotherapy. *Cancer Sci*. 2017.  
 373 [4] Azuma M, Ito D, Yagita H, Okumura K, Phillips JH, Lanier  
 374 LL, et al. B70 antigen is a second ligand for CTLA-4 and  
 375 CD28. *Nature*. 1993;366(6450):76-9.  
 376 [5] Butte MJ, Peña-Cruz V, Kim M-J, Freeman GJ, Sharpe AH.  
 377 Interaction of human PD-L1 and B7-1. *Molecular Immunol-  
 378 ogy*. 2008;45(13):3567-72.  
 379 [6] Sharma P, Allison JP. The future of immune checkpoint  
 380 therapy. *Science*. 2015 Apr 3;348(6230):56-61.  
 381 [7] Page DB, Postow MA, Callahan MK, Allison JP, Wolchok  
 382 JD. Immune modulation in cancer with antibodies. *Annual  
 383 Review of Medicine*. 2014;65(1):185-202.  
 384 [8] Snyder A, Makarov V, Merghoub T, Yuan J, Zaretsky JM,  
 385 Desrichard A, et al. Genetic basis for clinical response  
 386 to CTLA-4 blockade in melanoma. *N Engl J Med*. 2014;  
 387 371(23):2189-99.  
 388 [9] Ohaegbulam KC, Assal A, Lazar-Molnar E, Yao Y, Zang  
 389 X. Human cancer immunotherapy with antibodies to the  
 390 PD-1 and PD-L1 pathway. *Trends in Molecular Medicine*.  
 391 2015;21(1):24-33.  
 392 [10] Keir ME, Butte MJ, Freeman GJ, Sharpe AH. PD-1 and  
 393 its ligands in tolerance and immunity. *Annual Review of  
 394 Immunology*. 2008;26(1):677-704.  
 395 [11] Buchbinder EI, Desai A. CTLA-4 and PD-1 pathways: Sim-  
 396 ilarities, differences, and implications of their inhibition.  
 397 *American Journal of Clinical Oncology*. 2016;39(1):98-  
 398 106.  
 399 [12] Hanahan D, Weinberg RA. Hallmarks of cancer: The next  
 400 generation. *Cell*. 2011;144(5):646-74.  
 401 [13] Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell*.  
 402 2000;100(1):57-70.  
 403 [14] Larkin J, Chiarion-Sileni V, Gonzalez R, Grob JJ, Cowey  
 404 CL, Lao CD, et al. Combined nivolumab and ipilimumab or  
 405 monotherapy in untreated melanoma. *New England Journal  
 406 of Medicine*. 2015;373(1):23-34.  
 407 [15] Adler A, Gillon G, Lurie H, Shaham J, Loven D, Shachter Y,  
 408 et al. Active specific immunotherapy of renal cell carcinoma  
 409 patients: A prospective randomized study of immuno-  
 immunology and clinical aspects. *J Biol Response Mod*.  
 1987;6(6):610-24.  
 411 [16] Fyfe G, Fisher RI, Rosenberg SA, Sznol M, Parkinson  
 412 DR, Louie AC. Results of treatment of 255 patients with  
 413 metastatic renal cell carcinoma who received high-dose  
 414 recombinant interleukin-2 therapy. *J Clin Oncol*. 1995;  
 415 13(3):688-96.  
 416 [17] McDermott DF, Regan MM, Clark JI, Flaherty LE, Weiss  
 417 GR, Logan TF, et al. Randomized phase III trial of high-  
 418 dose interleukin-2 versus subcutaneous interleukin-2 and  
 419 interferon in patients with metastatic renal cell carcinoma.  
 420 *J Clin Oncol*. 2005;23(1):133-41.  
 421 [18] Motzer RJ, Escudier B, McDermott DF, George S, Ham-  
 422 mers HJ, Srinivas S, et al. Nivolumab versus everolimus  
 423 in advanced renal-cell carcinoma. *N Engl J Med*.  
 424 2015;373(19):1803-13.  
 425 [19] Curran MA, Montalvo W, Yagita H, Allison JP. PD-1 and  
 426 CTLA-4 combination blockade expands infiltrating T cells  
 427 and reduces regulatory T and myeloid cells within B16  
 428 melanoma tumors. *Proceedings of the National Academy  
 429 of Sciences*. 2010;107(9):4275-80.  
 430 [20] Hammers HJ, Plimack ER, Infante JR, Rini BI, McDermott  
 431 DF, Lewis LD, et al. Safety and efficacy of nivolumab  
 432 in combination with ipilimumab in metastatic renal cell  
 433 carcinoma: The checkMate 016 study. *Journal of Clinical  
 434 Oncology*. 2017;35(34):3851-8.  
 435 [21] Callahan MK, Kluger H, Postow MA, Segal NH, Lesokhin  
 436 A, Atkins MB, et al. Nivolumab plus ipilimumab in patients  
 437 with advanced melanoma: Updated survival, response, and  
 438 safety data in a phase I dose-escalation study. *J Clin Oncol*.  
 439 2018;36(4):391-8.  
 440 [22] Motzer RJ, Tannir NM, McDermott DF, Arén Frontera O,  
 441 Melichar B, Choueiri TK, et al. Nivolumab plus ipilimumab  
 442 versus sunitinib in advanced renal-cell carcinoma. *N Engl J  
 443 Med*. 2018;378(14):1277-90.  
 444 [23] Cella D, Grünwald V, Escudier B, Hammers HJ, George  
 445 S, Nathan P, et al. Patient-reported outcomes of patients  
 446 with advanced renal cell carcinoma treated with nivolumab  
 447 plus ipilimumab versus sunitinib (CheckMate 214):  
 448 A randomised, phase 3 trial. *The Lancet Oncology*.  
 449 2019;20(2):297-310.  
 450 [24] Wei SC, Duffy CR, Allison JP. Fundamental mechanisms  
 451 of immune checkpoint blockade therapy. *Cancer Discovery*.  
 452 2018;8(9):1069-86.  
 453 [25] Marin-Acevedo JA, Dholaria B, Soyano AE, Knutson KL,  
 454 Chumsri S, Lou Y. Next generation of immune check-  
 455 point therapy in cancer: New developments and challenges.  
 456 *Journal of Hematology & Oncology [Internet]*. 2018 Dec  
 457 [cited 2019 Jan 15];11(1). Available from: [https://jhoonline.  
 458 biomedcentral.com/articles/10.1186/s13045-018-0582-8](https://jhoonline.biomedcentral.com/articles/10.1186/s13045-018-0582-8).  
 459 [26] Diab A. Pivotal-02: Preliminary safety, efficacy and  
 460 biomarker results from dose escalation of the Phase 1/2  
 461 study of CD-122-biased agonist NKTR-214 plus nivolumab  
 462 in patients with locally advanced/metastatic melanoma,  
 463 renal cell carcinoma and non-small cell lung cancer. In  
 464 National Harbor, MD; 2017.  
 465 [27] Errico A. Melanoma: CheckMate 067—frontline  
 466 nivolumab improves PFS alone or in combination with ipil-  
 467 imumab. *Nature Reviews Clinical Oncology*. 2015;12(8):  
 468 435-5.  
 469 [28] Escudier B, Sharma P, McDermott DF, George S, Ham-  
 470 mers HJ, Srinivas S, et al. CheckMate 025 randomized phase  
 471 3 study: Outcomes by key baseline factors and prior ther-  
 472 apy for nivolumab versus everolimus in advanced renal cell  
 473 carcinoma. *European Urology*. 2017;72(6):962-71.  
 474

- 475 [29] Meeting Library | Pembrolizumab monotherapy as first- 496  
476 line therapy in advanced clear cell renal cell carcinoma 497  
477 (accRCC): Results from cohort A of KEYNOTE-427. 498  
478 [Internet]. [cited 2018 Jun 4]. Available from: <https://meetinglibrary.asco.org/record/160560/abstract>. 499
- 480 [30] Hodi FS, Chiarion-Sileni V, Gonzalez R, Grob J-J, 500  
481 Rutkowski P, Cowey CL, et al. Nivolumab plus ipilimumab 501  
482 or nivolumab alone versus ipilimumab alone in advanced 502  
483 melanoma (CheckMate 067): 4-year outcomes of a multi- 503  
484 centre, randomised, phase 3 trial. *The Lancet Oncology*. 504  
485 2018;19(11):1480-92. 505
- 486 [31] Robert C, Long GV, Schachter J, Arance A, Grob JJ, 506  
487 Mortier L, et al. Long-term outcomes in patients (pts) with 507  
488 ipilimumab (ipi)-naive advanced melanoma in the phase 508  
489 3 KEYNOTE-006 study who completed pembrolizumab 509  
490 (pembro) treatment. *Journal of Clinical Oncology*. 2017;35 510  
491 (15\_suppl):9504-4. 511
- 492 [32] Wolchok JD, Chiarion-Sileni V, Gonzalez R, Rutkowski P, 512  
493 Grob J-J, Cowey CL, et al. Overall survival with combined 513  
494 nivolumab and ipilimumab in advanced melanoma. *New 514*  
495 England Journal of Medicine. 2017;377(14):1345-56.
- [33] E Cortesi, F Cappuzzo, L. Galli, A. Bearz, S. Pignata. 892P 496  
- Treatment beyond progression in patients with advanced 497  
RCC participating in the Expanded Access Programme 498  
(EAP)E. Cortesi1, F. Cappuzzo2, L. Galli3, A. Bearz4, S. 499  
Annals of Oncology (2017) 28 (suppl.5): v295-v329. 2017. 500
- [34] Escudier B, Motzer RJ, Sharma P, Wagstaff J, Plimack 501  
ER, Hammers HJ, et al. Treatment beyond progression 502  
in patients with advanced renal cell carcinoma treated 503  
with nivolumab in checkMate 025. *European Urology*. 504  
2017;72(3):368-76. 505
- [35] Hyperprogressive disease (HPD) is a new pattern of progres- 506  
sion in cancer patients treated by anti-PD-1/PD-L1 | Clinical 507  
Cancer Research [Internet]. [cited 2019 Jan 15]. Available 508  
from: <http://clincancerres.aacrjournals.org/content/early/2016/11/08/1078-0432.CCR-16-1741>. 509
- [36] De Velasco G, Krajewski KM, Albiges L, Awad MM, Bell- 511  
munt J, Hodi FS, et al. The heterogeneity of radiologic 512  
response with anti-PD-1/PD-L1 therapy in metastatic Renal 513  
Cell Carcinoma (mRCC). *Cancer Immunol Res*. 2015. 514