

The difference in the dietary inflammatory index, functional food, and antioxidants intake between COVID -19 patients and healthy persons

Mahsa Mohajeri^{a,b}, Reza Mohajery^{b,c}, Ali Nemati^{a,d,*} and Farhad Pourfarzi^{a,*}

^a*Digestive Disease Research Center, Ardabil University of Medical Sciences, Ardabil, Iran*

^b*Academic Center for Education, Culture and Research, Ardabil, Iran*

^c*Energy Management Research Center, University of Mohaghegh Ardabili, Ardabil, Iran*

^d*Department of Clinical Biochemistry, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran*

Received 10 July 2021

Accepted 20 January 2022

Pre-press 29 January 2022

Abstract.

BACKGROUND: The healthy diet is important to maintain immunity against infection. This study aimed to assess and compare the consumption of functional foods, some antioxidants, and dietary inflammatory index between Iranian COVID-19 patients and healthy persons.

METHODS: This case-control study was conducted between 1000 (500 cases and 500 controls) adults aged 18–65 years in Iran, that were sampling based on the snowball method and their information was collected electronically. The dietary intake was assessed using the Food Frequency Questionnaire (FFQ).

RESULTS: There was a significant difference ($p=0.044$) in vitamin D consumption between healthy people and COVID-19 patients. Vitamin E intake in healthy participants was significantly ($p=0.041$) more than COVID-19 patients. There was a significant difference in Zinc ($p=0.011$), selenium ($p=0.021$), and vitamin C ($p=0.023$) between healthy persons and COVID-19 patients. Healthy participants' consumption of onion (56.5 ± 7.82 g/day), garlic (4.32 ± 0.01 g/day) and oat (6.32 ± 0.71 g/day) was significantly ($p \leq 0.05$) more than COVID-19 patients. With the increase of each unit in the score of the dietary inflammatory index, the risk of COVID-19 incidence increased 1.63 times (OR = 1.63 95%CI: 1.54–1.72). There was an inverse association between the consumption of antioxidants and functional foods with the risk of COVID-19 incidence in the study population ($p \leq 0.05$).

*Corresponding author: Ali Nemati, Digestive Disease Research Center, Ardabil University of Medical Sciences, Ardabil, Iran. E-mail: nutrition1391@gmail.com. and Farhad Pourfarzi, Department of Clinical Biochemistry, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran. E-mail: farhad.pourfarzi@gmail.com.

CONCLUSION: Healthy people consumption of antioxidants and functional foods was more than COVID-19 patients and there was a significant inverse association between the risk of COVID- 19 incidence with the consumption of functional foods and antioxidants. Increasing the dietary inflammatory index score increased the risk of COVID- 19 incidence. There is a need for further clinical trials to confirm the effect of consuming functional foods and antioxidants on the prevention or treatment of COVID-19.

Keywords: COVID-19, antioxidants, functional foods, dietary inflammatory index

1. Introduction

The results of studies have suggested that increased inflammation has an important role in the COVID-19 incidence [1, 2].

Dietary nutrients can affect the immune system through stimulation cells, a manifestation of the gene, and modification of signaling molecules [3, 4]. Many dietary components shape the responses of the body immune like as functional foods and dietary antioxidants that are important factors for the keep the function of the immunity system. These foods expand life quality by prevention of nutrition-related diseases [5–8]. Recent studies [9–11] have suggested that functional foods consumption can improve COVID-19 symptoms, but until now no study has examined these foods consumption impacts on COVID 19 severity of symptoms.

In addition to functional foods, other dietary components can reduce body inflammation levels, like antioxidants. Based on studies results, the antioxidants reduce oxidative stress and serum inflammatory markers. Since COVID -19 disease is also an inflammatory disease [12], the effect of these nutrients on the serum inflammation marker in these patients can be discussed. To date, no study has assessed the intake of functional foods, antioxidants, and dietary inflammation index in COVID -19 patients. Therefore, this study assessed and compared the consumption of functional foods, some antioxidants, and dietary inflammation index between Iranian COVID -19 patients and healthy persons.

2. Methods

This case-control (population base study) study was done between 1000 (500 cases and 500 controls) adults aged 18–65 years old that chosen based on the snowball method and their information was collected electronically. Electronic invitations were sent to people on WhatsApp and they were asked to send this invitation form to their friends and family members. At the beginning of the study, 2540 people participated in this survey through an electronic questionnaire (includes demographic characteristics, history of hypertension, diabetes, hyperlipidemia, COVID- 19 incidence history, and food frequency questionnaire). From all participants, 500 participants had a history of COVID -19 who were diagnosed using a valid laboratory test and have been treated out of the hospital. Then, among the healthy individuals who participated in the initial study 500 healthy individuals were selected by simple sampling. The matching was performed for the control group based on age and sex. All individuals completed a written electronic consent form to participate in the study. This study was approved by the ethics committee of Ardabil University of Medical Sciences and its ethics code is as follows IR.ARUMS.REC.1400.008. This project was financially supported by NIMAD project No.962249.

2.1. Dietary assessment

The nutritional status of study participants was assessed by a food frequency questionnaire (FFQ) with 138 items. In this questionnaire, there is a standard size for each food item, which is designed according to Willett

66 method. The study participants were asked to determine the frequency of their consumption of each food item
67 according to the amount consumed in the previous year. Depending on the type of food consumed, repeated
68 consumption per day, week, or month is asked. Past studies that have used this questionnaire have provided
69 acceptable values of results and a review of the validity of this questionnaire shows that the questionnaire has
70 acceptable validity. Based on the portion size of each food, the amounts consumed from each food item were
71 converted to grams per day, then this information was entered into the Nutritionist IV software and the amounts
72 of all nutrients (vitamin D, vitamin E, vitamin C, vitamin A, Selenium, Zinc) consumed were calculated with
73 this software [13].

74 2.2. *The Dietary Inflammatory Index (DII):*

75 Briefly, the DII is a scoring algorithm based on an extensive review of the literature published from 1950
76 to 2010, linking 1943 articles to a total of forty-five food parameters and including various macronutrients,
77 micronutrients, flavonoids, and food items. These dietary parameters were scored according to whether they
78 increased [1], decreased, or had no effect (0) on six inflammatory biomarkers. An overall food parameter-
79 specific inflammatory effect score was calculated and multiplied by a centered percentile value for each food.
80 This percentile was calculated by first linking the dietary data from a study to the regionally representative world
81 database intake, which was based on actual human consumption in eleven populations from different parts of the
82 world that provided a robust estimate of a mean and standard deviation for each parameter. These values then
83 become the multipliers to express an individual's exposure, relative to the 'standard global mean' as a z-score.
84 This was achieved by subtracting the 'standard global mean' from the amount reported and dividing this value
85 by the standard deviation. To minimize the effect of 'right skewing', this value was then converted to a centered
86 percentile score. The centered percentile score for each food parameter for each individual was multiplied by
87 the respective food parameter effect score that was derived from the literature review to obtain a food parameter-
88 specific DII score for an individual. All of the food parameter-specific DII scores were then summed to create
89 the overall DII score for every participant in the study. The greater the DII score, the more pro-inflammatory
90 the diet, and more negative values represent more anti-inflammatory diets. The DII score could take on values
91 ranging from 7.98 (maximally pro-inflammatory) to 28.87 (maximally anti-inflammatory). Construct validation
92 of the DII was performed using data derived from two different sources of dietary intake information, and serum
93 high-sensitivity CRP as the construct validator [14].

94 2.3. *Antioxidants consumption*

95 The food intake data from FFQ were converted to antioxidants intake (vitamin A, vitamin C, vitamin E,
96 vitamin D, zinc, selenium) using the Nutritionist IV software: Based on each food portion size, the amount of
97 food consumed was converted to gram and then these amounts were entered into the Nutritionist IV software
98 and by this software, the amounts of all micronutrients including vitamins and minerals were calculated.

99 2.4. *Functional foods intake*

100 The dietary information recorded in the FFQ was used to calculate the intake of some functional foods (oat,
101 garlic, onion, banana, turmeric, green tea). Based on each food portion size, its consumption was changed to
102 grams per day.

103 2.5. *Statistical methods*

104 For data analysis, the SPSS software version 18 was used. The normality of the variables was assessed by
105 the Kolmogorov-Smirnov test. Independent *T*-test, Chi-square test, and one-way ANOVA were used to compare

Table 1
Demographical characteristics of the participants according to the categories of the DII score by COVID-19

Variables	Healthy people				COVID-19 PATIENTS			
	Q1 N = 120	Q2 N = 298	Q3* N = 82	P	Q1 N = 20	Q2 N = 96	Q3* N = 384	P
DII mean \pm SD	18.9 \pm 5.2	19.6 3.8	20.6 4.9	0.041**	19.9 \pm 4.5	20.9 \pm 5.9	23.8 \pm 9.7	0.021**
Age (years) mean \pm SD	38.2 \pm 4.2	43.1 \pm 3.9	45.2 \pm 6.1	0.02**	38.4 \pm 4.2	42.8 \pm 5.8	32.8 \pm 9.7	0.042**
Family history of early CHD (%)	100(83.33%)	160(53.69%)	80(97.56%)	0.022#	5(25%)	42(43.75%)	201(52.34%)	0.014#
Hypertension n (%)	35(7%)	160(53.69%)	68(85%)	0.050#	10(50%)	38(39.58%)	220(57.29%)	0.031#
Dyslipidaemia n(%)	65(54.16%)	115(35.59%)	65(79.26%)	0.012#	5(25%)	36(37.5%)	195(50.78%)	0.025#
Diabetes n(%)	55(45.83%)	108(36.24%)	54(65.85%)	0.014#	4(20%)	61(63.54%)	119(30.98%)	0.027#
Marital status n(%)				0.004#				0.043#
Married	72(60%)	66	60		92	91	89	
Widowed	40(33.33%)	27	27		3	3	3	
Single/other	8(6.66%)	7	7		5	6	8	
Educational level n(%)				0.001#				0.061#
Primary education or less	53	86	87		22	31	29	
Secondary education	25	11	9		38	39	39	
Any college	22	3	4		40	30	32	

*Highest anti-inflammatory values of the DII. **: Based on one way ANOVA test, #: based on Chi-square test.

parametric and nonparametric/nominal variables between groups. The association between the COVID-19 incidence with consumption of functional foods and antioxidants intake was assessed by the multivariable logistic regression test. In all analyzes, a p -value less than 0.05 was considered statistically significant.

3. Results

This case-control study was conducted between 1000 (500 COVID-19 patients and 500 healthy adults) Iranian adults. There was no significant difference in the participants' age between groups of study ($p = 0.07$). The participants mean \pm SD of age was 42.1 \pm 8.2 years for COVID-19 patients and 42.3 \pm 7.9 years for the control group. The mean DII score for the control group was 20.4 \pm 3.5 and 22.9 \pm 4.1 for the COVID-19 patients ($p = 0.042$). Table-1 reported the participant's demographical characters according to the DII score categories by COVID-19. In both groups, the history of diabetes, cardiovascular disease, and hypertension were significantly ($p \leq 0.05$) higher in those with more DII scores.

Table-2 shows the results of comparing the consumption of antioxidants and functional foods in the two groups of study. There was a significant difference ($p = 0.044$) in vitamin D consumption between healthy people (3.14 \pm 0.14 Mg/day) and COVID-19 patients (2.02 \pm 0.22 Mg/day). Vitamin E intake of healthy participants (15.25 \pm 1.18 mg/day) was significantly ($p = 0.041$) more than COVID-19 patients (13.37 \pm 1.32 mg/day). There was a significant difference in Zinc ($p = 0.011$), selenium ($p = 0.021$), and vitamin C ($p = 0.023$) intake between healthy persons and COVID-19 patients. Healthy participants' consumption of onion (56.5 \pm 7.82 g/day), garlic (4.32 \pm 0.01 g/day) and oat (6.32 \pm 0.71 g/day) was significant ($p \leq 0.05$) more than COVID-19 patients.

Table 3 reported the association of COVID-19 incidence with dietary inflammatory index, antioxidants, and functional foods intake. The logistic regression indicated that with each gram increase in oat consumption, the risk of COVID-19 incidence in the study participants decreased by 13% (OR = 0.13 95% CI: 1.33–1.55). Each gram increase in onion consumption reduced the risk of COVID-19 incidence by 97% (OR = 0.97 95% CI:

Table 2
Dietary consumption in the COVID-19 patients and healthy people

Nutritional intake	COVID-19 Patients Mean \pm SD	Healthy persons Mean \pm SD	Recommended Dietary Allowance	P*
Antioxidants				
Vitamin C(mg/day)	63.53 \pm 2.35	78.3 \pm 1.44	75–90	0.023
Vitamin A) Mg/day)	702.4 \pm 15.44	702.3 \pm 13.12	900–700	0.063
Vitamin E(mg/day)	13.37 \pm 1.32	15.25 \pm 1.18	15	0.041
Vitamin D (Mg/day)	2.02 \pm 0.22	3.14 \pm 0.14	10	0.044
Zinc (mg/day)	7.23 \pm 0.45	8.1 \pm 0.28	11–8	0.011
Selenium (Mg/day)	42.21 \pm 2.21	51.02 \pm 3.1	55	0.021
Functional foods (g/day)				
Green Tea (cup/day)	0.51 \pm 0.002	0.52 \pm 0.005	–	0.054
Oat	6.32 \pm 0.71	7.44 \pm 0.68	–	0.024
Garlic	4.32 \pm 0.01	7.91 \pm 0.21	–	0.017
Onion	56.5 \pm 7.82	102.44 \pm 10.18	–	0.038
Turmeric	2.51 \pm 0.001	2.58 \pm 0.007	–	0.712
Banana	25.33 \pm 4.12	25.31 \pm 5.17	–	0.318

*: based on independent *T*-test.

Table 3
The COVID 19-incidence association as dependent variable with dietary inflammatory index, functional foods and antioxidants consumption

Independent variables	OR (95%CI)	P*
Dietary inflammatory index	1.63 (1.54–1.72)	\leq 0.0001
Green tea (cup)	0.001 (0.37–1.17)	0.99
Oat (g/d)	0.13 (1.33–1.55)	\leq 0.0001
Banana (g/d)	0.98 (0.43–1.11)	0.22
Turmeric (g/d)	0.86 (0.51–1.04)	0.194
Onion (g/d)	0.97 (1.14–1.38)	0.003
Vitamin D (mg/d)	0.87 (1.14–1.61)	\leq 0.0001
Vitamin A (mg/d)	0.89(0.97–1.24)	0.065
Vitamin C (mg/d)	0.95(1.10–1.34)	0.04
Selenium (mg/d)	0.91(1.95–2.21)	0.003
Zinc (mg/d)	0.92(1.34–1.82)	0.002

*: Based on multivariable logistic regression.

128 1.14–1.38). An increase of one unite in Zinc (OR = 0.92 95% CI : 1.34–1.82), Selenium (OR = 0.91 95 CI%:
129 1.95–2.21), vitamin C (OR = 0.95 95% CI: 1.10–1.34), vitamin E (OR = 0.86 95 CI% : 1.04–1.21) and vitamin
130 D (OR = 0.87 95% CI: 1.14–1.61) consumption also reduced the risk of COVID-19 incidence by 92%, 91%,
131 and 87%, respectively. Increasing each score in the dietary inflammatory index, increased the risk of Covid-19
132 incidence by 1.63 (OR = 1.63 95%CI: 1.54–1.72).

4. Discussion

This was the first study that assessed the DII, some functional foods, and antioxidant consumption in Iranian COVID-19 patients compared to healthy people. The results showed that the consumption of antioxidants and functional foods in COVID-19 patients was less than healthy individuals. Also, the score of dietary inflammation index in COVID-19 patients was more than healthy individuals. Also, the dietary inflammation index in COVID-19 patients was more than healthy individuals. The association between vitamin D and the incidence and even severity of COVID-19 has been well established in previous studies [15, 16].

Studies have suggested that a diet with more antioxidants can reduce the serum levels of inflammatory markers in patients and also increase patients' resistance to this viral infection by reducing oxidative stress in the body [17, 18]. Mrityunjaya et al in one study have expressed that a combination of some antioxidants including Zinc, vitamin D, vitamin C, curcumin, cinnamaldehyde, probiotics, selenium, lactoferrin, quercetin, etc, in the form of a food supplement can boost the immune system, prevent virus spread, and decrease the serum inflammation markers [9, 19, 20]. Jothimani et al. [21] in one study indicated that a significant number of COVID-19 patients were zinc deficient. These zinc-deficient patients developed more complications, and the deficiency was associated with a prolonged hospital stay and increased mortality.

In the results of this study, the green tea consumption of healthy people was more than COVID-19 patients, but this difference was not significant. Because green tea is rich in many antioxidants, it can protect the body against oxidative stress caused by the COVID-19 virus [22, 23]. Based on Szulińska et al. study, green tea can reduce the TNF- α , so suppress the body's inflammation [24]. Although no study has examined the effect of green tea consumption in improving or preventing COVID -19, according to the results of previous studies [25–27], it can be predicted that consuming green tea can increase the body's total antioxidant capacity and can prevent the increased serum pro-inflammation side effects in COVID-19 patients.

The onion, garlic, and oat consumption of healthy participants was more than COVID-19 patients. Garlic, oat, and onion are functional foods and according to many studies results, they have many benefits for the body. Pre-clinical data reported that garlic has potential antiviral activity against pathogenic viruses [18, 28, 29]. Functional foods contain phytochemicals that have anti-cancer, anti-bacterial, and anti-inflammatory effects, and their positive effects have been proven in all studies [11, 30, 31]. These foods are "items that modify the battered functions in the body" and increase the immune response and decrease the risk of many diseases. The functional foods and components have immune-boosting properties [32]. These natural compounds do not participate in the direct embarrassment response against COVID-19, but the complete response of immunity in COVID-19 prevention [33]. Micronutrients like vitamin C, D, probiotic foods, flavonoids, carotenoids, and herbs have been reported as functional foods [34].

The results of the present study showed that there is a significant relationship between the intake of vitamin C, vitamin D, vitamin E, and some functional foods such as garlic, onion, and oat with the risk of COVID-19 incidence. By increasing each unit in the amount of consumption of these foods and nutrients, the risks of COVID-19 incidence in the study participants were reduced. However, the risks of COVID-19 incidence in the study population increased with an increasing score of dietary inflammation index. Consistent with the present study result, Moloudi et al. in a study showed that patients with the maximum pro-inflammatory energy-adjusted E-DII score had 7.26 times greater odds of developing COVID-19, as compared to those in tertiles 1 (E-DII T3 v. E-DII T1: OR = 7.26; 95 % CI 2.64 to 9.94, $P < 0.001$). Also, a positive association between E-DII and C-reactive protein (CRP) was observed (BE-DII = 1.37, 95 % CI 0.72, 2.02), such that with each unit increase in E-E-DII, the CRP levels were increased by 1.37 units. Furthermore, a significant association was found between E-DII and the severity of disease (BE-DII = 0.03, 95 % CI 0.01, 0.06, 0.024).

Considering that the increase in inflammatory serum indices in COVID -19 patients is one of the reasons for severe respiratory symptoms, etc.[35], it is necessary to reduce these inflammatory indices in patients using a healthy diet. Studies have shown that an inflammatory diet increases the risk of chronic diseases. Several studies have indicated a converse association between healthy diets and serum indicators of inflammation, as well as a

180 direct association with 'unhealthy diet [8–14]. Exactly, more consumption of fruits and vegetables decreases the
181 CRP concentration [36]. On the contrary, more red meat consumption increases the serum inflammatory markers
182 [37, 38].

183 In this study results,

184 Because the diet's role in the control and prevention of COVID-19 is important, improving the dietary patterns
185 of people can prevent serious complications of this disease. This study is the first study in Iran that examined
186 the inflammation index of dietary, some antioxidants consumption, and functional food intake in people with a
187 history of COVID 19 incidence. One of the limitations of this study is the use of an electronic questionnaire. In
188 this questionnaire, the accuracy of information is less than a face-to-face interview. We suggest that next future
189 studies complete the information using face-to-face interviews.

190 5. Conclusion

191 This study showed that the dietary inflammatory index in COVID -19 patients was 19 more than healthy
192 individuals. The antioxidants and functional foods consumption in healthy people was more than COVID-19
193 patients and there was a significant inverse association between the risk of COVID- 19 incidence with the
194 consumption of functional foods and antioxidants. Increasing the dietary inflammatory index increased the risk
195 of COVID- 19 incidence. There is a need for further clinical trials to confirm the effect of consuming functional
196 foods and antioxidants on the prevention or treatment of COVID-1.

197 Acknowledgment

198 The authors of this article thank Ardabil University of Medical Sciences for their financial and spiritual support.

199 Conflict of interest

200 The authors declare that there is no conflict of interest.

201 References

- 202 [1] Zeng F, Huang Y, Guo Y, Yin M, Chen X, Xiao L, et al. Association of inflammatory markers with the severity of COVID-19: a
203 meta-analysis. *International Journal of Infectious Diseases*. 2020;96:467-74.
- 204 [2] Liu B, Li M, Zhou Z, Guan X, Xiang Y. Can we use interleukin-6 (IL-6) blockade for coronavirus disease 2019 (COVID-19)-induced
205 cytokine release syndrome (CRS)? *Journal of Autoimmunity*. 2020:102452.
- 206 [3] Calder PC, Carr AC, Gombart AF, Eggersdorfer M. Optimal nutritional status for a well-functioning immune system is an important
207 factor to protect against viral infections. *Nutrients*. 2020;12(4):1181.
- 208 [4] Naja F, Hamadeh R. Nutrition amid the COVID-19 pandemic: a multi-level framework for action. *European Journal of Clinical
209 Nutrition*. 2020;74(8):1117-21.
- 210 [5] Farag MA, Abdelwareth A, Sallam IE, El Shorbagi M, Jehmlich N, Fritz-Wallace K, et al. Metabolomics reveals impact of seven
211 functional foods on metabolic pathways in a gut microbiota model. *Journal of Advanced Research*. 2020;23:47-59.
- 212 [6] Mohajeri M, Horriatkhah E, Mohajery R. The effect of glutamine supplementation on serum levels of some inflammatory factors,
213 oxidative stress, and appetite in COVID-19 patients: a case-control study. *Inflammopharmacology*. 2021;29(6):1769-76.
- 214 [7] Mohajeri M, Houjehani S, Ghahremanzadeh M, Borghei MH, Moradi F, Barzegar A. Some behavioral risk factors of obesity in
215 Ardabil-Iran adults. *Obesity Medicine*. 2020;18:100167.
- 216 [8] Mohajeri M, Ghannadiazl F, Narimani S, Nemat A. The food choice determinants and adherence to Mediterranean diet in Iranian
217 adults before and during COVID-19 lockdown: population-based study. *Nutrition & Food Science*. 2021.

- 218 [9] Iddir M, Brito A, Dingo G, Fernandez Del Campo SS, Samouda H, La Frano MR, et al. Strengthening the immune system and reducing
219 inflammation and oxidative stress through diet and nutrition: considerations during the COVID-19 crisis. *Nutrients*. 2020;12(6):1562.
- 220 [10] Yang F, Zhang Y, Tariq A, Jiang X, Ahmed Z, Zhihao Z, et al. Food as medicine: A possible preventive measure against coronavirus
221 disease (COVID-19). *Phytotherapy Research*. 2020;34(12):3124-36.
- 222 [11] Díaz LD, Fernández-Ruiz V, Cámara M. An international regulatory review of food health-related claims in functional food products
223 labeling. *Journal of Functional Foods*. 2020;68:103896.
- 224 [12] Tay MZ, Poh CM, Rénia L, MacAry PA, Ng LF. The trinity of COVID-19: immunity, inflammation and intervention. *Nature Reviews*
225 *Immunology*. 2020;20(6):363-74.
- 226 [13] Mohammadi N, Sajjadi F, Maghroun M, Alikhasi H, Nilforoush Zadeh F, Sarrafzadegan N. Validation of a simplified food frequency
227 questionnaire for the assessment of dietary habits in Iranian adults: Isfahan Healthy Heart Program, Iran. *ARYA Atherosclerosis*.
228 2015;11(2):139.
- 229 [14] Shivappa N, Hébert JR, Rietzschel ER, De Buyzere ML, Langlois M, Debruyne E, et al. Associations between dietary inflammatory
230 index and inflammatory markers in the Asklepios Study. *British Journal of Nutrition*. 2015;113(4):665-71.
- 231 [15] Ebadi M, Montano-Loza AJ. Perspective: improving vitamin D status in the management of COVID-19. *European Journal of Clinical*
232 *Nutrition*. 2020;74(6):856-9.
- 233 [16] Vimala Swaran KS, Forouhi NG, Khunti K. Vitamin D and covid-19. *British Medical Journal Publishing Group*; 2021.
- 234 [17] DE FLORA S, Balansky R, LA MAESTRA S. Antioxidants and COVID-19. *Journal of Preventive Medicine and Hygiene*. 2021;62(1
235 Suppl 3):E34.
- 236 [18] Darenkaya M, Kolesnikova L, Kolesnikov S. The Association of Respiratory Viruses with Oxidative Stress and Antioxidants.
237 Implications for the COVID-19 Pandemic. *Current Pharmaceutical Design*. 2021;27(13):1618-27.
- 238 [19] Beltrán-García J, Osca-Verdegal R, Pallardó FV, Ferreres J, Rodríguez M, Mulet S, et al. Oxidative stress and inflammation in
239 COVID-19-associated sepsis: the potential role of anti-oxidant therapy in avoiding disease progression. *Antioxidants*. 2020;9(10):936.
- 240 [20] Zabetakis I, Lordan R, Norton C, Tsoupras A. COVID-19: the inflammation link and the role of nutrition in potential mitigation.
241 *Nutrients*. 2020;12(5):1466.
- 242 [21] Jothimani D, Kailasam E, Danielraj S, Nallathambi B, Ramachandran H, Sekar P, et al. COVID-19: Poor outcomes in patients with
243 zinc deficiency. *International Journal of Infectious Diseases*. 2020;100:343-9.
- 244 [22] Masek A, Chrzescijanska E, Latos M, Zaborski M, Podsedek A. Antioxidant and antiradical properties of green tea extract compounds.
245 *Int J Electrochem Sci*. 2017;12:6600-10.
- 246 [23] Li Y, Rahman SU, Huang Y, Zhang Y, Ming P, Zhu L, et al. Green tea polyphenols decrease weight gain, ameliorate alteration
247 of gut microbiota, and mitigate intestinal inflammation in canines with high-fat-diet-induced obesity. *The Journal of Nutritional*
248 *Biochemistry*. 2020;78:108324.
- 249 [24] Szulińska M, Stepień M, Kregielska-Narozna M, Suliburska J, Skrypnik D, Bak-Sosnowska M, et al. Effects of green tea supple-
250 mentation on inflammation markers, antioxidant status and blood pressure in NaCl-induced hypertensive rat model. *Food & Nutrition*
251 *Research*. 2017;61(1):1295525.
- 252 [25] Selvan DA, Mahendiran D, Kumar RS, Rahiman AK. Garlic, green tea and turmeric extracts-mediated green synthesis of silver
253 nanoparticles: Phytochemical, antioxidant and *in vitro* cytotoxicity studies. *Journal of Photochemistry and Photobiology B: Biology*.
254 2018;180:243-52.
- 255 [26] Jakubczyk K, Kochman J, Kwiatkowska A, Kałduńska J, Dec K, Kawczuga D, et al. Antioxidant properties and nutritional composition
256 of matcha green tea. *Foods*. 2020;9(4):483.
- 257 [27] Wang J, Yang Z, Celi P, Yan L, Ding X, Bai S, et al. Alteration of the antioxidant capacity and gut microbiota under high levels of
258 molybdenum and green tea polyphenols in laying hens. *Antioxidants*. 2019;8(10):503.
- 259 [28] Rouf R, Uddin SJ, Sarker DK, Islam MT, Ali ES, Shilpi JA, et al. Antiviral potential of garlic (*Allium sativum*) and its organosulfur
260 compounds: A systematic update of pre-clinical and clinical data. *Trends Food Sci Technol*. 2020;104:219-34.
- 261 [29] Shehzad A, Zahid A, Mahmood S, Suleria HAR. *Functional Foods: Concepts and Their Health Perspectives*. Human Health Benefits
262 of Plant Bioactive Compounds: Apple Academic Press; 2019. pp. 3-20.
- 263 [30] Maqsood S, Adiamo O, Ahmad M, Mudgil P. Bioactive compounds from date fruit and seed as potential nutraceutical and functional
264 food ingredients. *Food Chemistry*. 2020;308:125522.
- 265 [31] Rad AH, Maleki LA, Kafil HS, Zavoshti HF, Abbasi A. Postbiotics as novel health-promoting ingredients in functional foods. *Health*
266 *Promotion Perspectives*. 2020;10(1):3-4.
- 267 [32] BAYIR AG, AKSOY AN, KOÇYİĞİT A. The importance of polyphenols as functional food in health. *Bezmialem Science*.
268 2019;7(2):157.
- 269 [33] Temesi Á, Bacsó Á, Grunert KG, Lakner Z. Perceived correspondence of health effects as a new determinant influencing purchase
270 intention for functional food. *Nutrients*. 2019;11(4):740.

- 271 [34] Lencsényi KG, Bast A, de Boer A. Clarifying the health claim assessment procedure of EFSA will benefit functional food innovation.
272 *Journal of Functional Foods*. 2018;47:386-96.
- 273 [35] Zeng Z, Yu H, Chen H, Qi W, Chen L, Chen G, et al. Longitudinal changes of inflammatory parameters and their correlation with
274 disease severity and outcomes in patients with COVID-19 from Wuhan, China. *Critical Care*. 2020;24(1):1-12.
- 275 [36] Taborelli M, Polesel J, Parpinel M, Stocco C, Birri S, Serraino D, et al. Fruit and vegetables consumption is directly associated to
276 survival after prostate cancer. *Molecular Nutrition & Food Research*. 2017;61(4):1600816.
- 277 [37] Das DC, Jahan I, Uddin MG, Hossain MM, Chowdhury MAZ, Fardous Z, et al. Serum CRP, MDA, vitamin C, and trace elements in
278 Bangladeshi patients with rheumatoid arthritis. *Biological Trace Element Research*. 2021;199(1):76-84.
- 279 [38] Bahrami A, Bahrami-Taghanaki H, Khorasanchi Z, Tayefi M, Ferns GA, Sadeghnia HR, et al. the association between neuropsychological
280 function with serum vitamins A, D, and E and hs-CRP concentrations. *Journal of Molecular Neuroscience*. 2019;68(2):243-50.

Uncorrected Author Proof