

Effectiveness of spiritual care training for rehabilitation professionals: An exploratory controlled trial

Kate Fiona Jones^{a,g,*}, Julie Pryor^{a,b}, Candice Care-Unger^a, Joseph Descallar^{e,f} and Grahame Kenneth Simpson^{c,d,e}

^aRoyal Rehab, PO Box 6, Ryde, Sydney, NSW, Australia

^bSusan Wakil School of Nursing & Midwifery, University of Sydney, Sydney, NSW, Australia

^cBrain Injury Rehabilitation Research Group, Ingham Institute for Applied Medical Research, Sydney, NSW, Australia

^dSchool of Human Services and Social Work, Griffith University, Brisbane, QLD, Australia

^eIngham Institute for Applied Medical Research, Liverpool, NSW, Australia

^fSouth Western Sydney Clinical School, University of New South Wales, Liverpool, Australia

^gInstitute for Ethics and Society, University of Notre Dame, Sydney, Australia

Abstract.

BACKGROUND: Spirituality has been recognised to play an important role in neurorehabilitation, however research findings indicate that rehabilitation professionals do not feel well equipped to deliver spiritual care.

OBJECTIVE: To evaluate a spiritual care training program for rehabilitation professionals.

METHODS: An exploratory controlled trial was conducted. Participants enrolled in a two-module spiritual care training program. Spiritual care competency was measured with the Spiritual Care Competency Scale. Confidence and comfort levels were measured using the domains of the Spiritual Care Competency Scale. The Spirituality and Spiritual Care Rating Scale assessed participant attitudes and knowledge. Measures were administered three times: pre-program, post-program and six weeks after program completion.

RESULTS: The training was attended by 41 rehabilitation professionals working in spinal cord or traumatic brain injury. Thirty-two control group participants were recruited. Multilevel models found that for levels of spiritual care competency, confidence, comfort, and ratings on existential spirituality, pre intervention scores increased significantly in the intervention group at post intervention ($p < 0.05$) and were maintained at follow-up.

CONCLUSIONS: The results of this study demonstrated that a spiritual care training program was effective in increasing levels of self-reported competency, confidence and comfort in delivery of spiritual care for rehabilitation professionals.

Keywords: Spirituality, spiritual care, health professionals, spinal cord injury, traumatic brain injury, rehabilitation

1. Introduction

Traumatic brain injury and spinal cord injury are life changing injuries which can impact upon a person's physical, psychological, emotional or spiritual well-being. While much research has focused upon the negative impacts of neurotrauma, a growing

*Address for correspondence: Kate Jones, Institute for Ethics and Society, University of Notre Dame, PO Box 944, Broadway NSW 2007. Australia. Tel.: +61 414 875 163; E-mail: kate.jones1@nd.edu.au.

body of literature is emphasising the strengths and resilience of injured people and their family members (White, Driver, & Warren, 2008). One factor increasingly thought to contribute to resilience is spirituality (Fricchione & Nejad, 2012; Smith, Ortiz, Wiggins, Bernard, & Dalen, 2012; Walsh, 2003).

Spirituality has been described as ‘the aspect of humanity that refers to the way individuals seek and express meaning and purpose, and the way they experience their connectedness to the moment, to self, to others, to nature and to the significant or sacred’ (Puchalski et al., 2009). Spirituality and religion have generally been described as distinct but overlapping constructs, with religion considered to encapsulate “an institutionalised (i.e. systematic) pattern of values, beliefs, symbols, behaviours, and experiences that are oriented toward spiritual concerns, shared by a community, and transmitted over time in traditions” (Canda & Furman, 2009, p. 59). This positions spirituality as the broader of the two constructs, encompassing a range of different sources of meaning and connection, including but not limited to religious faith (Davis et al., 2015; Jones, Dorsett, Simpson, & Briggs, 2018).

The role of spirituality in promoting whole-of-person care is becoming evident within healthcare (Cobb, Puchalski, & Rumbold, 2012; Koenig, 2012). The results of two recent scoping reviews demonstrate that spirituality has been positively associated with quality of life, life satisfaction, mental and physical health, and resilience after both spinal cord injury (SCI) (Jones, Simpson, Briggs, & Dorsett, 2016) and traumatic brain injury (TBI) (Jones, Pryor, Care-Unger, & Simpson, 2018). Johnstone, Glass and Oliver (2007) have argued that addressing the spiritual needs of people affected by chronic disabilities, such as TBI and SCI, may be equally as important as addressing those needs in people with end-of-life conditions or illnesses. They suggested spirituality (or religion) may help such people “cope with their disability, give new meaning to their lives based on their newly acquired disabilities, and help them to establish new life goals” (p.1155).

Spiritual care has been described as “person centred care which seeks to help people (re)discover hope, resilience and inner strength in times of illness, injury, transition and loss” (NHS Education for Scotland, 2013). Despite the increasing awareness of the importance of spirituality after neurotrauma, existing research suggests spirituality is not well incorporated in neurorehabilitation practice (Jones, Dorsett, Briggs, & Simpson, 2018; Jones, Pryor, Care-Unger,

& Simpson, 2020). A recent study revealed that while rehabilitation health professionals acknowledged the importance of spirituality for patients, several barriers to addressing patients’ spiritual needs were identified; these included a need for more training (80%), not enough time (74%) and personal discomfort (61%) (Jones et al., 2020). These findings are consistent with results from other healthcare areas and disciplines, including palliative care doctors (Best, Butow, & Olver, 2016), acute care nurses (Gallison, Xu, Jurgens, & Boyle, 2012), social workers (Oxhandler, Parrish, Torres, & Achenbaum, 2015) and physiotherapists (Oakley, Katz, Sauer, Dent, & Millar, 2010). These studies have also demonstrated that overcoming these barriers in the delivery of spiritual care is important for healthcare professionals from a range of disciplines.

A number of spiritual care training programs and resources have been developed and trialled within healthcare settings to assist healthcare professionals to better address the spiritual needs of clients (NHS Education for Scotland, 2009). In a systematic review of the literature, Paal, Helo and Frick (2015) found that spiritual care training assisted participants to increase their awareness of personal spirituality and spiritual needs, clarify the role of spirituality and importance of spiritual care, and prepare trainees for spiritual encounters. However, Paal and colleagues also noted that few studies were well evaluated, and seldom involved a control group. Much of the training was conducted within the field of palliative care.

Professional development training in the contemporary health context has to compete with a broad range of other demands that health staff juggle in carrying out their daily duties. Within this context, multimodal presentation formats (online, face-to-face) employing brief training interventions are highly desirable. A few spiritual care programs have indicated that brief training in spiritual care can be effective in improving confidence and comfort levels in healthcare professionals. Cerra and Fitzpatrick (2008) observed that changes in healthcare professionals’ perceptions of spirituality were achieved after a two hour didactic lecture, while Meredith and colleagues (2012) reported changes in spiritual care and confidence after healthcare professionals attended a single workshop. Therefore, the aim of this study was to evaluate the effectiveness of a brief spiritual care training program to expand attitudes and knowledge regarding spirituality and spiritual care, and to increase rehabilitation professionals’ levels of competency, confidence and comfort in the

Table 1
Program outline

Module	Session Aims	Key Content	Format
1	To introduce the concept of spirituality, highlight its important role in rehabilitation, and present a range of different sources of spiritual strength that clients might draw upon	Spirituality and healthcare The importance of spirituality after traumatic injury What is spirituality? Sources of spiritual strength	<i>Self-study online</i> Written content Videod interviews of former clients
2	To build skills in spiritual care practice.	Understanding spirituality Introduction to spiritual care Introduction to spiritual care tools Role plays Looking after ourselves	<i>Workshop face-to-face</i> Didactic content Videod interviews Role plays Individual exercises

140 delivery of spiritual care. An underlying assumption
141 of the program was that spiritual care is relevant to all
142 healthcare disciplines, and therefore training should
143 be provided to all members of the multidisciplinary
144 team.

145 2. Methods

146 2.1. Participants

147 This study was an exploratory controlled trial. Ethical
148 approval was obtained from Northern Sydney
149 Local Health District Human Research Ethics Com-
150 mittee (LNR AU/1/5688313). Recruitment took place
151 between February and December 2019. The trial was
152 conducted across four specialised neurorehabilitation
153 units in Sydney Australia (2 TBI, 2 SCI). To limit
154 the possibility of contamination, two units (1 TBI, 1
155 SCI) were targeted for the training, with staff from the
156 other two units acting as controls. Invitations to partic-
157 ipate were distributed via email, or through direct
158 contact with study investigators, to all members of the
159 respective multidisciplinary teams. Written consent
160 was obtained from all participants, who participated
161 as volunteers.

162 2.2. Intervention

163 The Spiritual Care Training Program consisted of
164 two modules (see Table 1). Module 1 is a one fle-
165 hour computer-based self-study unit which includes
166 written information and video footage. Participants
167 are introduced to the concept of spirituality and pro-
168 vided with examples of how people with a traumatic
169 brain injury or spinal cord injury, and their family
170 members, have drawn upon different sources of spiri-
171 tuality in their adjustment. All participants completed
172 the self-study module (Module 1) before Module 2.

173 Module 2 is a 1.5 hour face-to-face workshop. It
174 includes didactic input, videod interviews with
175 former patients, the introduction of spiritual care
176 tools, and the opportunity to practise skills via role
177 plays. This content draws upon existing literature
178 and approaches to spiritual care training (Hodge,
179 2013; Puchalski & Romer, 2000). Program materi-
180 als emphasise that clients may draw upon a range of
181 sources of spiritual strength, including but not limited
182 to religious faith (Davis et al., 2015). In the role plays,
183 participants break into pairs and are provided with
184 case scenarios which depict conversations which may
185 arise with patients. They have opportunity to practise
186 taking the role of health professional or patient. Role
187 play practice incorporates exploration of the patient's
188 source of spiritual strength, the meaning this source
189 of spiritual strength currently holds for them, con-
190 nections and relationships that are important to them,
191 and how the patient would like their health profes-
192 sional to assist them to access their source of spiritual
193 strength. Participants are provided with the oppor-
194 tunity to reflect upon their own sources of spiritual
195 strength, and resources to use should they wish to
196 refer a patient for further support.

197 2.3. Measures

198 Spiritual care competency was the primary out-
199 come of interest. The Spiritual Care Competency
200 Scale (SCCS) (van Leeuwen, Tiesinga, Middel, Post,
201 & Jochemsen, 2009) is a valid and reliable 27 item
202 measure which rates participant perceptions of com-
203 petency in providing spiritual care. The 27 items
204 are scored on a five-point scale from "completely
205 disagree" to "completely agree" with total possible
206 scores ranging from 27 to 135. The scale consists
207 of six domains which measure: 1) assessment and
208 implementation of spiritual care; 2) professionalisa-
209 tion and improving the quality of spiritual care; 3)

personal support and patient counselling; 4) referral to professionals; 5) attitude towards patients' spirituality; and 6) communication. Scores are measured on a five-point scale from 1 "completely disagree" to 5 "fully agree". Cronbach's alpha for the six domains ranged from 0.56 to 0.82. The scale has good homogeneity, average inter-item correlations, and good test-retest reliability (van Leeuwen et al., 2009). It was originally designed for nursing staff, so minor adjustments were made to the wording to ensure its suitability for a wider range of professions.

Secondary outcomes of interest comprised participant levels of confidence and comfort, and attitudes and knowledge regarding spirituality and spiritual care. Participants were invited to rate their confidence and comfort levels from 0 to 10 based on the six Spiritual Care Competency Scale domains (van Leeuwen et al., 2009) listed above, with higher scores indicating higher levels of confidence or comfort. Participants' perceptions of spirituality and spiritual care were measured using the Spirituality and Spiritual Care Rating Scale (SSCRS) (McSherry, Draper, & Kendrick, 2002). The 17-item measure of spirituality and spiritual care uses a five-point scale ranging from 1 "strongly disagree" to 5 "strongly agree". A four factor model of the SSCRS (Ross et al., 2014) was used: Existential Spirituality (view that spirituality is concerned with people's sense of meaning, purpose, value, peace and creativity; 5 items); Religiosity (view that spirituality is only about religious beliefs; 3 items); Spiritual Care (view of spiritual care in its broadest sense including religious and existential elements, for example facilitating religious rituals and showing kindness; 5 items); and Personal Care (taking account of people's beliefs, values and dignity; 3 items), with one item contributing to the score of two of the subscales. Scores (total and subscale scores) are calculated by averaging the mean for the relevant items (all scores therefore range from 1–5). A broader view of spirituality and spiritual care is indicated by higher scores (Ross et al., 2014). The SSCRS has a modest level of internal consistency (Cronbach's alpha = 0.64) (McSherry, 1997) and has been used in a range of health settings, including rehabilitation (Austin, Macleod, Siddall, McSherry, & Egan, 2016).

2.4. Procedures

After signing the consent form, participants in the intervention group were provided with access to the online component (Module 1). They were then

provided with details to attend the program workshop (Module 2), which was scheduled approximately two weeks later. The scales were administered at three timepoints (pre-program, post-program, follow-up). The first timepoint (pre-program) occurred two weeks prior to Module 1, the second timepoint (post-program) immediately after Module 2, and the third timepoint (follow-up) four to six weeks after completing the training. The same measures were administered to the control group participants at the same time intervals. Data about demographic, discipline and work experience variables for both groups were collected at the pre-program timepoint. The question "Do you consider yourself a spiritual person? Please rank on a scale from 0 to 10 where 0 is 'not spiritual at all' and 10 is 'very spiritual'" was included to determine each participant's perceived level of spirituality.

2.5. Data analysis

Descriptive data were generated, and between-groups analysis at baseline on demographic variables was conducted. Multilevel models with piecewise slopes using restricted maximum likelihood (REML) were used to analyse each outcome measure over time. Each participant was considered level-2 in the models and the individual visits were level-1. A level-2 predictor representing whether the person was in the intervention or control group was added to each model. Two piecewise variables that indicate time from 1) pre intervention to post intervention and from 2) post intervention to follow-up were added to the model as level-1 variables. Interaction terms between group and each of the piecewise variables were added to assess differences in the outcomes between groups over time. Random intercepts were included in each model, and random slopes based on the piecewise variables were considered. $p < 0.05$ was considered statistically significant. The data analysis was generated using SAS Enterprise Guide software, Version 7.15 of the SAS System for Windows.

A satisfaction questionnaire measuring participant ratings of program content and usefulness was administered at the post-program timepoint. An open-ended question inviting participants to comment on the 'most significant change' they had observed since the training was added at the follow-up evaluation timepoint. A thematic analysis of this qualitative data was conducted according to guidelines provided by Braun and Clarke (2006) including familiarisation with the data; generating initial codes; searching for themes;

310 reviewing themes; defining and naming themes; and
311 producing a report.

312 3. Results

313 In relation to the intervention group, 47 reha-
314 bilitation professionals expressed initial interest in
315 participating in the training and provided consent.
316 Six of the 47 withdrew from the study prior to the
317 training due to sickness, work commitments or other
318 unexpected events, resulting in 41 rehabilitation pro-
319 fessionals who completed the training. A further 32
320 rehabilitation professionals were recruited to the con-
321 trol groups. See Fig. 1 for details of the numbers of
322 questionnaires completed at each time point by the
323 two groups.

324 Demographic details for all the participants are
325 reported in Table 2. Between group analyses (*t*-
326 test, chi square) revealed no significant differences
327 between the intervention and control groups on age,
328 gender, religious affiliation, patient group (TBI, SCI),

years of experience, or whether they considered
329 themselves to be a spiritual person.

330 For spiritual care competency, confidence, com-
331 fort, and the “existential factor” of the SSCRS,
332 pre intervention scores were not significantly dif-
333 ferent between the two groups ($p > 0.05$), however,
334 increased significantly in the intervention group at
335 post intervention ($p < 0.05$). The observed differences
336




Fig. 1. Flow diagram of the study.

Table 2
Demographic and professional details (N= 73)

Demographic items	Category	Intervention Group N= 41	Control Group N= 32
Gender (n,%)	Female	33 (80.5)	25 (78.1)
	Male	8 (19.5)	7 (21.9)
Age (n,%)	21–29	8 (19.5)	9 (28.1)
	30–39	10 (24.4)	12 (37.5)
	40–49	13 (31.7)	9 (28.1)
	50 and over	10 (24.4)	2 (6.3)
Type of patient group (n,%)	Spinal cord injury	27 (65.9)	20 (62.5)
	Traumatic brain injury	14 (34.1)	12 (37.5)
Setting (n, %)	Inpatient	33 (80.5)	32 (100.0)
	Community	8 (19.5)	
Area of expertise (discipline) (n,%)	Nursing	12 (29.3)	5 (15.6)
	Social work, psychology, case management	12 (29.3)	7 (21.9)
	Medical/other allied health	17 (41.5)	20 (62.5)
Work experience (years) (M,SD)		13.1 (9.85)	11.13 (9.77)
Qualification (n,%)	No bachelor degree	3 (7.3)	1 (3.1)
	Bachelor degree	24 (58.5)	20 (62.5)
	Master degree and above	14 (34.1)	11 (34.4)
Ethnicity (n,%)	Australian/New Zealander	27 (65.9)	19 (59.4)
	Asian	4 (9.8)	6 (18.8)
	European	5 (12.2)	7 (21.9)
	Other*	5 (12.2)	–
Born in Australia (n,%)	Yes	26 (63.4)	18 (56.3)
Religious affiliation (n,%)	None	11 (26.8)	10 (31.3)
	Christian	26 (63.4)	17 (53.1)
	Hindu	1 (2.4)	3 (9.4)
	Muslim	1 (2.4)	1 (2.4)
	Jewish	2 (4.9)	–
Previous spiritual care training (n,%)	Yes	3 (7.3)	2 (6.3)
Spiritual person 0–10 (M, SD)		6.2 (2.5)	5.4 (2.7)

Table 3
Comparison of outcomes between intervention and control groups over time

Outcome	Group	Pre-program			Post-program			Follow-up		
		Mean	Lower 95% CI	Upper 95% CI	Mean	Lower 95% CI	Upper 95% CI	Mean	Lower 95% CI	Upper 95% CI
SCCS	Intervention	94.88	91.38	98.38	112.56 ^c	109.35	115.77	111.15 ^c	107.78	114.52
	Control	94.50	90.54	98.46	92.71	88.9	96.53	92.57	88.75	96.39
Confidence	Intervention	33.22	30.18	36.25	46.95 ^c	44.08	49.82	46.97 ^c	44.02	49.93
	Control	32.22	28.78	35.65	31.90	28.52	35.28	32.76	29.38	36.13
Comfort	Intervention	36.32	33.03	39.61	47.83 ^c	44.83	50.83	47.72 ^c	44.67	50.77
	Control	33.72	30.00	37.44	32.46	28.96	35.96	32.96	29.46	36.46
Exist	Intervention	4.07	3.90	4.24	4.33 ^c	4.16	4.50	4.36 ^c	4.18	4.54
	Control	3.88	3.69	4.07	3.78	3.58	3.98	3.77	3.57	3.97
Religion	Intervention	1.82	1.65	1.99	1.64 ^a	1.48	1.81	1.63 ^c	1.45	1.81
	Control	1.94	1.75	2.13	1.94	1.74	2.15	2.10	1.89	2.30
Spiritual care	Intervention	4.31	4.15	4.48	4.69 ^c	4.56	4.81	4.57 ^c	4.44	4.71
	Control	4.18	4.00	4.37	4.10	3.95	4.25	4.13	3.98	4.28
Personal Care	Intervention	4.08 ^a	3.91	4.25	4.29 ^b	4.12	4.47	4.24 ^b	4.06	4.42
	Control	3.80	3.61	4.00	3.86	3.66	4.07	3.84	3.64	4.05

Note. CI, Confidence Interval; SCCS, Spiritual Care Competency Scale. A multilevel model was used to model the outcomes over time and compare the intervention and control groups at each time point. ^aIndicates mean is significantly different compared with the control group for that outcome at the same time point with $0.01 < p < 0.05$. ^bIndicates mean is significantly different compared with the control group for that outcome at the same time point with $0.001 < p < 0.01$. ^cIndicates mean is significantly different compared with the control group for that outcome at the same time point with $p < 0.001$.

between the groups at post intervention were maintained at follow-up. A similar trajectory was observed for the SSCRS “spiritual care” factor, with the exception of the score decreasing in the intervention group between post intervention and follow-up ($p < 0.05$), however, remaining significantly higher than the control group ($p < 0.001$). For the SSCRS factor “religion”, control group scores were significantly higher at post intervention ($p < 0.05$) and follow-up ($p < 0.001$). For the SSCRS factor “personal care”, intervention group scores were higher at pre intervention ($p < 0.05$), with differences increasing at post intervention ($p < 0.05$) and maintained at follow-up ($p < 0.05$) (see Table 3, Fig. 2).

The post-program questionnaire invited participants to rate and provide comments about workshop content and usefulness (see Table 4). Across all aspects of the workshop, the majority of participants rated the training as ‘good’ or ‘very good’. The ‘program overall’ was rated as ‘very good’ by the majority of participants. The lowest ranked aspect of the program was ‘the usefulness of the program in increasing my comfort levels’, however, most participants considered the program to be ‘good’ or ‘very good’ in raising confidence levels and in increasing knowledge and skills. Positive feedback was received regarding the introduction of a tool, the use of role plays and videos, as well as the time provided for reflection. Some participants also mentioned that the program confirmed that they

were already incorporating spiritual care into their practice.

Suggestions for improving the program included extending the duration of the program, providing information for referral to chaplaincy and other faith services, and advice on documentation in the medical record. When invited to comment on something they hoped to do better as a result of the program, many reported incorporating more meaningful questions into their practice and following clients up regarding their spiritual needs.

As part of the four to six-week follow-up, intervention group participants were invited to describe the most significant change they had noticed in their thinking or practice since the training (see Table 5). The 31 responses could be summarised by two key themes: increased awareness and understanding of spirituality as a broad concept, and increased confidence to provide spiritual care.

- i) Increased awareness and understanding of spirituality as a broad concept

Participants reported that following the program they were more aware about clients having spiritual needs, alert to the expression of spiritual needs, and more aware of the support they could provide. One participant explained:

It has helped me to be more aware of the breadth and depth of the term ‘spirituality’. To recognise in others that whilst they




Fig. 2. Comparison of intervention and control groups' scores on SCCS, and SSCRS factors.

Table 4
Workshop satisfaction ratings (N= 41)

Aspect of program	Very Poor (%)	Poor N (%)	Okay N (%)	Good N (%)	Very Good N (%)
Overall, I found the program today was	0 (0.0)	0 (0.0)	0 (0.0)	9 (21.9)	32 (78.0)
The time allocated to cover each section	0 (0.0)	0 (0.0)	1 (2.4)	15 (36.6)	25 (61.0)
The balance between theoretical and practical content	0 (0.0)	0 (0.0)	1 (2.4)	14 (34.1)	26 (63.4)
The usefulness of the content in relation to my workplace situation	0 (0.0)	0 (0.0)	1 (2.4)	11 (26.8)	29 (70.7)
The program content	0 (0.0)	0 (0.0)	1 (2.4)	14 (34.1)	26 (63.4)
The role play exercise	0 (0.0)	0 (0.0)	2 (4.9)	21 (51.2)	18 (43.9)
The level of interaction encouraged by the facilitator	0 (0.0)	0 (0.0)	0 (0)	11 (26.8)	30 (73.2)
The use of relevant language and case examples by the facilitator	0 (0.0)	0 (0.0)	1 (2.4)	7 (17.1)	33 (80.5)
The usefulness of the program in increasing my knowledge and skills	0 (0.0)	0 (0.0)	3 (7.3)	10 (23.4)	28 (68.3)
The usefulness of the program in increasing my confidence	0 (0.0)	0 (0.0)	2 (4.9)	13 (31.7)	26 (63.4)
The usefulness of the program in increasing my comfort levels	0 (0.0)	0 (0.0)	4 (9.8)	14 (34.1)	23 (56.1)

Table 5
Most significant change (N=31)

Increased awareness and understanding of spirituality	Increased confidence to provide spiritual care
<ul style="list-style-type: none"> ● Greater awareness of the broad definition of what spirituality can look like when dealing with clients ● Being more open and aware of clients' spiritual needs ● Identifying strategies for exploring spirituality during dietetic consult. Increased awareness of spiritual practices and its influence on food practices when consulting with patients. ● More aware about spirituality and used the SICA model ● I am more aware of asking clients about their spiritual needs and what gives them strength ● Being aware of what I say and do with clients and colleagues ● Increased awareness of the supports that I provide can be related to an individual spirituality and assisting my clients to consider these more actively. ● It has helped me to be more aware of the breadth and depth of the term 'spirituality'. To recognise in others that whilst they may not have a religious faith they still have a sense of well-being and connectedness that requires care, nurture and support and that we as health professionals can provide directly, support or facilitate. I appreciate more the concept of the 'whole person'. ● Being more aware of the breadth of spirituality, and looking for opportunities to assist. ● Increased awareness and confidence ● Has made me more aware of one spiritual needs in rehabilitation as little or as big as it may be. ● Greater knowledge and understanding ● I have a better understanding of what spirituality is. ● Better understanding what spirituality is and how it can be addressed in the inpatient setting ● Understanding the different aspects of spirituality and seeing how important it was to client's that it was addressed. Using a framework to assess spiritual needs, and how to assist client's during their rehab. ● Increasing my understanding of the different ways clients use spirituality as a form of hope and resilience ● Being more alert to clients expressing their spiritual needs in a variety of ways (ie language such as hope, worry) ● An important reminder to focus on the source of people's important life roles and areas of satisfaction or quality of life including spiritual beliefs. 	<ul style="list-style-type: none"> ● Making a conscious effort to implement spiritual care holistically in my practice ● I am more comfortable to acknowledge spiritual needs of clients in the context of their recovery from injury ● Being able to support clients with normalising their reflections on spiritual care post SCI and allowing time for them to explore this ● Being cognisant and supportive of a client's spiritual care needs. ● Therapy is increasingly focused on choice and control of clients ● Actively listen to clients ● Discussing spiritual needs to a greater extent. Asking more questions about what we can do to assist. If they discuss one aspect of spirituality (e.g. religion) and continue to enquire about other aspect (e.g. outside/nature). ● Realising the depth and breadth of what spirituality can involve has provided me with more confidence discussing this topic ● Providing opportunity for client to talk about spirituality in an informal way (e.g. what uplifts the client) ● Discussing spirituality during initial assessment ● Being better able to recognise a person's spiritual needs ● Actively listening to patients about spirituality ● Open discussions with work colleagues regarding concept of spirituality & supporting our clients

396 may not have a religious faith they still have
 397 a sense of well-being and connectedness
 398 that requires care, nurture and support, and
 399 that we as health professionals can provide
 400 directly, support or facilitate. I appreciate
 401 more the concept of the 'whole person'.

402 Another participant expressed that the train-
 403 ing had increased their understanding of "*the*
 404 *different ways clients use spirituality as a form*
 405 *of hope and resilience*". Another mentioned how
 406 understanding different aspects of spirituality
 407 had helped them to realise how important spiri-
 408 tuality is to clients.

409 ii) Increased confidence to provide spiritual care

410 The second identified theme was partici-
 411 pants feeling more confident to provide spiritual

412 care. One participant reported that the train-
 413 ing had helped them to support clients by
 414 "*normalising their reflections on spiritual care*
 415 *post SCI and allowing them to explore this*".
 416 Another mentioned that they felt more com-
 417 fortably acknowledging the spiritual needs of
 418 clients within the context of their recovery. Two
 419 participants reported that they were actively
 420 making a conscious effort to implement spiri-
 421 tual care or identify strategies for exploring
 422 spirituality in their work. One mentioned that
 423 they were "*asking more questions about what*
 424 *we can do to assist*" and exploring more than
 425 one source of spirituality with clients (for
 426 example, the natural world as well as religious
 beliefs).

4. Discussion

This exploratory study evaluated the effectiveness of a brief spiritual care training program to expand attitudes and knowledge regarding spirituality and spiritual care, and to increase rehabilitation professionals' perceived levels of competency, confidence and comfort in the delivery of spiritual care. To the best of our knowledge no other studies have trialled healthcare professional training in the area of spiritual care and neurorehabilitation. Significant increases in the primary outcome, spiritual care competency, were recorded for the intervention group at the post program timepoint and were not matched by the control group. These increases were maintained at follow-up. Similarly, in relation to the secondary outcomes the intervention group scored significantly higher scores on confidence and comfort, and demonstrated a greater understanding of spirituality and spiritual care at the post program timepoint than the control group. Participant satisfaction levels regarding the program content and usefulness were high. At follow-up participants in the intervention group could identify changes in both understanding and practice in their delivery of spiritual care.

Levels of spiritual care competency were significantly higher for the intervention group after attending the spiritual care training program. Increases in spiritual care competency have been reported in other studies investigating the effects of spiritual care training. A recent study by Pearce, Pargament, Oxhandler, Vieten and Wong (2019) with mental health providers (the majority of whom were psychologists, social workers, counsellors) found an online training program to be successful in improving spiritual care competencies. Spiritual care competencies improved at post-testing, after an eight-module online training program. Although the current program was much less time-intensive, similar results were achieved and maintained at follow-up, suggesting that even a small amount of training can bring about significant change.

Confidence and comfort levels in delivering spiritual care were also significantly higher for the intervention group at post and follow-up testing. Other spiritual care programs have indicated that brief training in spiritual care can be effective in improving confidence and comfort levels (Cerra & Fitzpatrick, 2008; Meredith et al., 2012). Perspectives on spirituality and spiritual care also changed for the intervention group, evident from participant scores on the SSCRS (McSherry et al., 2002). Compared

with the control group, the factors most likely to indicate change were participant ratings on Existential Spirituality (view that spirituality is concerned with people's sense of meaning, purpose, value, peace and creativity) and Spiritual Care (view of spiritual care in its broadest sense including religious and existential elements, for example facilitating religious rituals and showing kindness). These findings aligned well with the content of the program which encouraged participants to adopt broad definitions of spirituality and spiritual care in their practice. The factors which did not change were Religiosity (view that spirituality is only about religious beliefs) and Personal Care (taking account of people's beliefs, values and dignity). In fact, the scores for the control group on Religiosity were higher than the intervention group at post and follow-up timepoints. High scores on this item suggest that participants are more likely to hold the view 'that spirituality is only about religious beliefs'. Therefore, lower results for the intervention group around religiosity fit with the program content, which actively discouraged participants from considering spirituality as interchangeable with religion. Respecting people's beliefs, values and dignity is well incorporated into most healthcare professional training, and therefore little difference between groups on this factor was not surprising.

The qualitative findings of this study enrich the quantitative findings. Answers to the question about "most significant change" at follow-up suggest that shifting perceptions of spirituality and spiritual care may be linked with levels of confidence or comfort. Such a finding reinforces the notion that a small change in attitude and understanding can bring about benefits that extend beyond knowledge alone. Participant satisfaction levels were high, suggesting that a two-module program had good levels of acceptability for staff, and the program was well attended. The majority of healthcare professionals undertaking the training completed both modules.

This study had a number of limitations. Although this was a controlled trial, participants were not randomised to the two conditions. The intervention was delivered at one site and the number of staff participating in the training was modest. Participant skills or behaviour change were only evaluated by self-report at the follow-up time point and we were unable to determine whether clients reported any changes as a result. Furthermore, due to the time constraints of the project, a six-week follow-up period was decided upon. This did not allow all the participants to apply what they had learnt from the program. Despite these

530 limitations, the findings of this exploratory study
 531 would support larger, randomised controlled trials
 532 of spiritual care education programs in the field of
 533 rehabilitation.

534 5. Conclusion

535 The program's underlying principles was that
 536 spiritual care can be provided by all healthcare pro-
 537 fessionals and is relevant to staff in all areas of
 538 healthcare, including neurorehabilitation. Training
 539 which increases staff competency, comfort, confi-
 540 dence and understanding regarding the delivery of
 541 spiritual care will enhance the ability of healthcare
 542 services to embrace the needs of the whole person.
 543 That this training can be achieved over a brief period
 544 of time is promising and suggests that training need
 545 not be time-intensive or arduous for participants.
 546 Future research could expand the findings of this
 547 study by incorporating larger trials of the program and
 548 with rehabilitation professionals from a wider range
 549 of religious faith backgrounds. Intervention programs
 550 which address spirituality with rehabilitation clients
 551 and their family members directly would also be wor-
 552 thy of consideration. Such research will contribute to
 553 a growing acknowledgement that incorporating spir-
 554 itual care into rehabilitation practice is both valuable
 555 and achievable.

556 Acknowledgments

557 The authors would like to thank all the health pro-
 558 fessionals who participated in this study, and Royal
 559 Rehab, Liverpool Brain Injury Rehabilitation Unit,
 560 Prince of Wales Spinal Injuries Unit and Community
 561 of Christ for their support. We would also like to thank
 562 Jackie Francis for her assistance on this project.

563 Conflict of interest

564 The authors report no conflicts of interest.

565 References

566 Austin, P., Macleod, R., Siddall, P. J., McSherry, W., & Egan,
 567 R. (2016). The ability of hospital staff to recognise and meet
 568 patients' spiritual needs: a pilot study. *Journal for the Study of*
 569 *Spirituality*, 6(1), 20-37.

- Best, M., Butow, P., & Olver, I. (2016). Palliative care specialists' 570
 beliefs about spiritual care. *Support Care Cancer*, 24, 3295- 571
 3306. 572
- Braun, V., & Clarke, V. (2006). Using thematic analysis in psy- 573
 chology. *Qualitative Research in Psychology*, 3, 77-101. 574
- Canda, E. R., & Furman, L. D. (2009). *Spiritual diversity in social 575*
work practice: the heart of helping. Cary, NC: Oxford Univer- 576
 sity Press. 577
- Cerra, A., & Fitzpatrick, K. (2008). Can in-service education help 578
 prepare nurses for spiritual care? *Journal of Clinical Nursing*, 579
 25(4), 204-209. 580
- Cobb, M., Puchalski, C. M., & Rumbold, B. (Eds.). (2012). *Oxford 581*
textbook of spirituality in healthcare. Oxford: Oxford Univer- 582
 sity Press. 583
- Davis, D. E., Rice, K., Hook, J. N., Van Tongeren, D. R., DeBlaere, 584
 C., Choe, E., & Worthington, E. (2015). Development of the 585
 sources of spirituality scale. *Journal of Counselling Psychol- 586*
ogy, 62(3), 503-513. 587
- Fricchione, G., & Nejad, S. (2012). Resilience and coping. In M. 588
 Cobb, C. Puchalski, & B. Rumbold (Eds.), *Oxford Textbook 589*
of Spirituality in Healthcare (pp. 367-373). Oxford: Oxford 590
 University Press. 591
- Gallison, B., Xu, Y., Jurgens, C., & Boyle, S. (2012). Acute care 592
 nurses' spiritual care practices. *Journal of Holistic Nursing*, 593
 31(2), 95-103. 594
- Hodge, D. R. (2013). Implicit spiritual assessment: An alternative 595
 approach for assessing client spirituality. *Social Work*, 58(3), 596
 223-230. 597
- Johnstone, B., Glass, B. A., & Oliver, R. E. (2007). Religion 598
 and disability: clinical, research and training considerations 599
 for rehabilitation professionals. *Disability and Rehabilitation*, 600
 29(15), 1153-1163. 601
- Jones, K. F., Dorsett, P., Briggs, L., & Simpson, G. (2018). The 602
 role of spirituality in spinal cord injury (SCI) rehabilitation: 603
 exploring health professional perspectives. *Spinal Cord Series 604*
and Cases, 4(1), 54. doi:10.1038/s41394-018-0078-3 605
- Jones, K. F., Dorsett, P., Simpson, G., & Briggs, L. (2018). Moving 606
 forward on the journey: spirituality and family resilience after 607
 spinal cord injury. *Rehabilitation Psychology*, 63(4), 521-531. 608
 doi:10.1037/rep0000229 609
- Jones, K. F., Pryor, J., Care-Unger, C., & Simpson, G. K. (2018). 610
 Spirituality and its relationship with positive adjustment fol- 611
 lowing traumatic brain injury: a scoping review. *Brain Injury*, 612
 1-11. doi:10.1080/02699052.2018.1511066 613
- Jones, K. F., Pryor, J., Care-Unger, C., & Simpson, G. K. (2020). 614
 Rehabilitation health professionals' perceptions of spirituality 615
 and spiritual care: The results of an online survey. *NeuroReha- 616*
ilitation, 46(1), 17-30. doi:10.3233/NRE-192857 617
- Jones, K. F., Simpson, G. K., Briggs, L., & Dorsett, P. (2016). Does 618
 spirituality facilitate adjustment and resilience among individ- 619
 uals and their families after SCI? *Disability and Rehabilitation*, 620
 38(10), 921-935. 621
- Koenig, H. G. (2012). Religion, spirituality, and health: the 622
 research and clinical implications. *ISRN Psychiatry*, 2012, 623
 1-33. 624
- McSherry, W. (1997). *A descriptive survey of nurses' percep- 625*
tions of spirituality and spiritual care. The University of Hull, 626
 Hull. 627
- McSherry, W., Draper, P., & Kendrick, D. (2002). The construct 628
 validity of a rating scale designed to assess spirituality and 629
 spiritual care. *International Journal of Nursing Studies*, 39, 630
 723-734. 631

- 632 Meredith, P., Murray, J., Wilson, T., Mitchell, G., & Hutch, R.
633 (2012). Can spirituality be taught to health care professionals?
634 *Journal of Religion and Health*, 51, 879-889.
- 635 NHS Education for Scotland. (2009). *Spiritual Care Matters*. Edinburgh,
636 UK: NHS Education for Scotland.
- 637 NHS Education for Scotland. (2013). Spiritual Care. Retrieved
638 from [https://www.nes.scot.nhs.uk/education-and-training/by-](https://www.nes.scot.nhs.uk/education-and-training/by-discipline/spiritual-care.aspx)
639 [discipline/spiritual-care.aspx](https://www.nes.scot.nhs.uk/education-and-training/by-discipline/spiritual-care.aspx)
- 640 Oakley, E. T., Katz, G., Sauer, K., Dent, B., & Millar, A. (2010).
641 Physical therapists' perception of spirituality and patient care:
642 beliefs, practices, and perceived barriers. *Journal of Physical*
643 *Therapy Education*, 24(2), 45-52.
- 644 Oxhandler, H., Parrish, D., Torres, L., & Achenbaum, W. (2015).
645 The integration of clients' religion and spirituality in social
646 work practice: a national survey. *Social Work*, 60(3), 228-237.
- 647 Paal, P., Helo, Y., & Frick, E. (2015). Spiritual care training pro-
648 vided to healthcare professionals: A systematic review. *Journal*
649 *of Pastoral Care and Counselling*, 69(1), 19-30.
- 650 Pearce, M. J., Pargament, K., Oxhandler, H., Vieten, C., & Wong,
651 S. (2019). A novel training program for mental health providers
652 in religious and spiritual competencies. *Spirituality in Clinical*
653 *Practice*, 6(2), 73-82. doi:10.1037/scp0000208
- 654 Puchalski, C., Ferrell, B., Virani, R., Otis-Green, S., Baird, P.,
655 Bull, J., ... Sulmasy, D. (2009). Improving the quality of
656 spiritual care as a dimension of palliative care: The report of
657 the consensus conference. *Journal of Palliative Medicine*, 12,
658 885-904.
- 659 Puchalski, C., & Romer, A. L. (2000). Taking a spiritual history
660 allows clinicians to understand patients more fully. *Journal of*
661 *Palliative Medicine*, 3(1), 129-137.
- 662 Ross, L., Leeuwen, R., Baldacchino, D., Giske, T., McSherry, W.,
663 Narayanasamy, A., ... Schep-Akkerman, A. (2014). Student
664 nurses perceptions of spirituality and competence in delivering
665 spiritual care: A European pilot study. *Nurse Education Today*,
666 34, 697-702.
- 667 Smith, W. B., Ortiz, A. J., Wiggins, K. T., Bernard, J. F., & Dalen,
668 J. (2012). Spirituality, resilience, and positive emotions. In
669 L. J. Miller (Ed.), *The Oxford Handbook of Psychology and*
670 *Spirituality* (pp. 437-454). Oxford: Oxford University Press.
- 671 van Leeuwen, R., Tiesinga, L., Middel, B., Post, D., & Jochem-
672 sen, H. (2009). The validity and reliability of an instrument
673 to assess nursing competencies in spiritual care. *Journal of*
674 *Clinical Nursing*, 18, 2857-2869.
- 675 Walsh, F. (2003). Family resilience: a framework for clinical practice. *Family Process*, 42(1), 1-18.
- 676 White, B., Driver, S., & Warren, A. (2008). Considering resilience
677 in the rehabilitation of people with traumatic disabilities. *Reha-*
678 *bilitation Psychology*, 53(1), 9-17.
- 679