

# Aging research using the common marmoset: Focus on aging interventions

Corinna N. Ross<sup>a,b,\*</sup> and Adam B. Salmon<sup>b,c,d,\*</sup>

<sup>a</sup>*Department of Science and Mathematics, Texas A&M University San Antonio, San Antonio, TX, USA*

<sup>b</sup>*The Sam and Ann Barshop Institute for Longevity and Aging Studies and*

<sup>c</sup>*Department of Molecular Medicine, The University of Texas Health Science Center at San Antonio, San Antonio, TX, USA*

<sup>d</sup>*Geriatric Research, Education and Clinical Center, South Texas Veterans Health Care System, San Antonio, TX, USA*

**Abstract.** Traditional animal models have been used to make seminal discoveries in biomedical research including a better understanding of the biology of the aging process. However, translation of these findings from laboratory to clinical populations has likely been hindered due to fundamental biological and physiological differences between common laboratory animals and humans. Non-human primates (NHP) may serve as an effective bridge towards translation, and short-lived NHP like the common marmoset offer many advantages as models for aging research. Here, we address these advantages and discuss what is currently understood about the changes in physiology and pathology that occur with age in the marmoset. In addition, we discuss how aging research might best utilize this model resource, and outline an ongoing study to address whether pharmaceutical intervention can slow aging in the marmoset. With this manuscript, we clarify how common marmosets might assist researchers in geroscience as a potential model for pre-clinical translation.

**Keywords:** Marmoset, non-human primate, mTOR, rapamycin, longevity, resilience

## 1. Challenges of translation from laboratory animals (mostly rodents) to humans

Animal models have been a critical resource in the quest to evaluate physiological functional decline associated with aging and elucidate the cellular pathways and mechanisms associated with longevity. In general, these studies have focused most of their efforts on delineating the aging process using four primary laboratory organisms: yeast (*Saccharomyces cerevisiae*), roundworms (*Caenorhabditis elegans*), fruit flies (*Drosophila melanogaster*) and rodents including mice and rats (*Mus musculus* and *Rattus norvegicus*, respectively) [1]. These species have many practical advantages over other animal models

(including humans) as models of aging including their relatively short lifespan, the ability to breed and maintain large numbers of animals in the laboratory, and the ability to control environmental exposure for each individual (or a population). In addition, over the last few decades it has been shown clearly that the relative ease of manipulating genes of interest; *i.e.*, generating organisms with specific genetic mutations to address mechanisms of phenotype including longevity, is a powerful tool for the study of aging. Using these models has been extremely fruitful for geroscience research and has driven discoveries of conserved biochemical pathways associated with longevity, and allowed the comparative evaluation of the effect of aging on metabolism and development, mitochondrial stress pathways, adenosine monophosphate-activated protein kinase, and insulin-like growth factor [2, 3]. In particular, the widespread use of mice, and in particular genetic

\*Corresponding author: Corinna N. Ross, PhD. E-mail: cnross@tamusa.edu and Adam B. Salmon, PhD. E-mail: salmona@uthscsa.edu.

mutant lines of mice, have driven exponential growth in our understanding of the particular mechanistic pathways on the mammalian aging process [4]. In terms of “translational” approaches to addressing aging and longevity in humans, mice and rats have served a central role in identification and elucidation of interventional and therapeutic treatments on aging and health-span. Most notably, the gold standard for extending lifespan in a model organism is the use of caloric restriction and indeed the seminal studies of caloric restriction have largely been performed using mice and rats [5–9]. In addition, many new drugs of geroscience interest have been tested in rodents and continue to reveal details regarding aging and disease progression, and have yielded novel findings about the anti-aging effect of intervention with pharmaceutical and nutraceutical compounds [10–17]. In many ways, interventional approaches utilizing pharmaceuticals seem to be the most “turn-key” for benefiting human health in terms of their relative ease of translation for clinical consideration.

However, there remain many concerns regarding using rodents as a model of human model of aging (in addition to disease, physiology, pathology, *etc.*) as a foundation for designing treatments and interventions to improve *human* health. Unfortunately, a litany of findings in rodent translational research have revealed that many of the drugs tested in rodents do not translate successfully to humans, either having few similar effects or having toxicity in humans [18]. Further, the development of inbred strains of mice has introduced the perpetuation of many traits that are a by-product of selection for breeding, rather than traits of interest. Many of these traits are undetected or unknown and may be detrimental to longevity studies or have unknown impact on the research of interest [19]. Lastly, there is a distinct dichotomy between rodents and humans in terms of the spectrum of diseases that naturally occur with old-age or otherwise. Particularly challenging are the wide-range of diseases and pathologies that plague aging humans that do not naturally exist in rodents (*e.g.*, neurodegenerative diseases including Alzheimer’s disease, cardiovascular disease, diabetes, and a host of others) due to differences between species in their basic physiology, the dramatic differences in their maximum ages, diet and environmental conditions or other unknown variables. Specifically because of such challenges, there has been significant investment in drawing from comparative biology to identify additional animal models that may serve as stronger bridges to clinical translation.

## 2. How non-human primates are useful (critical) to this process

Non-human primate (NHP) models are of particular interest for additional pre-clinical testing towards translation due to their close evolutionary history with humans. Old World monkeys and apes are the closest living relatives to humans and share a closer evolutionary history with humans than other mammals, and thus more similar genetic, biochemical, behavioral and phenotypic outcomes. Old world monkeys split from the ancestors of the human lineage around 25 million years ago, and apes, specifically chimpanzees, split as recently as six million years ago [20, 21]. In the limited studies that have examined the aging characteristics of captive chimpanzees, our closest living relatives, the age-related diseases in this species are described as similar to that of humans including the development of heart disease, cancer, and diabetes with advancing age [22–29]. However, chimpanzees are very long lived, with animals in captivity living up to 60 years, thus making them comparable in difficulty to working with human populations [21]. Additionally, both wild and captive populations of chimpanzees are now classified by the International Union for Conservation of Nature (IUCN) as endangered animals and the National Institutes of Health (NIH) have effectively ceased funding for invasive research on chimpanzees. In part for these reasons, research using chimpanzees in the laboratory setting has significantly diminished in many countries including the U.S.

Other Old World monkeys, including rhesus macaques, have traditionally been of greater focus for biomedical translational research including aging studies [30–35]. Rhesus macaques have a maximum lifespan of approximately 40 years in captivity and their aging phenotypes have been well characterized [36, 37]. In addition, a means to monitor cognitive functional decline in this species has been developed and is hypothesized to model the similar decline in human cognition with age [32]. The rhesus macaque has also been used to test whether the benefits of caloric restriction on healthy aging translate to NHP species [30]. While there continues to be some discussion on the equivocal nature of caloric restrictions’ effect on NHP lifespan in two ongoing research groups, these studies have largely confirmed that calorie restriction reduces the prevalence of cardiovascular disease, type 2 diabetes, and neoplasias [38–45]. Due in part to the long lifespan of this species and the large time and financial commitment

to these studies, it is unlikely that such in depth NHP lifespan studies will be able to be repeated.

In an attempt to reduce the heavy commitments required for aging NHP studies, several shorter-lived NHP species have been suggested for longevity research including the bush baby (*Galago senegalensis*), the grey mouse lemur (*Microcebus murinus*), and the common marmoset (*Callithrix jacchus*) [46, 47]. These species all have relatively shorter lifespans (for each, the maximum lifespan is ~15–20 years) compared to Old World monkeys and apes, and they offer a number of advantages due to their small size and rapid reproduction [46, 47]. Of note, it was recently reported that the lifespan of the grey mouse lemur could be extended by calorie restriction but that this intervention did not alter cognitive or motor function [48]. However, of these species, the common marmoset has recently emerged as a model for aging related research due to growing investigations on the changes in physiology, pathology and health that occur in this species [49–54].

### 3. Marmosets in aging research

The common marmoset has many characteristics that make it attractive as a non-human primate model for aging research. As mentioned above, this species has a lifespan that is roughly half that of other NHP species commonly used in biomedical research (Fig. 1). While still quite long compared to rodents, this length of time is certainly more amenable to use within the career of a single researcher. Moreover, there is fiduciary benefit to the shorter lifespan (*i.e.*, half the amount of time paying per diems as other long-lived NHP). Along those lines, per diem costs tend to be significantly lower for marmosets than for other NHP, due in part to the smaller cage and space requirements. Their small size also has benefits in terms of pharmaceutical interventions (*i.e.*, less drug is required per animal) as well as in terms of staff safety. Marmosets are also the fastest reproducing anthropoid primate giving birth to fraternal twins on average twice per year [55]. This fast reproduction allows for relatively rapid growth of colonies for use in biomedical research. Further, these characteristics allowed for the development of an specific pathogen free (SPF) barrier maintained colony of marmosets for aging research which to date has displayed extended lifespan when compared to conventionally housed marmosets [56]. These features




Fig. 1. Maximum reported lifespans for four primary primate species used in aging research.

then support the notion that marmosets offer a unique NHP model with many advantages for studying the physiological changes associated with aging.

As part of the development of this species as an aging model, there is a growing understanding of the basic aging phenotypes of aging marmosets. For example, a number of skeletal changes associated with aging have been described in marmosets. Marmosets lack adult growth plates and display similar human age related structural changes in skeletal bone [57]. When treated with a bisphosphate therapy, trabecular number and volume increased in marmosets exhibiting age-related bone loss [57]. Further, marmoset bones undergo degeneration of structure associated with bone aging in such bones as the vertebrae [58].

With advancing age, marmosets have been reported to lose weight and show dramatic changes in body composition [54, 59]. These patterns are very similar to the types of changes in body weight and composition that occur in the latter half of normal lifespan in humans [60]. Interestingly, the maximum weight achieved by an individual marmoset is significantly associated with the likelihood of survival in older animals, with animals having a peak mass of less than 400 g having decreased survival after eight years of age [54]. Perhaps related to this, marmosets that exhibit increased rates of stretching behavior have a significantly higher 6-month mortality than animals do not display this behavior [50]. Rather than a beneficial behavior, this type of stretching is hypothesized to be a posture adopted due to discomfort in the marmoset. It is an atypical posture in which the marmosets hang for extended periods of time from their forelimbs rather than maintaining a more typical marmoset quadrupedal positioning. It is still unclear

whether this can be attributed to a specific pathology but it would certainly be of further interest to identify how marmoset activity and ambulatory behavior are impacted by physiological declines associated with aging.

There has also been increasing interest in marmosets as a model to evaluate a number of age-related neurological diseases including Parkinson's disease, Huntington's disease, Alzheimer's disease, stroke, multiple sclerosis and spinal cord injury. Marmosets exhibit decreased adult neurogenesis in the dentate gyrus that is evident before the animals would typically be classified as aged [61]. Ultrastructural examinations of the frontal cortex and hippocampus for animals more than 12 years of age found widespread accumulation of lipofuscin in the glial cells, perivascular macrophages and pericytes [62]. A recent study described an increase in  $\alpha$ -synuclein aggregations in the olfactory bulb and hippocampus of aged marmosets [63].  $\beta$ -amyloid deposition has been described in the brain of marmosets over the age of seven [64], and this data is often used as a criterion for defining marmosets as aged when they are eight years old. However, Ridley [65] did not detect  $\beta$ -amyloid in animals under the age of 10 in their colony and very little deposition in older animals. Marmoset brains have also been evaluated for the presence of amyloid beta markers and tau hyperphosphorylation [66] revealing diffuse amyloid plaques throughout the cortex of the aged marmosets, but not in the younger age groups. Conformational changes in tau were detected in all subjects in this study but these changes increased in frequency with aging. Dystrophic microglia were also significantly more likely to exhibit tau hyperphosphorylation than were active microglia in this study. However, the inconsistency between the presence of neurological markers at specific ages in the marmosets suggests that there may be environmental differences associated with the rate of aging between the colonies that have been examined.

As a potential translational model, there have also been numerous attempts to apply standard human clinical biomarkers of aging to marmosets. To date the only reported change in marmoset blood chemistry associated with age has been a significant decrease in albumin with advancing age [50]. Serum albumin decreases are of particular interest as they are a highly predictive of a risk of death in otherwise healthy aging humans [67]. In the largest study to date of age-related changes in marmoset blood chemistry, a cross-sectional analysis of 60 marmosets ranging in

age from two to 13 years old identified 2500 metabolites [68, 69]. Connectivity between the metabolites was found to decrease with age and the abundance of several of the metabolites significantly declined with age [70]. With age, marmosets tend to display increased mean arterial pressure and diastolic pressure, but no age effect on the systolic pressure [71]. Finally, testosterone has been found to decrease in aging male marmosets, but they remain capable of reproduction, and no known phenotypic changes are associated with the decreased testosterone [55, 72–74].

An analysis of the pathologies associated with age in marmosets at the New England National Primate Center and the Southwest National Primate Research Center revealed dramatic shifts in the associated causes of death as animals aged in the colony [50, 54]. Deaths in young adults under the age of 6 years were most likely to be due to injury, inflammatory bowel disease (IBD) and infection. Both colonies reported rare occurrences of neoplasia and diabetes. In contrast, for marmosets over the age of 6 years the most likely causes of death were infection and IBD, with increasing rates of pathologies typically thought of as age related pathologies in humans and other model organisms including neoplasia, amyloidosis, diabetes, cardiac and renal failure. It must be noted that IBD is not a primary cause of mortality in humans and this reflects one of the challenges in using the marmoset as an aging model. On the other hand, the etiology of IBD in the marmoset has been difficult to ascertain and may be driven in part by susceptibility to infection in the community housing setting. Supporting this idea, we recently reported that marmosets maintained in SPF conditions showed no deaths associated with inflammatory gastrointestinal or infectious disease unlike marmosets maintained in a standard colony setting [56]. A brief summary and comparison of most likely causes of adult death among mice, marmosets, and humans are provided in Fig. 2.

While the extent of physiological and behavioral markers that have been evaluated for changes with aging is ever expanding, many things remain to be examined. For example, a common evaluation to assess frailty or health in geriatric patients is the use of walking speed and grip strength. Translating these relatively simple human assessments to marmosets has proven to be challenging. Other indicators of physiological and social resilience are also still lacking and need further refinement and development to determine their trajectories with age. Tools to assess




			
<b><u>Species</u></b>	<b><i>M. musculus</i></b>	<b><i>C. jacchus</i></b>	<b><i>H. sapiens</i></b>
<b><u>Max lifespan</u></b>	<b>3-4 y.</b>	<b>15-20 y.</b>	<b>&gt;120 y.</b>
<b><u>Most common causes adult mortality</u></b>	<b>Neoplasia</b>	<b>Kidney disease Heart disease Cancer Hepatitis Amyloidosis Inflammatory bowel/wasting</b>	<b>Heart disease Neoplasia Chronic respiratory Stroke Alzheimer's disease Diabetes Kidney disease</b>

Fig. 2. Common causes of death among laboratory mice, laboratory marmosets, and humans (clipart images from www.istockphotos.com).

cognitive health are not as varied and detailed in the marmoset as they are for human evaluations. Many of the cognitive tools used to evaluate Old World monkeys or apes are beyond the ability of the marmoset to be trained for or to complete, and interpretations of results can be controversial. Development of new cognitive tests and adaptations of human, monkey and mouse assessments are ongoing developments.

#### 4. Scientific approaches to study aging in marmosets

It is now clear that longevity can be altered using three primary means in traditional laboratory animals: genes, diet, and pharmaceutical interventions [13, 75, 76]. These interventional methods are invaluable tools for studying the basic biology of aging and there is growing evidence that they can be translated to marmoset studies with some considerations.

While genetic modifications have been valuable tools in invertebrate and mouse models of aging, one of the biggest disadvantages historically for NHP biomedical studies has been the inability to produce genetically modified individuals to examine targeted areas of interest in ways that are either relatively cheap or easy (or hopefully both). There are a few reports of transgenic Old World monkeys that exist

but the production of a single living offspring in these species is generally extraordinarily costly both financially and in terms of time commitment [77, 78]. Moreover, in terms of aging studies the phenotypic symptoms of the gene of interest may not appear until mid- to old-age which could be on the orders of decades. However, transgenic marmosets have existed for nearly a decade with the production of the lentiviral-induced GFP transgenic marmosets first reported in 2009 [79]. In this mutant monkey, the GFP transgene was distributed throughout the somatic cells in addition to a successful germ-line transmission that was verified within 2 years of the birth of the initial infants. Since this first report, a number of facilities have produced transgenic lines of marmosets using lentiviral transduction, CRISPR/Cas and Tet-on systems. A process that has a great deal of potential for the production of transgenic marmosets is somatic cell nuclear transfer (SCNT) because it may decrease the likelihood of producing mosaic individuals, but to date these attempts have not been successful [80]. Regarding aging research, many of the transgenic lines proposed and developed using marmosets have focused on neurologic disease and modeling of neurodegenerative disorders such as Alzheimer's and Parkinson's. Lentiviral transgenic induction of Tet-on human ataxin 3 genes has produced young marmosets with neurodegenerative disease phenotype [81, 82]. Recently, marmosets that express transgenic

343  
344  
345  
346  
347  
348  
349  
350

351  
352

353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367

368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396

397 induction of intracellular calcium indicators have  
398 been developed in order to greatly advance the tech-  
399 nology and imaging possibilities [83]. While the  
400 developing transgenic marmoset models is still cer-  
401 tainly much more expensive and time consuming than  
402 developing similar rodent models, the ability to exam-  
403 ine neurodegenerative disorders and cognitive deficit  
404 in a NHP will greatly advance our understanding of  
405 these processes in human aging. Further refinement of  
406 transgenic marmoset generation will also help expand  
407 the scientific focus of such models to other areas of  
408 aging research.

409 As mentioned above, calorie restriction has been  
410 one of most important tools used in aging research  
411 to understand the basic mechanisms of animal  
412 aging. Dietary interventions, including both dietary  
413 restriction and over-nutrition, have been adapted to  
414 marmoset studies to test the physiological outcomes  
415 of such interventions [84, 85]. However, these effects  
416 have not been evaluated in association with aging  
417 or age-related disease to date. Dietary restriction  
418 in marmosets has been limited to evaluations of the  
419 timing of caloric restriction during pregnancy and lac-  
420 tation to infant outcome [84]. While it is possible to  
421 restrict marmosets calorically, for some short studies  
422 such as measuring food intake or evaluation of daily  
423 food patterning, it is extremely challenging to use  
424 this intervention under normal housing conditions for  
425 this animal; *i.e.*, social housing. Marmosets are one of  
426 the few primates that has been found to actively share  
427 food with other family members and with infants [86].  
428 This particular trait of marmosets makes it difficult to  
429 account for food taken from a hopper by an individ-  
430 ual animal because even though they remove it from  
431 a dish, they are not necessarily eating the food by  
432 themselves. Single housing animals, as would typ-  
433 ically be done in rodent calorie restriction, is not  
434 preferred in marmosets due to the social habits (and  
435 requirements) of this species. One potential approach  
436 to food restriction might be separating the individual  
437 of interest during feeding and then removing all food  
438 from the family housing prior to returning the animal.  
439 However, this technique is extremely labor intensive  
440 and it is likely that the entire group's feeding patterns  
441 would be shifted in ways that would be challenging  
442 to document. Another possible approach would be  
443 to use time-restricted feeding or intermittent fasting,  
444 though the status of these approaches in comparison  
445 to "traditional" calorie restriction is still equivocal.  
446 Thus, there is still opportunity to develop appropri-  
447 ate caloric restriction paradigms in the marmoset for  
448 future testing.

449 Pharmaceutical interventions are by far the most  
450 easily translated to marmoset studies of aging and  
451 age-related disease. The marmoset is already a valu-  
452 able NHP model for evaluating preclinical drug  
453 pharmacokinetics, teratology and toxicity in part due  
454 to its small size and thus need for smaller doses  
455 of drugs compared to other NHP [87–89]. Their  
456 metabolic profile and primate physiology allows  
457 drugs to be tested in the same delivery modality  
458 as will be used in human dosing, making transla-  
459 tion to clinical trials smoother. Marmosets have been  
460 key to the development of drugs for use in neu-  
461 roscience, immunology and infectious disease [89].  
462 Drug development and testing for multiple sclero-  
463 sis, Parkinson's, stroke, spinal injury, amyloidosis,  
464 hepatitis, encephalitis, viral hemorrhagic fevers and  
465 bone disease have all relied on marmosets [89]. Thus,  
466 an easy case can be made for using this particular  
467 approach as the "low-hanging fruit" for refinement  
468 of the marmoset as a valuable NHP for studying the  
469 basic biology of aging. In this regard, we have put  
470 together an aging cohort of marmosets to test the very  
471 idea of whether drug intervention can delay the aging  
472 process in this NHP.

## 473 5. Designing a study to address whether 474 rapamycin slows aging in marmosets

475 The work of the Interventions Testing Program  
476 and others has identified rapamycin as one of the  
477 most consistent (so far) pharmaceutical interventions  
478 capable of extending lifespan and improving health  
479 in mice [13–15, 17, 90]. Rapamycin is an inhibitor of  
480 mTOR (mechanistic target of rapamycin) signaling  
481 and this compound, along with some of its analogues,  
482 have been shown to delay (if not rejuvenate) the  
483 physiological dysfunction and pathologies associated  
484 with aging across multiple organs in the aging mouse  
485 [90–96]. Collectively, these outcomes have provided  
486 the first proof of principle that longevity in mammals  
487 can be extended, if not slowed outright, by admin-  
488 istration of a pharmaceutical agent. As described  
489 above, there are significant challenges in applying the  
490 results of intervention testing in rodents towards the  
491 benefit of human health. There is some evidence that  
492 rapamycin or its orthologs might provide some bene-  
493 fit to generally healthy older human populations with  
494 limited side-effects [97, 98]. While clinical based trial  
495 may reveal additional benefit, they are unlikely in  
496 the near future to be capable of determining whether  
497 this intervention increases longevity or slows aging

498 among humans directly. As discussed here, there  
499 then is potentially tremendous benefit in addressing  
500 whether interventions that extend longevity in mice,  
501 such as rapamycin, similarly extend lifespan in a NHP  
502 species.

503 To this end, we have enrolled a cohort of middle-  
504 aged marmosets into a long-term study to test the  
505 effect of rapamycin on longevity and healthy aging in  
506 this species. During the development of this project,  
507 we previously characterized the pharmacokinetics of  
508 daily oral treatment with rapamycin encapsulated  
509 in a slow-release enteric coating (as used in the  
510 Intervention Testing Program's mouse studies). We  
511 reported that marmosets treated with an oral dose of  
512 1.0 mg/kg body weight/day rapamycin showed cir-  
513 culating trough rapamycin concentrations similar to  
514 those reported in mouse lifespan studies as well as  
515 similar to the reported therapeutic ranges for humans  
516 given rapamycin as chemotherapy [99]. Moreover, we  
517 showed this dose of rapamycin was sufficient to inhibit  
518 mTOR signaling *in vivo* both in the short (weeks) and  
519 long (14 months) term. In this pilot study, we also  
520 reported little to no signs of intolerability among ani-  
521 mals treated with rapamycin (total 7 animals of mixed  
522 sexes) and no additional veterinary interventions were  
523 required of animals who had taken daily treatment  
524 with this drug for approximately 14 months.

525 Our results from this pilot study in large part  
526 confirmed that we could deliver rapamycin to mar-  
527 mosets in such a way to at least achieve similar  
528 blood concentrations (and *in vivo* mTOR inhibition)  
529 similar to that reported in mouse longevity studies  
530 using this drug. While this suggests the basic bio-  
531 chemical properties of this drug intervention can be  
532 translated between these species, other results from  
533 our pilot study support the notion that physiologi-  
534 cal outcomes of such an intervention may differ  
535 between species. For example, one of the relatively  
536 consistent ancillary effects of rapamycin treatment in  
537 mice has been development of glucose intolerance,  
538 likely due to an inhibitory effect of rapamycin on  
539 mTORC2 signaling in the liver [15, 100–102]. How-  
540 ever, in marmosets treated with rapamycin we found  
541 no evidence of glucose intolerance measured by oral  
542 glucose tolerance tests or indices of insulin resis-  
543 tance [103]. It is unclear exactly why we found these  
544 discrepancies between species, though one possibil-  
545 ity might be basic anatomical differences between  
546 mice and NHP. For example, the liver, which has  
547 been reported to be at least partially responsible for  
548 rapamycin-mediated glucose intolerance, makes up  
549 a greater contribution of the total mass of mice in

550 comparison to marmosets. In this regard, marmoset  
551 relative liver mass is much more similar to humans  
552 than to mice. While this provides no assurance that  
553 results from marmosets will directly translate to  
554 humans, it does hint at the potential value of utilizing  
555 a model species with basic biological characteristics  
556 similar to humans in biomedical research.

557 In a similar vein, there has been speculation that  
558 at least part of the beneficial effects of rapamycin  
559 on longevity are mediated through mTOR's effects  
560 on protein homeostasis (or proteostasis). While the  
561 relationships between mTOR and autophagy and  
562 mTOR and protein translation are well-understood,  
563 there is also growing evidence that mTOR may reg-  
564 ulate proteostasis through other pathways such as  
565 ubiquitin-proteasome and protein chaperone acti-  
566 vation. Indeed, previous reports have suggested  
567 that chronic rapamycin treatment can stimulate  
568 autophagy, induce proteasome activity and increase  
569 expression of protein chaperones [104–107]. In mar-  
570 mosets treated with rapamycin, we found evidence  
571 for mild stimulation of autophagy in some, though  
572 not all, tissues compared to control animals though  
573 little to no evidence that other protein degradation  
574 pathways, including proteasome and protein chaper-  
575 ones, were affected [108]. These data then suggest  
576 that if rapamycin has an effect on healthy aging in  
577 the marmoset, at least some of this effect might be  
578 contributed to activation of autophagy and provides  
579 a mechanistic target of action for further study.

580 The goal of our ongoing marmoset study is to  
581 directly address the question of whether interven-  
582 tion with rapamycin will benefit longevity or healthy  
583 aging in this species. As mentioned above, we have  
584 recruited and enrolled a cohort of middle-aged mar-  
585 mosets into a long-term study to test this question. We  
586 have designed this experiment starting with older ani-  
587 mals (approximately 5–7 years of age in this species)  
588 for three main reasons. 1) This provides a reasonable  
589 chance to see effects on longevity in this species over  
590 the next 5–10 years. 2) This is similar to the original  
591 Interventions Testing Program report on rapamycin  
592 which started in middle-aged mice and showed a ben-  
593 efiticial effect of this drug on longevity [13]. 3) It is  
594 reasonable to assume that translation of aging inter-  
595 ventions in humans might target those individuals that  
596 have reached a period considered later in life [97].  
597 Our cohort is of mixed sex with roughly equivalent  
598 numbers of male and female animals. We are treat-  
599 ing half of the cohort with rapamycin administered  
600 daily at dose of 1 mg/kg body weight which we have  
601 shown is comparable to doses known to extend mouse

602 lifespan [99], the other half of this cohort is control  
603 (*i.e.*, treated daily with the agent used to encapsulate  
604 rapamycin only). We are tracking physiological and  
605 behavioral phenotypes as well as health outcomes and  
606 longevity over the next several years.

607 While our primary goal is to assess effects of  
608 this intervention on longevity in the marmoset, we  
609 also have the opportunity to examine the effects of  
610 rapamycin on “healthy aging” or the progression of  
611 age-related physiological decline. In mice, rapamycin  
612 has been suggested to delay this progression (or  
613 even reverse age-related loss) across several organ  
614 systems with cardiac [91, 93], hematopoietic stem  
615 cells [96], the immune system [94], and periodontal  
616 bone [109] benefits among the many reported positive  
617 outcomes. In our cohort of aging marmosets,  
618 we are testing the effect of both advancing age and  
619 rapamycin intervention across multiple minimally-  
620 invasive assessments of animal health, frailty and  
621 resilience. For example, changes in blood chemistry,  
622 complete blood counts, blood lipids, etc. are being  
623 tested through regular blood draws similar to what  
624 would be performed during routine physical assess-  
625 ments in a clinical setting. In addition, we have  
626 defined scheduled tests of glucose metabolism, car-  
627 diovascular function, musculoskeletal function and  
628 inflammatory processes outlined for repeated test-  
629 ing throughout the remaining lifetime of this cohort.  
630 These repeated longitudinal tests also give us an  
631 opportunity to address long-term administration of  
632 rapamycin to a relatively healthy population has sim-  
633 ilar potential for side-effects as in clinical use to treat  
634 specific disease conditions. For example, the most  
635 commonly cited potential side effects of this drug are  
636 metabolic dysfunction (including new-onset type 2  
637 diabetes and dyslipidemia) and immunological sup-  
638 pression. As mentioned above, we have found so  
639 far no evidence to suggest rapamycin is impairing  
640 glucose metabolism at least through approximately  
641 1 year of treatment and no evidence for dyslipi-  
642 demia [51]. We continue to monitor these outcomes  
643 in our long-term treatment cohort. While our design  
644 does not allow us to directly test changes in immune  
645 response with age and rapamycin in the marmoset,  
646 we do monitor changes in blood cell count (includ-  
647 ing subsets of leukocytes) during our semi-annual  
648 health checkups. Again, to date we have found little  
649 evidence that chronic rapamycin significantly alters  
650 these parameters. Functional tests of immune cell  
651 function *in vivo* would at least tangentially address  
652 whether rapamycin inherently alters the ability of the  
653 immune cells to address challenge.

654 At death, animals will undergo full pathological  
655 assessment as well as to tissue collection to assess  
656 the effect of chronic intervention with rapamycin  
657 on multiple markers that represent the “pillars of  
658 aging” [110]. Thus, over the course of this study we  
659 should develop a complete picture of not only the  
660 broad effects of aging on the common marmoset but  
661 also whether pharmaceutical interventions known to  
662 extend lifespan in mice have similar effects in a NHP  
663 species.

## 664 6. Conclusions

665 Marmosets offer a unique non-human primate  
666 model in which to evaluate both cross-sectional and  
667 longitudinal effects of aging on measures of longevity  
668 and healthspan. While areas of interest remain for  
669 which there are currently no assessment tools for use  
670 in the marmoset, recent developments have rapidly  
671 expanded tools that are available. Marmosets have  
672 significant advantages including relatively short life  
673 span and ease of handling that make them ideal for  
674 this type of work. We look forward to the continued  
675 expansion of tools and knowledge that are becom-  
676 ing available for the marmosets as models for aging  
677 research.

## 678 Acknowledgments

679 During the preparation of this manuscript, CNR  
680 received support from a grant from the National  
681 Institute of Health (P30 AG013319); ABS received  
682 support from grants from the National Institute of  
683 Health (R01 AG050797 and R01 AG057431), the San  
684 Antonio Nathan Shock Center and Claude A. Pepper  
685 Center, and from the Geriatric Research, Education  
686 and Clinical Center of the South Texas Veterans  
687 Health Care System. This material is the result of  
688 work supported with resources and the use of facil-  
689 ities at South Texas Veterans Health Care System,  
690 San Antonio, Texas. The contents do not represent  
691 the views of the U.S. Department of Veterans Affairs  
692 or the United States Government.

## 693 References

- 694 [1] Wang AM, Promislow DE, Kaeberlein M. Fertile  
695 waters for aging research. *Cell*. 2015;160(5):814-5. doi:  
696 10.1016/j.cell.2015.02.026. PubMed PMID: 25723160.



- 697 [2] Bitto A, Wang AM, Bennett CF, Kaeberlein M. Biochem-  
698 ical genetic pathways that modulate aging in multiple  
699 species. *Cold Spring Harbor Perspectives in Medicine*.  
700 2015;5(11). doi: 10.1101/cshperspect.a025114  
701 [3] Santulli G, Borrás C, Bousquet J, Calzá L, Cano A, Illario  
702 M, et al. Models for preclinical studies in aging-related dis-  
703 orders: One is not for all. *Translational Medicine @ UniSa*.  
704 2015;13:4-12. PubMed PMID: PMC4811343.  
705 [4] Yuan R, Peters LL, Paigen B. Mice as a mammalian model  
706 for research on the genetics of aging. *ILAR J*. 2011;52(1):4-  
707 15. PubMed PMID: 21411853; PubMed Central PMCID:  
708 PMCPMC3074346.  
709 [5] Koopman JJ, van Heemst D, van Bodegom D, Bonkowski  
710 MS, Sun LY, Bartke A. Measuring aging rates of mice sub-  
711 jected to caloric restriction and genetic disruption of growth  
712 hormone signaling. *Aging (Albany NY)*. 2016;8(3):539-46.  
713 doi: 10.18632/aging.100919. PubMed PMID: 26959761;  
714 PubMed Central PMCID: PMCPMC4833144.  
715 [6] Morselli E, Maiuri MC, Markaki M, Megalou E, Pasparaki  
716 A, Palikaras K, et al. Caloric restriction and resveratrol  
717 promote longevity through the Sirtuin-1-dependent  
718 induction of autophagy. *Cell Death Dis*. 2010;1:e10. doi:  
719 10.1038/cddis.2009.8. PubMed PMID: 21364612; PubMed  
720 Central PMCID: PMCPMC3032517.  
721 [7] Wuttke D, Connor R, Vora C, Craig T, Li Y, Wood S, et al.  
722 Dissecting the gene network of dietary restriction to iden-  
723 tify evolutionarily conserved pathways and new functional  
724 genes. *PLoS Genet*. 2012;8(8):e1002834. doi: 10.1371/  
725 journal.pgen.1002834. PubMed PMID: 22912585;  
726 PubMed Central PMCID: PMCPMC3415404.  
727 [8] McDonald RB, Ramsey JJ. Honoring clive McCay and 75  
728 years of calorie restriction research. *The Journal of Nutri-  
729 tion*. 2010;140(7):1205-10. doi: 10.3945/jn.110.122804.  
730 PubMed PMID: PMC2884327.  
731 [9] Yu BP, Masoro EJ, Murata I, Bertrand HA, Lynd FT. Life  
732 span study of SPF Fischer 344 male rats fed ad libitum or  
733 restricted diets: Longevity, growth, lean body mass and dis-  
734 ease. *Journal of gerontology*. 1982;37(2):130-41. PubMed  
735 PMID: 7056998.  
736 [10] Hasty P, Livi CB, Dodds SG, Jones D, Strong R, Javors M,  
737 et al. eRapa restores a normal life span in a FAP mouse  
738 model. *Cancer Prevention Research*. 2014;7(1):169-78.  
739 doi: 10.1158/1940-6207.CAPR-13-0299. PubMed PMID:  
740 24282255; PubMed Central PMCID: PMC4058993.  
741 [11] Kennedy BK, Pennypacker JK. Drugs that modulate  
742 aging: The promising yet difficult path ahead. *Transl  
743 Res*. 2014;163(5):456-65. doi: 10.1016/j.trsl.2013.11.007.  
744 PubMed PMID: 24316383; PubMed Central PMCID:  
745 PMCPMC4004650.  
746 [12] Sharp ZD, Bartke A. Evidence for down-regulation of phos-  
747 phoinositide 3-kinase/Akt/mammalian target of rapamycin  
748 (PI3K/Akt/mTOR)-dependent translation regulatory sig-  
749 naling pathways in Ames dwarf mice. *J Gerontol A Biol Sci  
750 Med Sci*. 2005;60(3):293-300. PubMed PMID: 15860463.  
751 [13] Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle  
752 CM, Flurkey K, et al. Rapamycin fed late in life  
753 extends lifespan in genetically heterogeneous mice.  
754 *Nature*. 2009;460(7253):392-5. doi: 10.1038/nature08221.  
755 PubMed PMID: 19587680; PubMed Central PMCID:  
756 PMC2786175.  
757 [14] Miller RA, Harrison DE, Astle CM, Baur JA, Boyd AR,  
758 de Cabo R, et al. Rapamycin, but not resveratrol or sim-  
759 vastatin, extends life span of genetically heterogeneous  
760 mice. *The Journals of Gerontology Series A: Biologi-  
761 cal Sciences and Medical Sciences*. 2011;66A(2):191-201.  
762 doi: 10.1093/gerona/gdq178. PubMed Central PMCID:  
763 PMC3021372.  
764 [15] Miller RA, Harrison DE, Astle CM, Fernandez E, Flurkey  
765 K, Han M, et al. Rapamycin-mediated lifespan increase in  
766 mice is dose and sex dependent and metabolically distinct  
767 from dietary restriction. *Aging Cell*. 2014;13(3):468-  
768 77. doi: 10.1111/accel.12194. PubMed Central PMCID:  
769 PMC4032600.  
770 [16] Harrison DE, Strong R, Allison DB, Ames BN, Astle  
771 CM, Atamna H, et al. Acarbose, 17-alpha-estradiol, and  
772 nordihydroguaiaretic acid extend mouse lifespan prefer-  
773 entially in males. *Aging Cell*. 2014;13(2):273-82. doi:  
774 10.1111/accel.12170. PubMed PMID: 24245565; PubMed  
775 Central PMCID: PMC3954939.  
776 [17] Strong R, Miller RA, Antebi A, Astle CM, Bogue  
777 M, Denzel MS, et al. Longer lifespan in male mice  
778 treated with a weakly estrogenic agonist, an antioxi-  
779 dant, an alpha-glucosidase inhibitor or a Nrf2-inducer.  
780 *Aging Cell*. 2016;15(5):872-84. doi: 10.1111/accel.12496.  
781 PubMed PMID: 27312235.  
782 [18] Bracken MB. Why animal studies are often poor predic-  
783 tors of human reactions to exposure. *Journal of the Royal  
784 Society of Medicine*. 2009;102(3):120-2. doi: 10.1258/  
785 jrsm.2008.08k033. PubMed PMID: PMC2746847.  
786 [19] Zeiss CJ, Allore HG, Beck AP. Established patterns of  
787 animal study design undermine translation of disease-  
788 modifying therapies for Parkinson's disease. *PLoS One*.  
789 2017;12(2):e0171790. doi: 10.1371/journal.pone.0171790.  
790 PubMed PMID: 28182759.  
791 [20] Chatterjee HJ, Ho SYW, Barnes I, Groves C. Estim-  
792 ating the phylogeny and divergence times of primates  
793 using a supermatrix approach. *BMC Evolutionary Biology*.  
794 2009;9(1):259. doi: 10.1186/1471-2148-9-259  
795 [21] Finch C, Austad SN. Primate aging in the mammalian  
796 scheme: The puzzle of extreme variation in brain aging.  
797 *AGE*. 2012;34:1075-91.  
798 [22] Bellino FL, Wise PM. Nonhuman primate models of  
799 menopause workshop. *Biol Reprod*. 2003;68(1):10-8.  
800 PubMed PMID: 12493689.  
801 [23] Chilton J, Wilcox A, Lammey M, Meyer D. Char-  
802 acterization of a cardiorenal-like syndrome in aged chim-  
803 panzees (*Pan troglodytes*). *Vet Pathol*. 2016;53(2):417-  
804 24. doi: 10.1177/0300985815618435. PubMed PMID:  
805 26792841.  
806 [24] Ely JJ, Zavaskis T, Lammey ML. Hypertension increases  
807 with aging and obesity in chimpanzees (*Pan troglodytes*).  
808 *Zoo Biol*. 2013;32(1):79-87. doi: 10.1002/zoo.21044.  
809 PubMed PMID: 22968757; PubMed Central PMCID:  
810 PMCPMC3537917.  
811 [25] Gilissen EP, Leroy K, Yilmaz Z, Kovari E, Bouras C,  
812 Boom A, et al. A neuronal aging pattern unique to  
813 humans and common chimpanzees. *Brain Struct Funct*.  
814 2016;221(1):647-64. doi: 10.1007/s00429-014-0931-5.  
815 PubMed PMID: 25381006.  
816 [26] Lowenstine LJ, McManamon R, Terio KA. Comparative  
817 pathology of aging great apes: Bonobos, chimpanzees,  
818 gorillas, and orangutans. *Vet Pathol*. 2016;53(2):250-  
819 76. doi: 10.1177/0300985815612154. PubMed PMID:  
820 26721908.

- 821 [27] Nunamaker EA, Lee DR, Lammey ML. Chronic diseases  
822 in captive geriatric female Chimpanzees (Pan troglodytes).  
823 *Comp Med.* 2012;62(2):131-6. PubMed PMID: 22546920;  
824 PubMed Central PMCID: PMCPMC3318251.
- 825 [28] Sleeper MM, Drobatz K, Lee DR, Lammey ML. Echocar-  
826 diography parameters of clinically normal adult captive  
827 chimpanzees (Pan troglodytes). *J Am Vet Med Assoc.*  
828 2014;244(8):956-60. doi: 10.2460/javma.244.8.956.  
829 PubMed PMID: 24697773.
- 830 [29] Videan EN, Fritz J, Murphy J. Effects of aging on  
831 hematology and serum clinical chemistry in chimpanzees  
832 (Pan troglodytes). *Am J Primatol.* 2008;70(4):327-38. doi:  
833 10.1002/ajp.20494. PubMed PMID: 17943981.
- 834 [30] Didier ES, MacLean AG, Mohan M, Didier PJ, Lack-  
835 ner AA, Kuroda MJ. Contributions of nonhuman primates  
836 to research on aging. *Vet Pathol.* 2016;53(2):277-90. doi:  
837 10.1177/0300985815622974. PubMed PMID: 26869153;  
838 PubMed Central PMCID: PMCPMC5027759.
- 839 [31] Finch C, Austad S. Primate models for human brain aging  
840 and neurological diseases. *Annual Review of Gerontology*  
841 *and Geriatrics.* 2014;34(1):139-70. doi: 10.1891/0198-  
842 8794.34.139
- 843 [32] Lacreuse A, Herndon JG. Nonhuman primate models of  
844 cognitive aging. 2009:1-30. doi: 10.1007/978-1-59745-  
845 422-3\_2
- 846 [33] Lane MA. Nonhuman primate models in biogerontol-  
847 ogy. *Exp Gerontol.* 2000;35(5):533-41. PubMed PMID:  
848 10978676.
- 849 [34] Mattison JA, Vaughan KL. An overview of nonhuman  
850 primates in aging research. *Exp Gerontol.* 2016. doi:  
851 10.1016/j.exger.2016.12.005. PubMed PMID: 27956088.
- 852 [35] Verdier JM, Acquatella I, Lautier C, Devau G, Trouche S,  
853 Lasbleiz C, et al. Lessons from the analysis of nonhuman  
854 primates for understanding human aging and neurode-  
855 generative diseases. *Front Neurosci.* 2015;9:64. doi:  
856 10.3389/fnins.2015.00064. PubMed PMID: 25788873;  
857 PubMed Central PMCID: PMCPMC4349082.
- 858 [36] Hoffman CL, Higham JP, Mas-Rivera A, Ayala JE,  
859 Maestripieri D. Terminal investment and senescence in rhesus  
860 macaques (*Macaca mulatta*) on Cayo Santiago. *Behav*  
861 *Ecol.* 2010;21(5):972-8. doi: 10.1093/beheco/arq098.  
862 PubMed PMID: 22475990; PubMed Central PMCID:  
863 PMCPMC2920293.
- 864 [37] Maestripieri D, Hoffman CL. Chronic stress, allostatic  
865 load, and aging in nonhuman primates. *Dev Psychopathol.*  
866 2011;23(4):1187-95. doi: 10.1017/S0954579411000551.  
867 PubMed PMID: 22018089; PubMed Central PMCID:  
868 PMCPMC3942660.
- 869 [38] Colman RJ, Anderson RM. Nonhuman primate caloric  
870 restriction. *Antioxid Redox Signal.* 2011;14(2):229-39.  
871 doi: 10.1089/ars.2010.3224. PubMed PMID: 20698791;  
872 PubMed Central PMCID: PMCPMC3000242.
- 873 [39] Colman RJ, Beasley TM, Allison DB, Weindruch R.  
874 Attenuation of sarcopenia by dietary restriction in rhesus  
875 monkeys. *J Gerontol A Biol Sci Med Sci.* 2008;63(6):556-  
876 9. PubMed PMID: 18559628; PubMed Central PMCID:  
877 PMCPMC2812805.
- 878 [40] Colman RJ, Beasley TM, Allison DB, Weindruch R.  
879 Skeletal effects of long-term caloric restriction in rhesus  
880 monkeys. *Age (Dordr).* 2012;34(5):1133-43. doi:  
881 10.1007/s11357-011-9354-x. PubMed PMID: 22189911;  
882 PubMed Central PMCID: PMCPMC3448987.
- [41] Colman RJ, Beasley TM, Kemnitz JW, Johnson SC, 883  
Weindruch R, Anderson RM. Caloric restriction reduces 884  
age-related and all-cause mortality in rhesus monkeys. 885  
*Nat Commun.* 2014;5:3557. doi: 10.1038/ncomms4557. 886  
PubMed PMID: 24691430; PubMed Central PMCID: 887  
PMCPMC3988801. 888
- [42] Kemnitz JW. Calorie restriction and aging in nonhuman 889  
primates. *Institute for Laboratory Animal Research Journal.* 890  
2011;52(1):66-77. 891
- [43] Mattison JA, Colman RJ, Beasley TM, Allison DB, Kem- 892  
nitz JW, Roth GS, et al. Caloric restriction improves 893  
health and survival of rhesus monkeys. *Nat Commu-* 894  
*n.* 2017;8:14063. doi: 10.1038/ncomms14063. PubMed 895  
PMID: 28094793. 896
- [44] Mattison JA, Roth GS, Beasley TM, Tilmont EM, 897  
Handy AM, Herbert RL, et al. Impact of caloric restric- 898  
tion on health and survival in rhesus monkeys from the 899  
NIA study. *Nature.* 2012;489(7415):318-21. doi: 900  
10.1038/nature11432. PubMed PMID: 22932268; PubMed 901  
Central PMCID: PMCPMC3832985. 902
- [45] Smucny DA, Allison DB, Ingram DK, Roth GS, Kem- 903  
nitz JW, Kohama SG, et al. Changes in blood chemistry 904  
and hematology variables during aging in captive rhesus 905  
macaques (*Macaca mulatta*). *Journal of Medical Primatol-* 906  
*ogy.* 2001;30:161-73. 907
- [46] Austad SN. Small nonhuman primates as potential models 908  
of human aging. *Institute for Laboratory Animal Research* 909  
*Journal.* 1997;38(3):142-7. 910
- [47] Fischer KE, Austad SN. The development of small primate 911  
models for aging research. *Institute for Laboratory Animal* 912  
*Research Journal.* 2011;52(1):78-88. 913
- [48] Pifferi F, Terrien J, Marchal J, Dal-Pan A, Djelti F, Hardy I, 914  
et al. Caloric restriction increases lifespan but affects brain 915  
integrity in grey mouse lemur primates. *Communications* 916  
*Biology.* 2018;1(1):30. doi: 10.1038/s42003-018-0024-8 917
- [49] Mansfield KG. Marmoset models commonly used 918  
in biomedical research. *Comparative Medicine.* 919  
2003;53(4):383-92. 920
- [50] Ross CN, Davis K, Dobek G, Tardif SD. Aging pheno- 921  
types of common marmosets (*Callithrix jacchus*). *Journal* 922  
*of Aging Research.* 2012;2012:567143. 923
- [51] Ross CN, Salmon AB, Strong R, Fernandez E, Javors M, 924  
Richardson A, et al. Metabolic consequences of long-term 925  
rapamycin exposure on common marmoset monkeys (*Cal-* 926  
*lithrix jacchus*). *Aging.* 2015;7(11):1-10. 927
- [52] Salmon AB. Moving toward 'common' use of the marmoset 928  
as a non-human primate aging model. *Pathobiol Aging* 929  
*Age Relat Dis.* 2016;6:32758. doi: 10.3402/pba.v6.32758. 930  
PubMed PMID: 27452489; PubMed Central PMCID: 931  
PMCPMC4958916. 932
- [53] Tardif SD, Abee CR, Mansfield KG. Workshop summary: 933  
Neotropical primates in biomedical research. *Institute for* 934  
*Laboratory Animal Research Journal.* 2011;52(3):386-92. 935
- [54] Tardif SD, Mansfield KG, Ratnam R, Ross CN, Ziegler 936  
TE. The marmoset as a model of aging and age-related 937  
disease. *Institute for Laboratory Animal Research Journal.* 938  
2011;52(1):54-65. 939
- [55] Tardif SD, Arauajuo A, Arruda MF, French JA, Sousa 940  
MB, Yamamoto K. Reproduction and aging in marmosets 941  
and tamarins. In: Atsalis S, Margulis SW, editors. *Primate* 942  
*Reproductive Aging: Interdisciplinary topics in gerontol-* 943  
*ogy and geriatrics.* Karger; 2008, pp. 29-48. 944

- 945 [56] Ross CN, Austad S, Brasky K, Brown CJ, Forney LJ, Gel- 1007  
 946 fond JA, et al. The development of a specific pathogen free 1008  
 947 (SPF) barrier colony of marmosets (*Callithrix jacchus*) for 1009  
 948 aging research. *Aging* (Albany NY). 2017;9(12):2544-58. 1010  
 949 doi: 10.18632/aging.101340. PubMed PMID: 29227963; 1011  
 950 PubMed Central PMCID: PMC5764392. 1012
- 951 [57] Bagi CM, Volberg M, Moalli M, Shen V, Olson E, Hanson 1013  
 952 N, et al. Age-related changes in marmoset trabecular and 1014  
 953 cortical bone and response to alendronate therapy resemble 1015  
 954 human bone physiology and architecture. *Anat Rec* 1016  
 955 (Hoboken). 2007;290(8):1005-16. doi: 10.1002/ar.20561. 1017  
 956 PubMed PMID: 17610276. 1018
- 957 [58] Bernick S, Caillet R, Levy BM. The maturation and aging 1019  
 958 of the vertebrae of marmosets. *Spine*. 1980;5(6):519-24. 1020
- 959 [59] Power RA, Power ML, Layne DG, Jaquish CE, Oftedal OT, 1021  
 960 Tardif SD. Relations among measures of body composition, 1022  
 961 age, and sex in the common marmoset monkey (*Callithrix* 1023  
 962 *jacchus*). *Comparative Medicine*. 2001;51(3):281-23. 1024
- 963 [60] Frisard MI, Broussard A, Davies SS, Roberts LJ, Rood J, de 1025  
 964 Jonge L, et al. Aging, resting metabolic rate, and oxidative 1026  
 965 damage: Results from the Louisiana healthy aging study. 1027  
 966 *The Journals of Gerontology Series A, Biological Sciences* 1028  
 967 *and Medical Sciences*. 2007;62(7):752-9. PubMed PMID: 1029  
 968 PMC2692620. 1030
- 969 [61] Leuner B, Kozorovitskiy Y, Gross CG, Gould E. Dimin- 1031  
 970 ished adult neurogenesis in the marmoset brain precedes 1032  
 971 old age. *Proc Natl Acad Sci U S A*. 2007;104(43):17169- 1033  
 972 73. doi: 10.1073/pnas.0708228104. PubMed PMID: 1034  
 973 17940008; PubMed Central PMCID: PMC2040400. 1035
- 974 [62] Honavar M, Lantos PL. Ultrastructural changes in the 1036  
 975 frontal cortex and hippocampus in the aging marmoset. 1037  
 976 *Mechanisms of Aging and Development*. 1987;41:161-75. 1038
- 977 [63] Kobayashi R, Takahashi-Fujigasaki J, Shiozawa S, Hara- 1039  
 978 miyauchi C, Inoue T, Okano HJ, et al. Alpha-Synuclein 1040  
 979 aggregation in the olfactory bulb of middle-aged com- 1041  
 980 mon marmoset. *Neurosci Res*. 2016;106:55-61. doi: 1042  
 981 10.1016/j.neures.2015.11.006. PubMed PMID: 26643383. 1043
- 982 [64] Geula C, Nagykerly N, Wu C-K. Amyloid-B deposits in 1044  
 983 the cerebral cortex of the aged common marmoset (*Callithrix* 1045  
 984 *jacchus*): Incidence and chemical composition. *Acta* 1046  
 985 *Neuropathol*. 2002;103:48-58. 1047
- 986 [65] Ridley RM, Baker HF, Windle CP, Cummings RM. Very 1048  
 987 long term studies of the seeding of beta-amyloidosis in pri- 1049  
 988 mates. *Journal of Neural Transmission*. 2006;113:1243-51. 1050
- 989 [66] Rodriguez-Callejas JD, Fuchs E, Perez-Cruz C. Evidence 1051  
 990 of Tau hyperphosphorylation and dystrophic microglia in 1052  
 991 the common marmoset. *Frontiers in Aging Neuroscience*. 1053  
 992 2016;8:315. 1054
- 993 [67] Klonoff-Cohen H, Barrett-Connor EL, Edelstein SL. Albu- 1055  
 994 min levels as a predictor of mortality in the healthy elderly. 1056  
 995 *J Clin Epidemiol*. 1992;45(3):207-12. PubMed PMID: 1057  
 996 1569417. 1058
- 997 [68] Go YM, Liang Y, Uppal K, Soltow QA, Promislow 1059  
 998 DE, Wachtman LM, et al. Metabolic characterization 1060  
 999 of the common marmoset (*Callithrix jacchus*). *PLoS* 1061  
 1000 *One*. 2015;10(11):e0142916. doi: 10.1371/journal.pone. 1062  
 1001 0142916. PubMed PMID: 26581102; PubMed Central 1063  
 1002 PMCID: PMC4651467. 1064
- 1003 [69] Hoffman JM, Tran V, Wachtman LM, Green CL, Jones DP, 1065  
 1004 Promislow DE. A longitudinal analysis of the effects of 1066  
 1005 age on the blood plasma metabolome in the common mar- 1067  
 1006 moset, *Callithrix jacchus*. *Exp Gerontol*. 2016;76:17-24. 1068
- doi: 10.1016/j.exger.2016.01.007. PubMed PMID: 26805607; PubMed Central PMCID: PMC4775367.
- [70] Soltow QA, Jones DP, Promislow DE. A network 1009  
 perspective on metabolism and aging. *Integrative and* 1010  
*Comparative Biology*. 2010;50(5):844-54. 1011
- [71] Mietsch M, Baldauf K, Reitemeier S, Suchowski M, Schoon 1012  
 HA, Einspanier A. Blood pressure as prognostic marker for 1013  
 body condition, cardiovascular, and metabolic diseases in 1014  
 the common marmoset (*Callithrix jacchus*). *J Med Primatol*. 1015  
 2016;45(3):126-38. doi: 10.1111/jmp.12215. PubMed 1016  
 PMID: 27144322. 1017
- [72] Kelnar CJ, McKinnell C, Walker M, Morris KD, Wallace 1018  
 WH, Saunders PT, et al. Testicular changes during infantile 1019  
 'quiescence' in the marmoset and their gonadotrophin 1020  
 dependence: A model for investigating susceptibility of the 1021  
 prepubertal human testis to cancer therapy? *Hum Reprod*. 1022  
 2002;17(5):1367-78. PubMed PMID: 11980767. 1023
- [73] Kholkute SD, Aitken RJ, Lunn SF. Plasma testosterone 1024  
 response to hCG stimulation in the male marmoset 1025  
 monkey (*Callithrix jacchus jacchus*). *J Reprod Fertil*. 1026  
 1983;67(2):457-63. PubMed PMID: 6403699. 1027
- [74] Nunes S, Fite JE, French JA. Variation in steroid 1028  
 hormones associated with infant care behaviour and 1029  
 experience in male marmosets (*Callithrix kuhlii*). *Anim* 1030  
*Behav*. 2000;60(6):857-65. doi: 10.1006/anbe.2000.1524. 1031  
 PubMed PMID: 11124885. 1032
- [75] Kenyon CJ. The genetics of ageing. *Nature*. 2010;464:504. 1033  
 doi: 10.1038/nature08980 1034
- [76] Weindruch R, Walford RL, Fligiel S, Guthrie D. The retar- 1035  
 dation of aging in mice by dietary restriction: Longevity, 1036  
 cancer, immunity and lifetime energy intake. *The Journal of* 1037  
*Nutrition*. 1986;116(4):641-54. doi: 10.1093/jn/116.4.641 1038
- [77] Yang SH, Cheng PH, Banta H, Piotrowska-Nitsche 1039  
 K, Yang JJ, Cheng EC, et al. Towards a transgenic 1040  
 model of Huntington's disease in a non-human primate. 1041  
*Nature*. 2008;453(7197):921-4. doi: 10.1038/nature06975. 1042  
 PubMed PMID: 18488016; PubMed Central PMCID: 1043  
 PMC2652570. 1044
- [78] Seita Y, Tsukiyama T, Iwatani C, Tsuchiya H, Matsushita 1045  
 J, Azami T, et al. Generation of transgenic cynomolgus 1046  
 monkeys that express green fluorescent protein throughout 1047  
 the whole body. *Scientific Reports*. 2016;6:24868. doi: 1048  
 10.1038/srep24868https://www.nature.com/articles/srep 1049  
 24868#supplementary-information. 1050
- [79] Sasaki E, Suemizu H, Shimada A, Hanazawa K, 1051  
 Oiwa R, Kamioka M, et al. Generation of trans- 1052  
 genic non-human primates with germline transmission. 1053  
*Nature*. 2009;459(7246):523-7. doi: 10.1038/nature08090. 1054  
 PubMed PMID: 19478777. 1055
- [80] Sasaki E. Prospects for genetically modified non-human 1056  
 primate models, including the common marmoset. *Neurosci* 1057  
*Res*. 2015;93:110-5. doi: 10.1016/j.neures.2015. 1058  
 01.011. PubMed PMID: 25683291. 1059
- [81] Tomioka I, Ishibashi H, Minakawa EN, Motohashi 1060  
 HH, Takayama O, Saito Y, et al. Transgenic mon- 1061  
 key model of the polyglutamine diseases recapitulating 1062  
 progressive neurological symptoms. *eNeuro*. 2017;4(2). 1063  
 doi: 10.1523/ENEURO.0250-16.2017. PubMed PMID: 1064  
 28374014; PubMed Central PMCID: PMC5368386. 1065
- [82] Tomioka I, Nogami N, Nakatani T, Owari K, Fujita 1066  
 N, Motohashi H, et al. Generation of transgenic mar- 1067  
 mosets using a tetracyclin-inducible transgene expression 1068

- system as a neurodegenerative disease model. *Biol Reprod.* 2017;97(5):772-80. doi: 10.1093/biolre/iox129. PubMed PMID: 29045563.
- [83] Park JE, Zhang XF, Choi SH, Okahara J, Sasaki E, Silva AC. Generation of transgenic marmosets expressing genetically encoded calcium indicators. *Sci Rep.* 2016;6:34931. doi: 10.1038/srep34931. PubMed PMID: 27725685; PubMed Central PMCID: PMC4693453.
- [84] Tardif S, Power M, Layne D, Smucny D, Ziegler T. Energy restriction initiated at different gestational ages has varying effects on maternal weight gain and pregnancy outcome in common marmoset monkeys (*Callithrix jacchus*). *The British journal of nutrition.* 2004;92(5):841-9. PubMed PMID: 15533274.
- [85] Tardif SD, Power ML, Ross CN, Rutherford JN, Layne-Colon DG, Paulik MA. Characterization of obese phenotypes in a small nonhuman primate, the common marmoset (*Callithrix jacchus*). *Obesity (Silver Spring).* 2009;17(8):1499-505. doi: 10.1038/oby.2009.77. PubMed PMID: 19325546; PubMed Central PMCID: PMC4693453.
- [86] Miller CT, Freiwald WA, Leopold DA, Mitchell JF, Silva AC, Wang X. Marmosets: A neuroscientific model of human social behavior. *Neuron.* 2016;90(2):219-33. doi: 10.1016/j.neuron.2016.03.018. PubMed PMID: PMC4840471.
- [87] Soars MG, Riley RJ, Burchell B. Evaluation of the marmoset as a model species for drug glucuronidation. *Xenobiotica; the Fate of Foreign Compounds in Biological Systems.* 2001;31(12):849-60. doi: 10.1080/00498250110069690. PubMed PMID: 11780760.
- [88] VanderVeen N, Paran C, Appelhans A, Krasinkiewicz J, Lemons R, Appelman H, et al. Marmosets as a preclinical model for testing "off-label" use of doxycycline to turn on Flt3L expression from high-capacity adenovirus vectors. *Molecular Therapy — Methods & Clinical Development.* 2014;1. doi: 10.1038/mtm.2013.10 http://www.nature.com/articles/mtm201310#supplementary-information. PubMed Central PMCID: PMC4111110.
- [89] Orsi A, Rees D, Andreini I, Venturella S, Cinelli S, Oberto G. Overview of the marmoset as a model in nonclinical development of pharmaceutical products. *Regulatory Toxicology and Pharmacology.* 2011;59(1):19-27. doi: https://doi.org/10.1016/j.yrtph.2010.12.003
- [90] Wilkinson JE, Burmeister L, Brooks SV, Chan C-C, Friedline S, Harrison DE, et al. Rapamycin slows aging in mice. *Aging Cell.* 2012;11(4):675-82. doi: 10.1111/j.1474-9726.2012.00832.x
- [91] Flynn JM, O'Leary MN, Zambataro CA, Academia EC, Presley MP, Garrett BJ, et al. Late-life rapamycin treatment reverses age-related heart dysfunction. *Aging Cell.* 2013;12(5):851-62. doi: 10.1111/ace1.12109. PubMed Central PMCID: PMC4098908.
- [92] Zhang Y, Bokov A, Gelfond J, Soto V, Ikeno Y, Hubbard G, et al. Rapamycin extends life and health in C57BL/6 mice. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences.* 2014;69A(2):119-30. doi: 10.1093/gerona/glt056. PubMed Central PMCID: PMC4038246.
- [93] Dai D-F, Karunadharm PP, Chiao YA, Basisty N, Crispin D, Hsieh EJ, et al. Altered proteome turnover and remodeling by short-term caloric restriction or rapamycin rejuvenate the aging heart. *Aging Cell.* 2014;13(3):529-39. doi: 10.1111/ace1.12203. PubMed Central PMCID: PMC4040127
- [94] Hurez V, Dao V, Liu A, Pandeswara S, Gelfond J, Sun L, et al. Chronic mTOR inhibition in mice with rapamycin alters T, B, myeloid, and innate lymphoid cells and gut flora and prolongs life of immune-deficient mice. *Aging Cell.* 2015;14(6):945-56. doi: 10.1111/ace1.12380. PubMed PMID: 26315673; PubMed Central PMCID: PMC4693453.
- [95] Chiao YA, Kolwicz SC, Basisty N, Gagnidze A, Zhang J, Gu H, et al. Rapamycin transiently induces mitochondrial remodeling to reprogram energy metabolism in old hearts. *Aging (Albany NY).* 2016;8(2):314-27. PubMed PMID: 26872208; PubMed Central PMCID: PMC4789585.
- [96] Chen C, Liu Y, Liu Y, Zheng P. mTOR regulation and therapeutic rejuvenation of aging hematopoietic stem cells. *Sci Signal.* 2009;2(98):ra75. doi: 10.1126/scisignal.2000559
- [97] Mannick JB, Del Giudice G, Lattanzi M, Valiante NM, Praestgaard J, Huang B, et al. mTOR inhibition improves immune function in the elderly. *Science Translational Medicine.* 2014;6(268):268ra179. doi: 10.1126/scitranslmed.3009892
- [98] Kraig E, Linehan LA, Liang H, Romo TQ, Liu Q, Wu Y, et al. A randomized control trial to establish the feasibility and safety of rapamycin treatment in an older human cohort: Immunological, physical performance, and cognitive effects. *Exp Gerontol.* 2018;105:53-69. doi: 10.1016/j.exger.2017.12.026. PubMed PMID: 29408453; PubMed Central PMCID: PMC5869166.
- [99] Tardif S, Ross C, Bergman P, Fernandez E, Javors M, Salmon A, et al. Testing efficacy of administration of the antiaging drug rapamycin in a nonhuman primate, the common marmoset. *J Gerontol A Biol Sci Med Sci.* 2015;70(5):577-87. doi: 10.1093/gerona/glu101. PubMed PMID: 25038772; PubMed Central PMCID: PMC4400395.
- [100] Liu Y, Diaz V, Fernandez E, Strong R, Ye L, Baur JA, et al. Rapamycin-induced metabolic defects are reversible in both lean and obese mice. *Aging (Albany NY).* 2014;6(9):742-54. PubMed PMID: 25324470.
- [101] Lamming DW, Ye L, Katajisto P, Goncalves MD, Saitoh M, Stevens DM, et al. Rapamycin-induced insulin resistance is mediated by mTORC2 loss and uncoupled from longevity. *Science.* 2012;335(6076):1638-43. doi: 10.1126/science.1215135. PubMed Central PMCID: PMC3324089.
- [102] Weiss R, Fernandez E, Liu Y, Strong R, Salmon AB. Metformin reduces glucose intolerance caused by rapamycin treatment in genetically heterogeneous female mice. *Aging (Albany NY).* 2018. doi: 10.18632/aging.101401. PubMed PMID: 29579736.
- [103] Ross C, Salmon A, Strong R, Fernandez E, Javors M, Richardson A, et al. Metabolic consequences of long-term rapamycin exposure on common marmoset monkeys (*Callithrix jacchus*). *Aging (Albany NY).* 2015;7(11):964-73. PubMed PMID: 26568298; PubMed Central PMCID: PMC4694066.
- [104] Zhao J, Zhai B, Gygi SP, Goldberg AL. mTOR inhibition activates overall protein degradation by the ubiquitin proteasome system as well as by autophagy. *Proc Natl Acad Sci U S A.* 2015;112(52):15790-7. doi: 10.1073/

- 1193 pnas.1521919112. PubMed PMID: 26669439; PubMed  
1194 Central PMCID: PMC4703015.
- 1195 [105] Diaz-Troya S, Perez-Perez ME, Florencio FJ, Crespo JL.  
1196 The role of TOR in autophagy regulation from yeast  
1197 to plants and mammals. *Autophagy*. 2008;4(7):851-65.  
1198 PubMed PMID: 18670193.
- 1199 [106] Conn CS, Qian SB. mTOR signaling in protein home-  
1200 ostasis: Less is more? *Cell Cycle*. 2011;10(12):1940-7.  
1201 PubMed PMID: 21555915; PubMed Central PMCID:  
1202 PMC3154417.
- 1203 [107] Zhang Y, Nicholatos J, Dreier JR, Ricoult SJ, Widen-  
1204 maier SB, Hotamisligil GS, et al. Coordinated regulation  
1205 of protein synthesis and degradation by mTORC1.  
1206 *Nature*. 2014;513(7518):440-3. doi: 10.1038/nature13492.  
1207 PubMed PMID: 25043031; PubMed Central PMCID:  
1208 PMC4402229.
- [108] Lelegren M, Liu Y, Ross C, Tardif S, Salmon AB. 1209  
Pharmaceutical inhibition of mTOR in the common mar- 1210  
moset: Effect of rapamycin on regulators of proteostasis 1211  
in a non-human primate. *Pathobiol Aging Age Relat Dis*. 1212  
2016;6:31793. doi: 10.3402/pba.v6.31793. PubMed PMID: 1213  
27341957; PubMed Central PMCID: PMC4920937. 1214
- [109] An JY, Quarles EK, Mekvanich S, Kang A, Liu A, Santos 1215  
D, et al. Rapamycin treatment attenuates age-associated 1216  
periodontitis in mice. *GeroScience*. 2017;39(4):457- 1217  
63. doi: 10.1007/s11357-017-9994-6. PubMed PMID: 1218  
PMC5636779. 1219
- [110] López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer 1220  
G. The hallmarks of aging. *Cell*. 153(6):1194-217. doi: 1221  
10.1016/j.cell.2013.05.039 1222