

Review Article

Does objectively measured prolonged standing for desk work result in lower ratings of perceived low back pain than sitting? A systematic review and meta-analysis

Diana De Carvalho^{a,*}, Ryan Greene^a, Michelle Swab^b and Marshall Godwin^c

^a*Clinical Epidemiology, Discipline of Medicine, Faculty of Medicine, Memorial University of Newfoundland, St. John's, NL, Canada*

^b*Health Science Library, Faculty of Medicine, Memorial University of Newfoundland, St. John's, NL, Canada*

^c*Primary Healthcare Research Unit, Faculty of Medicine, Memorial University of Newfoundland, St. John's, NL, Canada*

Received 18 January 2019

Accepted 2 September 2019

Abstract.

BACKGROUND: Prolonged sitting has been shown to induce transient low back pain (LBP). Height adjustable office desks now present the opportunity to replace sitting with standing in the workplace. Since standing has also been associated with LBP, this may not be an advisable alternative.

OBJECTIVE: To determine if objectively measured prolonged exposures to desk work while standing, compared to sitting, results in lower perceived LBP in healthy adults.

METHODS: A systematic search of several databases was conducted. Two independent reviewers screened titles/abstracts and conducted a quality assessment. The results of three studies were pooled using an inverse variance random-effects meta-analysis. Heterogeneity was tested using the Chi-squared test and I^2 statistic.

RESULTS: Objectively measured prolonged standing postures during desk work did not induce significantly less perceived LBP compared to seated postures (standardized mean difference 0.60, 95% CI -0.68 to 1.87, $p = 0.36$.) There was significant heterogeneity, $I^2 = 90\%$.

CONCLUSIONS: It appears that replacing seated desk work postures with standing for prolonged periods of time would not be recommended. Larger studies, including a wider age range and health history, conducted in the field with objective measures is recommended to obtain more generalizable data on which to base ergonomic standards for work postures.

Keywords: Transient pain, office work, objectively measured posture, low back pain

*Address for correspondence: Diana De Carvalho, Clinical Epidemiology Discipline of Medicine, Faculty of Medicine, Memorial University of Newfoundland, 300 Prince Philip

Drive St. John's NL A1B 3V6, Canada. E-mail: Diana.decarvalho@med.mun.ca.

1. Introduction

Globally, low back pain (LBP) is the leading cause of years lost to disability [1]; is responsible for billions of health care dollars annually [2, 3]; and is not easily resolved, with a significant number of cases progressing to chronicity [4, 5]. *In vivo* basic science research has demonstrated that prolonged sitting can induce transient perceived back pain in young, healthy populations [6–10]. Considering 40% of the workforce in developed countries [11] are seated for more than two thirds of the workday [12–14] solutions for those who develop pain in sitting are important. To this end, the introduction of height adjustable office desks to workplaces has provided an option to replace seated postures with standing. However, prolonged standing has also been shown to be associated with LBP [15] and may not be a helpful alternative.

To date, there has been a number of epidemiological studies that have looked at the association of either standing or sitting postures with LBP [16–20]. However, there has been no systematic evaluation of the literature that directly compared seated and standing desk work postures in terms of immediate LBP development. Further, the studies that have examined the association of either sitting or standing postures with back pain have relied on self-report exposure time or have assumed posture duration based on an occupation title. There is evidence from the literature to suggest that these methods of time determination far underestimate actual durations [21]. The objective measure of sitting and standing time, either by direct observation, timed laboratory trial or wearable sensor would provide a more robust dose/response picture of exposure to postures with back pain. Therefore, the purpose of this study is to conduct a systematic review and meta-analysis to determine whether objectively measured prolonged standing results in less perceived low back pain compared to prolonged sitting.

1.1. Research question

Does objectively measured (by laboratory controlled time trial, direct observation or wearable sensor) exposures of prolonged (\geq one hour) desk work while standing result in lower perceived back pain (upper or lower back determined by pain scale rating) compared to sitting in adults (>18 years of age) with no history of low back pain?

2. Methods

2.1. Identification of studies

A systematic review of the literature was conducted, with the support of a health services librarian (MS), in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [22]. The electronic databases PubMed (MEDLINE), EMBASE, SPORTDiscus and CINAHL were searched on October 20, 2017 and updated on November 14, 2017 and September 2, 2018. No language restrictions were used and all articles published from database inception to the search date were included. Both observational studies (cross-sectional, cohort, case control) and randomized control trials were sought. The search strategy included all possible versions and combinations of the terms “back pain”, “discomfort”, “upper back”, “lower back”, “objective measure”, “sensor”, “laboratory”, “sitting”, “standing”, “motion analysis” and “video” in either the title or abstract. The specific search strategies for each database are included in the appendix. The reference lists of included articles were also screened to locate additional papers missed by the initial search. We considered sitting and standing in any context (i.e. occupational space, laboratory, leisure time etc.) in any country so long as the criteria of “objectively measured time” was met and the outcome of perceived back pain could be directly related to the exposure.

2.2. Title and abstract screen

Exact duplicates of article were identified and removed using RefWorks by the health sciences librarian (MS). Two authors (DD and RG) reviewed the titles and abstracts of the remaining articles and compiled a list of papers that appeared to fit the inclusion criteria. If there was any question of acceptability at this point, the full paper was accessed in order to confirm study details. Reasons for exclusion were documented. The reviewers met to discuss their findings. Discrepancies were resolved through discussion and with consultation of a third reviewer (MS) to reach a final decision when necessary.

Two independent reviewers (DD and RG) extracted data for each for the included studies using a standardized form including the following information: study setting, population demographics and baseline characteristics, details of intervention and control conditions, methodology, recruitment rates and study

dropout numbers, outcome measures (including units and variance).

2.3. Assessment of quality

Following data extraction for each paper, two independent reviewers (DD and RG) completed a quality assessment of all included articles. Specifically, a critical appraisal checklist from the Joanna Briggs Institute was used to assess these quasi-experimental studies [23]. The reviewers met to discuss their findings. Discrepancies were resolved through discussion and with consultation of a third reviewer (MS) to reach a final decision.

2.4. Quantitative analysis

Perceived pain data (continuous data) from included studies were then pooled using an inverse variance random-effects meta-analysis calculating the 95% confidence intervals and two-sided p -values accordingly. Heterogeneity was tested using the Chi-squared test and I^2 statistic and we considered 0.25, 0.5 and 0.75 as low, moderate and high levels of heterogeneity respectively for the I^2 . To visually assess publication bias, a funnel plot was generated. Review Manager 5.3 was used to calculate the analysis and generate the resulting tables and figures an online version of R was used to calculate the Fail-Safe N and Kendall's tau for the funnel plot.

3. Results

The search was performed on October 20, 2017 and re-run on November 14, 2017 and September 2, 2018. Figure 1 presents the PRISMA flowchart for this process. A total of 208 articles were found from all databases and 93 were identified and removed as exact duplicates. One new article was identified through a search of reference lists. Of these 116 articles, four [24–27] were identified as satisfying the inclusion with no exclusions and a full text review was completed to extract data (Table 1) and perform the quality assessment (Table 2). All studies were laboratory-controlled cross-sectional experiments that used each participant as an internal control in order to examine outcome variables between three work posture conditions for one or more hours: standing, sitting and either sit/stand protocol or a perching-type sit/stand chair. For the purposes of this

analysis, only the sitting and standing conditions were used. All studies had measures of perceived back pain discomfort that could be directly related to the posture exposure. Sample sizes of all studies were small, ranging from 12 to 24 participants and only one, Karakolis et al., balanced the population with respect to gender. While all studies included young, healthy individuals, only Karakolis et al. presented detailed exclusion criteria.

All included studies were found to have acceptable quality. At this stage Chester et al., 2002, was excluded from the qualitative and quantitative analyses. This decision was made based on the fact that perceived ratings of comfort were used instead of discomfort and the assumption could not be made that inverting comfort values would approximate discomfort. Further, no standard deviations or p -values for the mean results were provided which directly limits inclusion into the quantitative analysis.

3.1. Qualitative synthesis

With the exception of the study by Le and Marras, which found higher rates of perceived pain in the prolonged standing condition, the majority of included studies [25–27] found little difference in pain ratings between postures.

3.2. Quantitative synthesis

Data from the three included studies [25–27], including a total 56 participants, were pooled for the outcome measure of perceived low back pain using standardized mean differences (Fig. 2). Prolonged standing postures during the completion of a standardized work task in a laboratory controlled environment did not induce significantly less perceived low back pain compared to seated postures (standardized mean difference 0.60, 95% CI –0.68 to 1.87, $p = 0.36$). There was significant heterogeneity observed in this analysis. Specifically, I^2 was 90% which indicates a large amount of heterogeneity not attributable to random error.

A visual analysis of the Forrest Plot (Fig. 2) highlighted Le and Marras (2016) as potentially driving this heterogeneity, perhaps due to the presence of a backrest in the seated condition compared to no backrest in the remaining two studies. Considering this, there was a question of whether this methodological difference impacted the results. Therefore, a secondary analysis was done removing the Le and




Fig. 1. PRISMA flow diagram for search results and selection of included articles for qualitative and quantitative analysis.

217 Marras study. Doing this removed the heterogeneity
218 previously seen ($I^2 = 0\%$), however, did not change
219 the overall result. No statistical difference was found
220 between prolonged standing and sitting in terms
221 of the outcome perceived ratings of low back pain
222 ($p = 0.73$).

223 The potential for publication bias was assessed
224 visually to be low using a funnel plot (Fig. 3). The
225 plot is symmetrical and covers a fair range of standard-
226 ized mean differences. Further, rank correlation test
227 for funnel plot asymmetry, Kendall's tau, was found
228 to be low ($\tau = 0.3333$, $p = 1.0000$). The Fail-Safe N
229 calculation using the Rosenthal Approach found that
230 8 studies would be needed to change the result from
231 insignificant to significant.

232 4. Discussion

233 The results of this study suggest that prolonged
234 standing for desk work does not translate into lower
235 rates of perceived low back pain compared to sit-
236 ting. Both postures induce low back pain that appear
237 to be directly linked to the exposure. Furthermore,
238 it appears that perceived pain induced in all studies
239 are large enough to be clinically meaningful: hav-
240 ing a change of more than two points on a discrete
241 scale and more than 2 cm or 20 mm on a 10 cm or
242 100 mm visual analog scale. Therefore, this suggests
243 that neither posture would be recommended for pro-
244 longed periods of deskwork. To be prudent, however,
245 the results of this study should be interpreted with

Table 1
Details of each study included for qualitative analysis

Article	Study Type	Population	Intervention	Outcome	Results and Conclusions
Seo et al. 1996, Japan	Cross-Sectional	12 healthy subjects (8 males and 4 females), average age 24.1 ± 1.2 years with no current edematous disease or lower leg injury.	Participants were exposed to 3 laboratory-controlled posture exposures in a random order, standing sitting (stool type chair with no backrest) and buttock chair sitting while completing desk work (completing jigsaw puzzles) for 1 hour on three different days.	Lower limb swelling (bioelectric impedance) and subjective ratings of leg dullness, back pain discrete body fatigue.	Subjective complaints increased for all conditions. Higher rates of back pain were found for the buttock chair (5 SD 1) condition and was similar for both standing (3 SD 1) and ordinary chair sitting (3 SD 1).
Karakolis et al. 2016, Canada	Cross-Sectional	24 healthy participants (12 male and 12 female) average age 22.6 ± 1.7 (male) and 23.8 ± 3.0 (female). Participants were excluded if they had experienced an episode of severe non-specific low back pain with in the last 6 months that caused them to miss at least one day of school or work, had backpain at the time of the study, self-identified as developing low back pain from sitting that would lead them to avoid prolonged seated exposures (i.e. a long drive), held a job that involved prolonged standing exposures for more than 10 hours per week or had upper extremity pain that limited their ability to perform typing/mousing tasks.	Participants were exposed to 3, 1 hour, laboratory-controlled posture exposures while completing a standardized desk work task (typing and mousing) on three different days in three postures in a random order: standing, sitting (office chair with backrest removed and alternating standing/sitting in a 3:1 ratio).	Spine posture, perceived low back discomfort (rated on a 100 mm visual analog scale with anchors of 0 = no pain and 100 = worst pain), L4–L5 joint loading work productivity.	All conditions resulted in increasing levels of discomfort. Overall, there was higher discomfort during sitting (27.66 SD 15.77) compared to standing (25.79 , SD 14.70) with the least discomfort perceived in the sit/stand condition (17.15 SD 13.39). A Significant interaction between gender and posture was found (males more discomfort in standing and females more discomfort in sitting).
Le and Marras 2016, USA	Cross-Sectional	20 healthy participants (10 men and 10 women), average age 26.5 ± 8.5 years, with no history of previous or current low back pain in the past 6 months.	Participants exposed to 3 conditions, in a counter-balanced presentation, of sitting (office chair with backrest), standing and perching while completing a standardized desk work task in a laboratory-controlled study. All conditions were completed on the same testing day, for 1 hour duration each, with a 20 minute washout period in between (duration of washout based on pilot work).	Postural transitions, spinal loads, discomfort for various body regions (measured with a 10 cm visual analog scale) and task performance.	Standing had the highest reports of discomfort (23 SD 8), followed by perching (8 SD 5) and seating (9 SD 6).

Table 2

The checklist for quasi-experimental studies (non-randomized experimental studies) from the Joanna Briggs Institute was used to assess quality for the four articles identified as satisfying the inclusion/exclusion criteria of this study. Final results, following review and consultation by two independent researchers, are presented. Based on these findings one study, Chester et al. 2002, was excluded with justification provided

Joanna Briggs Institute Critical Appraisal Checklist for Quasi-Experimental Studies	Articles			
	Chester et al. 2002	Seo et al. 1996	Karakolis et al. 2016	Le and Marras 2016
	Leg swelling, comfort and fatigue when sitting, standing and sit/standing	Leg swelling during continuous standing and sitting work without restricting leg movement.	A comparison of trunk biomechanics, musculoskeletal discomfort and productivity during simulated sit-stand office work.	Evaluating the low back biomechanics of three different office workstations: seated, standing, and perching.
1 Is it clear in the study what is the “cause” and what is the “effect” (i.e. there is no confusion about what variable comes first)?	yes	yes	yes	yes
2 Were the participants included in any comparisons similar?	yes	yes	yes	yes
3 Were the participants included in any comparisons receiving similar treatment/care, other than the exposure or intervention of interest?	yes	yes	yes	yes
4 Was there a control group?	not applicable	not applicable	not applicable	not applicable
5 Were there multiple measurements of the outcome both pre and post the intervention/exposure?	yes	yes	yes	yes
6 Was follow up complete, and if not, were differences between groups in terms of their follow up adequately described and analyzed?	yes	yes	yes	yes
7 Were the outcomes of participants included in any comparisons measured in the same way?	yes	yes	yes	yes
8 Were outcomes measured in a reliable way?	yes	yes	yes	yes
9 Was appropriate statistical analysis used?	yes	yes	yes	yes
Overall Appraisal				
Include		✓	✓	✓
Exclude	✓			
Seek Further Info				
Reasons for Exclusion	(1) Perceived comfort is presented, not a direct opposite of discomfort/pain. (2) No standard deviations are given for mean values.			




Fig. 2. *Top*: Forest plot and measures of heterogeneity for perceived ratings of low back discomfort in laboratory-controlled and objectively measured conditions of prolonged standing versus prolonged sitting desk work postures in healthy adults for all three included studies. *Bottom*: Forest plot and measures of heterogeneity for perceived ratings of low back discomfort in laboratory-controlled and objectively measured conditions of prolonged standing versus prolonged sitting desk work postures in healthy adults with data for Le and Marras (2016) removed.




Fig. 3. To visually assess for publication bias, a funnel plot of the three studies included in the quantitative analysis has been prepared. However, since the plot is balanced with a wide range in study results (Kendall's Tau = 0.3333, $p = 1.0000$), it appears that the risk of publication bias in this case is small. The Fail-Safe N was calculated to be 8 using the Rosenthal approach.

caution, given the cross-sectional study design, small number of included studies, and small sample size of the included studies.

The result that prolonged standing for desk work is no better than sitting in terms of perceived back pain supports opinions that have been expressed in the literature, namely that both standing and sitting, statically sustained for prolonged periods of time can lead to pain and potential injury [28]. To date, no meta-analyses of perceived low back pain in response to sitting and/or standing have been conducted. However, previous systematic reviews, have reached conflicting conclusions about the relationship between posture and pain during desk work. A

number of reports have concluded there is a relationship between sitting [18, 29, 30] and low back pain as well as standing [15, 31] and low back pain while others have found conflicting results for these postures [16, 17, 19, 32]. The relationship with back pain is indeed complex, with many confounding variables such as health history, previous back injury, psychological, social, economic and occupational factors playing a role. While this work highlights that prolonged standing may not be a suitable replacement for prolonged sitting at work, more studies, using higher level designs, are needed to generate high quality evidence upon which to base occupational standards and recommendations.

The heterogeneity in the main analysis was high ($I^2 = 90\%$). While the included studies were very similar in terms of experimental setting, health status, age range and objective measures there were some key differences that could have contributed to this finding. For instance, Karakolis et al. found significant differences in perceived discomfort ratings between men and women: with men perceiving higher rates of back pain in standing and women perceiving higher rates in sitting. Since most articles included in this analysis contained datasets with unbalanced genders and did not present data separately for males and females, there is the possibility that these differences contributed to the observed heterogeneity. However, considering the heterogeneity dropped to 0% with the removal of the Le and Marras study, it is reasonable to conclude that the factor driving the large heterogeneity must have been the presence of a backrest. The sensitivity of the results to the presence/absence of a backrest during sitting was tested during the analysis by

246
247
248
249
250
251
252
253
254
255
256
257
258
259

260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293

removing the study by Le and Marras and re-analyzing the data. While the result was unaffected in this case, it can be argued that this factor needs to be better addressed with more research before drawing a firm conclusion; especially since logically the ability to recline back and unload body stress on a support would result in lower amounts of perceived back pain.

This study was strengthened by using data from studies in which the exposure was objectively measured. This is particularly important as recent evidence suggests that self-recall for posture exposure, beyond TV viewing, is poor [33]. However, this study was limited by a several factors. First, there were a small number of (small) cross-sectional studies that fit the inclusion criteria (total participant sample = 56) and, while a within-control design was used, there were no true control groups in the design of any of these studies. Secondly, the data extracted from two papers (Le and Marras 2016 and Seo et al. 1996) were estimated from graphs, as tabulated data were not provided. Further, since only Karakolis et al. presented data separately by gender, a subgroup analysis could not be performed in this meta-analysis. Future work should take care to study and present data for males and females since Karakolis et al. found a significant interaction for gender and posture and these underlying differences can contribute to heterogeneity and may warrant gender-specific recommendations for guidelines.

5. Conclusion

Replacing sitting with standing for deskwork would not be recommended based on the results of this study; indeed both postures appear to be a problem. However, this interpretation must be accepted with caution due to the study type, number and size or the studies included in this analysis. To best answer this question, larger studies, with objective measures conducted preferably in a field setting and including a wider age range and sub-grouped by gender and health history should be performed to generate high quality evidence upon which to base ergonomic guidelines.

Conflict of interest

The authors have no conflicts of interest or funding to declare for this project.

References

- [1] GBD 2013 DALYs and HALE Collaborators, Murray CJ, Barber RM, Foreman KJ, Abbasoglu Ozgoren A, Abd-Allah F, et al. Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990–2013: quantifying the epidemiological transition. *Lancet*. 2015;386(10009):2145-91.
- [2] Dagenais S, Caro J, Haldeman S. A systematic review of low back pain cost of illness studies in the United States and internationally. *Spine J*. 2008;8(1):8-20.
- [3] Luo X, Pietrobon R, Sun SX, Liu GG, Hey L. Estimates and patterns of direct health care expenditures among individuals with back pain in the United States. *Spine (Phila Pa1976)*. 2004;29(1):79-86.
- [4] Pengel LH, Herbert RD, Maher CG, Refshauge KM. Acute low back pain: systematic review of its prognosis. *BMJ*. 2003;327(7410):323.
- [5] Von Korff M, Saunders K. The course of back pain in primary care. *Spine (Phila Pa1976)*. 1996;21(24):2833-7; discussion 2838-9.
- [6] Fenety A, Putnam C, Walker JM. In-chair movement: validity, reliability and implications for measuring sitting discomfort. *ApplErgon*. 2000;31(4):383-93.
- [7] Beach TA, Parkinson RJ, Stothart JP, Callaghan JP. Effects of prolonged sitting on the passive flexion stiffness of the *in vivo* lumbar spine. *Spine J*. 2005;5(2):145-54.
- [8] De Carvalho DE, Callaghan JP. Passive stiffness changes in the lumbar spine and effect of gender during prolonged simulated driving. *IntJIndErgonomics*. 2011;41(6):617-24.
- [9] Dunk NM, Callaghan JP. Gender-based differences in postural responses to seated exposures. *ClinBiomech(Bristol, Avon)*. 2005;20(10):1101-10.
- [10] Gregory DE, Dunk NM, Callaghan JP. Stability ball versus office chair: comparison of muscle activation and lumbar spine posture during prolonged sitting. *HumFactors*. 2006;48(1):142-53.
- [11] van der Ploeg HP, Moller SV, Hannerz H, van der Beek AJ, Holtermann A. Temporal changes in occupational sitting time in the Danish workforce and associations with all-cause mortality: results from the Danish work environment cohort study. *IntJBehavNutrPhysAct*. 2015;12(Journal Article):71-015-0233-1.
- [12] Clemes SA, O'Connell SE, Edwardson CL. Office workers' objectively measured sedentary behavior and physical activity during and outside working hours. *JOccupEnvironMed*. 2014;56(3):298-303.
- [13] Jans MP, Proper KI, Hildebrandt VH. Sedentary behavior in Dutch workers: differences between occupations and business sectors. *AmJPrevMed*. 2007;33(6):450-4.
- [14] Kazi A, Duncan M, Clemes S, Haslam C. A survey of sitting time among UK employees. *OccupMed(Lond)*. 2014;64(7):497-502.
- [15] Nelson-Wong E, Callaghan JP. Transient low back pain development during standing predicts future clinical low back pain in previously asymptomatic individuals. *Spine (Phila Pa1976)*. 2014;39(6):E379-83.
- [16] Hartvigsen J, Leboeuf-Yde C, Lings S, Corder EH. Is sitting-while-at-work associated with low back pain? A systematic, critical literature review. *ScandJPublic Health*. 2000;28(3):230-9.

- 403 [17] Chen SM, Liu MF, Cook J, Bass S, Lo SK. Sedentary
404 lifestyle as a risk factor for low back pain: a system-
405 atic review. *IntArchOccupEnvironHealth*. 2009;82(7):
406 797-806.
- 407 [18] Frymoyer JW, Pope MH, Costanza MC, Rosen JC, Goggin
408 JE, Wilder DG. Epidemiologic studies of low-back pain.
409 *Spine*. 1980;5(5):419-23.
- 410 [19] Roffey DM, Wai EK, Bishop P, Kwon BK, Dagenais S.
411 Causal assessment of occupational sitting and low back
412 pain: results of a systematic review. *Spine J*. 2010;10(3):
413 252-61.
- 414 [20] Magora A. Investigation of the relation between low back
415 pain and occupation. 3. Physical requirements: sitting,
416 standing and weight lifting. *IMS IndMedSurg*. 1972;41(12):
417 5-9.
- 418 [21] Foley B, Engelen L, Gale J, Bauman A, Mackey M.
419 Sedentary Behavior and Musculoskeletal Discomfort Are
420 Reduced When Office Workers Trial an Activity-Based
421 Work Environment. *JOccupEnvironMed*. 2016;58(9):
422 924-31.
- 423 [22] Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA
424 Group. Preferred reporting items for systematic reviews
425 and meta-analyses: the PRISMA statement. *IntJSurg*.
426 2010;8(5):336-41.
- 427 [23] Aromataris E, Munn Z. Joanna Briggs Institute Reviewer's
428 Manual [Internet]. 2017. (Tufanaru C, Munn Z, Aromataris
429 E, Campbell J, Hopp L, editors. Systematic Reviews of
430 effectiveness.). Available from: https://joannabriggs.org/assets/docs/critical-appraisal-tools/JBI-Quasi-Experimental-Appraisal_Tool2017.pdf
- 431 [24] Chester MR, Rys MJ, Konz SA. Leg swelling, comfort and
432 fatigue when sitting, standing, and sit/standing. *IntJIndErgon*.
433 2002;29(5):289-96.
- 434
- 435 [25] Karakolis T, Barrett J, Callaghan JP. A comparison of trunk
436 biomechanics, musculoskeletal discomfort and productiv-
437 ity during simulated sit-stand office work. *Ergonomics*.
438 2016;59(10):1275-87.
- 439 [26] Le P, Marras WS. Evaluating the low back biomechanics
440 of three different office workstations: Seated, standing, and
441 perching. *Appl Ergon*. 2016;56:170-8.
- 442 [27] Seo A, Kakehashi M, Tsuru S, Yoshinaga F. Leg Swelling
443 during Continuous Standing and Sitting Work without
444 Restricting Leg Movement. *Journal of Occupational Health*.
445 1996;38(4):186-9.
- 446 [28] Callaghan J, De Carvalho D, Gallagher K, Karakolis T,
447 Nelson-Wong E. Is Standing the Solution to Sedentary
448 Office Work? *Ergonomics in Design*. 2015;23(3)(Journal
449 Article):20-4.
- 450 [29] Damkot DK, Pope MH, Lord J, Frymoyer JW. The relation-
451 ship between work history, work environment and low-back
452 pain in men. *Spine*. 1984;9(4):395-9.
- 453 [30] Wilder DG, Pope MH. Epidemiological and aetiological
454 aspects of low back pain in vibration environments - an
455 update. *ClinBiomech(Bristol, Avon)*. 1996;11(2):61-73.
- 456 [31] Coenen P, Parry S, Willenberg L, Shi JW, Romero L, Black-
457 wood DM, et al. Associations of prolonged standing with
458 musculoskeletal symptoms-A systematic review of labora-
459 tory studies. *Gait Posture*. 2017;58(Journal Article):310-8.
- 460 [32] Roffey DM, Wai EK, Bishop P, Kwon BK, Dagenais S.
461 Causal assessment of occupational standing or walking and
462 low back pain: results of a systematic review. *Spine J*.
463 2010;10(3):262-72.
- 464 [33] Clark BK, Thorp AA, Winkler EA, Gardiner PA, Healy GN,
465 Owen N, et al. Validity of self-reported measures of work-
466 place sitting time and breaks in sitting time. *MedSciSports
467 Exerc*. 2011;43(10):1907-12.

468 **Appendix**469 *PubMed (Medline)*

470 (“Back Pain”[Mesh] OR ((discomfort[tw] OR
471 pain*[tw] OR ache*[tw] OR aching[tw]) AND (back
472 [tw] OR thoracic[tw] OR thorax[tw] OR lumbar
473 [tw])) OR backache*[tw] OR dorsalgia[tw] OR LBP
474 [tw] OR lumbago[tw] OR “Sciatica” [Mesh] OR
475 sciatica[tw] OR “Radiculopathy” [Mesh] OR radicu-
476 lopath*[tw]) AND (sitting[tw]) AND (standing[tw])
477 AND (“Monitoring, Ambulatory” [Mesh] OR “Mon-
478 itoring, Physiologic” [Mesh] OR “Video Recording”
479 [Mesh] OR “Task Performance and Analysis” [Mesh]
480 OR “Accelerometry” [Mesh] OR “Actigraphy” [Me-
481 sh] OR acceleromet*[tw] OR actigraph*[tw] OR gy-
482 roscop*[tw] OR sensor[tw] OR sensors[tw] OR
483 device*[tw] OR wearable[tw] OR inertial[tw] OR
484 “motion capture” [tw] OR “motion analysis” [tw]
485 OR lab[tw] OR laborator*[tw] OR video[tw] OR
486 videorecord*[tw] OR videotap*[tw] OR “objectively
487 measured” [tw])

488 *Embase*

489 (‘backache’/exp OR ((discomfort:ab,ti OR pain*:
490 ab,ti OR ache*:ab,ti OR aching:ab,ti) AND (back:
491 ab,ti OR thoracic:ab,ti OR thorax:ab,ti OR lumbar:
492 ab,ti)) OR backache*:ab,ti OR dorsalgia:ab,ti OR
493 lbp:ab,ti OR lumbago:ab,ti OR ‘sciatica’/de OR sci-
494 atica:ab,ti OR ‘radiculopathy’/exp OR radiculopath*:
495 ab,ti) AND sitting:ab,ti AND standing:ab,ti AND
496 (‘ambulatory monitoring’/de OR ‘physiologic moni-
497 toring’/exp OR ‘videorecording’/de OR ‘task perfor-
498 mance’/de OR ‘accelerometry’/de OR ‘actimetry’/de
499 OR acceleromet*:ab,ti OR actigraph*:ab,ti OR gyro-
500 scop*:ab,ti OR sensor:ab,ti OR sensors:ab,ti OR de-
501 vice*:ab,ti OR wearable:ab,ti OR inertial:ab,ti OR
502 ‘motion capture’:ab,ti OR ‘motion analysis’:ab,ti OR
503 lab:ab,ti OR laborator*:ab,ti OR video*:ab,ti OR ‘ob-
504 jectively measured’:ab,ti)

505 *Cinahl*

506 (MH “Back Pain” OR ((TI discomfort OR TI pain*
507 OR TI ache* OR TI aching OR AB discomfort OR
508 AB pain* OR AB ache* OR AB aching) AND (TI
509 back OR TI thoracic OR TI thorax OR TI lumbar
510 OR AB back OR AB thoracic OR AB thorax OR AB
lumbar)) OR TI backache* OR AB backache* OR TI

dorsalgia OR AB dorsalgia OR TI LBP OR AB LBP
OR TI lumbago OR AB lumbago OR MH “Sciatica”
OR TI sciatica OR AB sciatica OR MH “Radicu-
lopathy” OR TI radiculopath* OR AB radiculopath*)
AND (MH “Sitting” OR TI sitting OR AB sitting)
AND (MH “Standing” OR TI standing OR AB stand-
ing) AND (MH “Monitoring, Physiologic” OR MH
“Videorecording” OR MH “Task Performance and
Analysis” OR MH “Accelerometry” OR MH “Actig-
raphy” OR TI acceleromet* OR AB acceleromet* OR
TI actigraph* OR AB actigraph* OR TI gyroscop*
OR AB gyroscop* OR TI sensor OR AB sensor OR
TI sensors OR AB sensors OR TI device* OR AB
device* OR TI wearable OR AB wearable OR TI in-
ertial OR AB inertial OR TI “motion capture” OR AB
“motion capture” OR TI “motion analysis” OR AB
“motion analysis” OR TI lab OR AB lab OR TI lab-
orator* OR AB laborator* OR TI video* OR AB
video* OR TI “objectively measured” OR AB “objec-
tively measured”)

511 *SPORTDiscus*

(DE “BACKACHE” OR DE “LUMBAR pain” OR
((TI discomfort OR TI pain* OR TI ache* OR TI a-
ching OR AB discomfort OR AB pain* OR AB ache*
OR AB aching) AND (TI back OR TI thoracic OR TI
thorax OR TI lumbar OR AB back OR AB thoracic
OR AB thorax OR AB lumbar)) OR TI backache*
OR AB backache* OR TI dorsalgia OR AB dorsalgia
OR TI LBP OR AB LBP OR TI lumbago OR AB
lumbago OR DE “SCIATICA” OR TI sciatica OR AB
sciatica OR DE “RADICULOPATHY” OR TI radicu-
lopath* OR AB radiculopath*) AND (DE “SITTING
position” OR TI sitting OR AB sitting) AND (DE
“STANDING position” OR TI standing OR AB stand-
ing) AND (DE “PATIENT monitoring” OR DE “AC-
CELEROMETERS” OR DE “SPEEDOMETERS”
OR TI acceleromet* OR AB acceleromet* OR TI acti-
graph* OR AB actigraph* OR TI gyroscop* OR AB
gyroscop* OR TI sensor OR AB sensor OR TI sen-
sors OR AB sensors OR TI device* OR AB device*
OR TI wearable OR AB wearable OR TI inertial OR
AB inertial OR TI “motion capture” OR AB “motion
capture” OR TI “motion analysis” OR AB “motion
analysis” OR TI lab OR AB lab OR TI laborator*
OR AB laborator* OR TI video* OR AB video* OR
TI “objectively measured” OR AB “objectively
measured”)