

Erratum

CYP1A1 genetic polymorphisms in Ecuador, South America

By C. Paz-y-Miño, M. Arévalo, M.J. Muñoz G. and P.E. Leone

[*Disease Markers* 21(2) 2005, pp. 57–59]

Tables 1 and 2 have been omitted from the article.

Table 1
CYP 1A1 allelic frequencies

	N	Alleles	Frequencies	rare Allele
<i>Msp I</i>				
m2m2	30	60	0.17	
m1m2	40	80	0.22	0.70m2
m2m2	103	206	0.59	
<i>Ile462Val</i>				
ile/ile	4	8	0.01	
ile/val	164	328	0.94	0.5val
val/val	6	12	0.03	

Table 2
Allele frequency of *CYP1A1* polymorphism in different populations

Population	N	chromosomes	allele m2	N	chromosomes	allele val
Ecuador	174	348	0.70	174	348	0.50
<i>Brazil [2]</i>						
Xavante	21	42	0.95	21	42	0.97
Wai Wai	26	52	0.81	26	52	0.81
Zoró	30	60	0.87	30	60	0.76
Gavião	30	60	0.72	30	60	0.59
Surui	24	48	0.96	24	48	0.54
<i>Chile</i>						
Mapuche [6]	84	168	0.83	84	168	0.77
General [10]	194	388	0.30	180	360	0.37
Japan [5]	106	212	0.33	106	212	0.22
<i>China</i>						
[12]				1089	2178	0.19
[9]				414	828	0.29
[12]	91	182	0.37			
<i>Hawaii</i>						
[5]	22	44	0.45	22	44	0.25
[13]	311	622	0.26			
Taiwan [14]	470	940	0.35	470	940	0.27
Spain [15]	115	230	0.38	115	230	0.047
<i>Caucasic</i>						
Sweden [16]	70	140	0.1	100	200	0.18
RepChec [17]	645	1290	0.08			
UK [18]	3303	6606	0.036			
[5]	60	120	0.12	60	120	0.05