

Review

Role and Mechanism of Vitamin A Metabolism in the Pathophysiology of Parkinson's Disease

Anaïs Marie^a, Morgane Darricau^{a,b,1}, Katia Touyarot^{a,2},
Louise C. Parr-Brownlie^{c,d,3} and Clémentine Bosch-Bouju^{a,4,*}

^aUniversity Bordeaux, INRAE, Bordeaux INP, NutriNeuro, UMR 1286, Bordeaux, France

^bUniversity Bordeaux, Institut des Maladies Neurodégénératives, UMR 5293, Bordeaux, France

^cDepartment of Anatomy, Brain Health Research Centre, University of Otago, Dunedin, New Zealand

^dBrain Research New Zealand (Center of Research Excellence), Dunedin, New Zealand

Accepted 12 May 2021

Pre-press 5 June 2021

Abstract. Evidence shows that altered retinoic acid signaling may contribute to the pathogenesis and pathophysiology of Parkinson's disease (PD). Retinoic acid is the bioactive derivative of the lipophilic vitamin A. Vitamin A is involved in several important homeostatic processes, such as cell differentiation, antioxidant activity, inflammation and neuronal plasticity. The role of vitamin A and its derivatives in the pathogenesis and pathophysiology of neurodegenerative diseases, and their potential as therapeutics, has drawn attention for more than 10 years. However, the literature sits in disparate fields. Vitamin A could act at the crossroad of multiple environmental and genetic factors of PD. The purpose of this review is to outline what is known about the role of vitamin A metabolism in the pathogenesis and pathophysiology of PD. We examine key biological systems and mechanisms that are under the control of vitamin A and its derivatives, which are (or could be) exploited for therapeutic potential in PD: the survival of dopaminergic neurons, oxidative stress, neuroinflammation, circadian rhythms, homeostasis of the enteric nervous system, and hormonal systems. We focus on the pivotal role of ALDH1A1, an enzyme expressed by dopaminergic neurons for the detoxification of these neurons, which is under the control of retinoic acid. By providing an integrated summary, this review will guide future studies on the potential role of vitamin A in the management of symptoms, health and wellbeing for PD patients.

Keywords: Neuroinflammation, vitamin A, retinoic acid, ALDH1A1, oxidative stress, RAR RXR receptors

INTRODUCTION

Even though pharmacological treatments exist to control motor symptoms in Parkinson's disease (PD),

patients often require more holistic care in order to improve their daily quality of life and wellbeing. As such, nutrition becomes increasingly a focus for patients, medical teams and caregivers. In addition to influencing the daily management of the disease, nutrition is a potential disease-modifying factor that can either slow down the course of neurodegeneration when it is optimized, or worsen it when inadequate. Therefore, it is critical to understand what is suitable for parkinsonian patients in terms of food habits, diet, and nutrients to maximize health and wellbeing. So far, there is no consensus

¹ORCID: 0000-0002-6663-3156

²ORCID: 0000-0003-4310-4152

³ORCID: 0000-0002-3001-7669

⁴ORCID: 0000-0001-8869-768X

*Correspondence to: Clémentine Bosch-Bouju, Univ. Bordeaux, INRAE, Bordeaux INP, NutriNeuro, UMR 1286, F-33000, Bordeaux, France. Tel.: +335 5757 9235; E-mail: cbosch@enscbp.fr.

for an ideal diet or crucial ingredient. However, in recent years, researchers and clinicians have focused attention on candidate nutrients, such as omega-3 polyunsaturated fatty acids or vitamins. Widespread evidence, mainly in the preclinical literature, suggests that altered vitamin A signaling may contribute to the pathophysiology of PD. However, no clear consensus has been made on the underlying mechanisms.

Vitamin A is a lipophilic molecule that is exclusively provided by food. Vitamin A is the unique precursor of retinoic acid (RA), a molecule that modulates gene transcription through its binding to nuclear receptors [1]. Through its derivatives, vitamin A is involved in many important homeostatic processes in the organism, such as cell differentiation, antioxidant power, inflammation and neuronal plasticity. The role of vitamin A derivatives in the pathophysiology of neurodegenerative diseases, and their potential as therapeutics, has been carefully studied for more than 10 years, especially for Alzheimer's disease, and to a lesser extent PD [2–8]. However, the literature sits in disparate fields, does not focus on the importance of vitamin A metabolism, and has not been concisely summarized with the view of maximizing PD patient wellbeing.

The purpose of this review is to outline what is known about the role of vitamin A metabolism in the pathogenesis and pathophysiology of PD. We examine key biological systems and mechanisms that are under the control of vitamin A and its bioactive derivative, RA, which are (or could be) exploited for therapeutic potential in PD: the survival of dopaminergic neurons in the substantia nigra *pars compacta* (SNc), oxidative stress, and function of immune, enteric nervous, and hormonal systems. By providing an integrated summary, this review will guide future studies on the potential role of vitamin A in the management of symptoms, health and wellbeing for PD patients.

CLINICAL AND PRECLINICAL EVIDENCE FOR A ROLE OF VITAMIN A IN THE PATHOGENESIS AND PATHOPHYSIOLOGY OF PD

Metabolism and function of vitamin A

Vitamin A (retinol) is a fat-soluble vitamin that is essential for the organism. Vitamin A is found in foods derived from animals (meat, milk, eggs) in the form of retinol esters, or in vegetable sources in the

form of carotenoids that are metabolized to retinol by intestinal enzymes [9, 10]. The bioactive metabolites of vitamin A form the group of retinoid compounds or retinoids (Fig. 1). The main active metabolite is all-*trans* RA, called RA in this review, but other isomers also display biological activity, such as 9-*cis* or 13-*cis*-RA [9] (Fig. 1). RA acts through nuclear receptors of the steroid receptor family, namely RA receptors (RAR), which can exist in 3 different subtypes: RAR- α , RAR- β and RAR- γ [3]. The receptor subtypes differ mainly by their different locations in the body, and particularly in different brain areas [11]. RAR act as heterodimers with retinoid-X receptor (RXR) to bind DNA on RA response element (RARE) sequences located in promoter regions to activate (or repress) gene expression. RXRs also exist in 3 isoforms, RXR- α , RXR- β and RXR- γ and they function as a heterodimer partner for many other nuclear receptors [12], such as thyroid hormone receptor. The signaling of RXR is very complex and not yet fully understood. Of note, multiple ligands have been found for RXR, including retinoids, but the endogenous ligand that is most likely to activate RXR is 9-*cis*-dihydro RA (9CDHRA) [13]. Furthermore, RXR can also function without ligands in some cases [12–14].

Through its nuclear receptors (more often RAR-RXR heterodimers), RA controls the transcription of hundreds of genes, belonging to various gene pathways, including dopamine signaling and plasticity [3, 15]. Throughout life, RA is involved in cell growth, differentiation and cell death: during development, the action of RA is crucial since it is involved in the patterning of tissues, especially in the central nervous system [1, 5]; at adult age, RA continues to be involved in the maintenance and homeostasis of cells [16], and controls inflammatory responses, hormone actions, reproduction and vision [16]. The endocrine system is also tightly related to the retinoid system because RXR forms heterodimers (sometimes in an obligatory role) for some hormone nuclear receptors, such as thyroid or vitamin D receptors [17].

Vitamin A and RA are stored in liver cells, and are released on demand to act at target organs, such as the brain. Vitamin A is transformed into retinaldehyde by retinol dehydrogenase (RDH) enzymes, and retinaldehyde is further transformed in RA by retinaldehyde dehydrogenase (RALDH) enzymes [5, 10] (Fig. 2). Due to their lipophilic nature, retinoids are carried by specific proteins in the blood (retinol binding proteins, RBP) and in cells (cellular retinol




Fig. 1. Chemical structures of the main retinoids metabolized in the body. Retinol (vitamin A) is directly provided by animal sources. Carotenoids, such as β -carotene, are precursors of retinol from vegetable sources that can be converted into retinol by the organism. Retinol is metabolized into retinal by enzymes of the RDHs family, which can also be converted back to retinol. Retinal can be metabolized into 11 cis retinal. A key bioactive metabolite produced from retinal is RA (*all trans* retinoic acid), which is irreversibly metabolized by a RALDH protein (ALDH1A1 belongs to RALDHs). RA can be metabolized in 11-Cis Retinal, which is an active metabolite in the retina, and in 9-cis RA, another key active metabolite for the brain. Alternatively, retinol can be transformed in 9CDHRA, an endogenous RXR ligand. RDHs, retinol dehydrogenases; RALDHs, retinaldehydes dehydrogenases.

binding proteins, CRBP, and cellular retinoic acid binding proteins, CRABP) to be taken to, and have action, at other organs (Fig. 2). Intriguingly, extracellular carriers (RBP) and the transmembrane transporter (STRA6 is stimulated by retinoic acid 6) have been identified for retinol, but not for RA, yet RA is clearly present in the blood [18]. In addition, STRA6 is not present on every cell, suggesting passive diffusion or another transporter, not yet identified, play a role [19].

In the brain, all the machinery for retinoid metabolism/catabolism, receptors and transport are present, with a specific distribution of these proteins depending on the brain structure and cell type [5, 10, 11, 20, 21] (Fig. 2). Remarkably in the hippocampus, RA has been reported to be synthesized by the meninges whereas RA has action at hippocampal neurons [20]. In the dopaminergic nigro-striatal pathway, it is likely that RA synthesis and RA action are also unevenly distributed: the enzymes for produc-

tion of RA are concentrated in SNc dopaminergic neurons [3, 21, 22], whereas RA receptors are mostly expressed in striatal neurons [11]. This is in line with the paracrine action of RA that has been described [1, 23], and recent data show that RA can act as a retrograde signaling molecule in this pathway [24]. In addition, RA signaling has been involved in the patterning of a specific dopaminergic nigrostriatal pathway [25]. Of note, a recent study reports that alpha-synuclein is a RA-carrier protein that regulates transcriptional activity of retinoids [26]. This evidence is in line with increasing literature showing that alpha-synuclein is a lipid-binding protein [27]. Yet, alpha-synuclein aggregates have been found to be expressed trans-synaptically in humans with Lewy-bodies dementia [28]. Thus, alpha-synuclein could be an extracellular carrier of RA in the nigrostriatal pathway, but this has not been fully proven yet (Fig. 2). In the study [26], RA triggers the translocation of alpha-synuclein in the nucleus in SH-SY5Y.




Fig. 2. Model of RA metabolism and signaling in the nigro-striatal pathway. Retinol (vitamin A) is transported in the extracellular fluids by RBP, and internalized into the cells by STRA6. Retinol can also diffuse across the plasma membrane or through other transporters, not yet identified. Retinol is then bound to CRBP and metabolized into retinal by enzymes of the RDHs family. Retinal is further metabolized into RA by a RALDH protein (notably ALDH1A1 in a sub-population of SNc neurons). RA is then transported to the nucleus bound to CRABP protein. In the nucleus, RA binds to RA receptors (RARs), which activate the control of gene expression by RAR/RXR dimers on genes with a RARE sequence in their promoter. Furthermore, as a trans-synaptic factor, RA can travel trans-synaptically from SNc neurons to striatum neurons [24]. From the literature, it is possible that alpha-synuclein may serve as a cargo protein for the trans-synaptic transport of RA [26]. Finally, RA can be degraded by the Cyp26B1 enzyme. RBP, retinol binding proteins; STRA6, transporter stimulated by retinoic acid 6; CRBP, cellular retinol binding protein; CRABP, cellular retinoic acid binding protein; RDHs, retinol dehydrogenases; RALDHs, retinaldehyde dehydrogenases family (including ALDH1A1); RARE, retinoic acid receptor response element.

Here, we postulate that a lack of RA could impede nuclear translocation of alpha-synuclein, thus favoring accumulation and aggregation. The balance of RA's therapeutic and pathogenic roles needs to be elucidated.

As for other target organs, retinoid signaling is important for brain function throughout life. During development, retinoid signaling is a determinant for neuronal development, maturation and differentiation [1]. In particular, RA signaling is involved in the neurogenesis and differentiation of striatal neurons [24, 29–32]. Studies with transgenic mice demonstrate that RAR β was necessary for the differentiation of D1⁺ medium size spiny neurons of the striatum [29],

as well as the formation of striosomes [32]. Other studies show that ALDH1A1 was necessary for the expression of μ opioid receptors (highly expressed in striosomes) [24] while RALDH3 induced the GABAergic phenotype of striatal neurons through the synthesis of RA [31]. An *in vitro* study also showed that DARPP-32, a marker of medium size spiny neurons, was induced by RA [30]. The late phase of dopaminergic neuron differentiation is also linked to proteins involved in RA signaling, such as ALDH1A1 or NURR1 (an orphan nuclear receptor expressed in dopaminergic neurons that dimerizes with RXR) [33–36]. Interestingly, a recent study with embryonic carcinoma tissue showed that differentiation of cells

can follow striatal or dopaminergic fates, depending on the RAR subtypes that are activated [37].

In the adult brain, retinoid signaling is involved in synaptic plasticity [3, 5, 38]. On the one hand, vitamin A deficiency or dysfunction in RAR/RXR signaling (transgenic RXR^{-/-} and/or RAR^{-/-} mice) alters long term plasticity in the hippocampus [20, 39–42]. On the other hand, RA controls synaptic scaling, an essential mechanism that allows homeostatic synaptic plasticity [38]. When a neuronal network is deprived of excitatory activity, the strength of synapses is increased in order to compensate for this low activity, which is called synaptic scaling. RA is a major actor of synaptic scaling, through various processes depending on the system studied, such as externalization of AMPA receptors or internalization of GABA_A receptors [43–49]. Of note, these mechanisms can involve non-genomic action of RA and their receptors by stimulating local dendritic protein synthesis [38, 45, 50]. These mechanisms have been described in cell cultures or in the hippocampus, but rarely in the basal ganglia. One recent study showed that RA regulates levels of AMPA receptors in cultured nucleus accumbens neurons [51].

With aging, retinoid signaling remains necessary for brain function, however, aging processes and senescence eventually reduces vitamin A function and efficacy. Indeed, data show that the liver reduces its ability to release vitamin A and RA with age, which can lead to decreased bioavailability for target organs, especially the brain [18, 52, 53]. This leads to reduced neurogenesis and reduced dendritic tree arborization for pyramidal neurons in the hippocampus [18, 54]. Thus, the decreased bioavailability of vitamin A with age could contribute to age-related cognitive decline, and the development of neurodegenerative processes that underlie Alzheimer's disease and PD. Interestingly, increasing dietary vitamin A intake can prevent the decrease of vitamin A bioavailability [18, 53, 54]. Indeed, nutritional studies show that dietary vitamin A is stored in the liver and a proportion of vitamin A is directly transported to the brain through the portal vein system [55–57]. A recent study showed that this 'secondary pathway' for vitamin A absorption is particularly dedicated to the brain [57]. Moreover, vitamin A supplementation increases the amount of vitamin A absorption through this pathway, favoring brain supply and 'escaping' or by-passing liver storage [57]. Therefore, vitamin A supplementation may be a therapeutic lever to improve bioavailability of vitamin A, and thus retinoid signaling, in the brain.

Vitamin A and PD: Clinical studies

Epidemiological and clinical studies investigating the link between vitamin A or retinoids/carotenoids and PD have not shown a clear effect (Table 1). Original studies, meta-analyses and reviews have mainly found an absence of link between the disease and the dietary intake [58–66]. More recent studies found an association between low dietary intake of β -carotene, or low levels of blood α -carotene, β -carotene and lycopene with the progression of PD [67, 68]. These studies are based on nutritional intake or retinol blood levels, and therefore they do not provide information about the vitamin A status of patients, i.e., the bioavailability of vitamin A and RA for target organs, such as the brain. The level of retinol in the blood is correlated to nutritional intake, but is also tightly controlled by homeostatic processes to maintain a constant value around 2 μ g/ml [52]. Thus, the retinol level in the blood is not a reliable marker of vitamin A and RA functionality. More relevant measures would be gene expression of RA receptors (RAR/RXR) or retinol level in the cerebral spinal fluid [69]. Measuring RA concentration in the blood or the cerebral spinal fluid would also be more informative, but it is highly challenging due to its biochemical instability [70].

Of note, ALDH1A1, a synthetic enzyme for RA, has been described as a biomarker for PD, since its reduced expression has been found in the blood and brain of PD patients [71–73] (see below).

Vitamin A and PD: Preclinical studies

Despite the lack of association between dietary vitamin A and PD in humans, important links have been established between the pathophysiology of the disease and proteins involved in the metabolism of vitamin A and retinoids. Preclinical evidence for vitamin A's role in the pathogenesis of PD has been established by various manipulations of vitamin A and its pathways: dietary supplementation, dietary deficiency, knock-out mice for retinoid receptors, treatments with vitamin A derivatives *in vivo* or *in vitro*.

First, evidence came from the discovery that the development and fate of the nigro-striatal pathway is under the control of nuclear receptors from the retinoid receptor family, namely RAR, RXR and Nurr1 [8, 34, 36, 74, 75]. Of note, the striatum is the brain structure that expresses the most RAR β and RXR γ [11] and expression of D1 and D2 recep-

Table 1

Summary of the main clinical studies that have directly or indirectly investigated the relationship between vitamin A and Parkinson's disease

Design	Factors quantified	sample size	Method	SNc changes*	Main outcome(s)	Study
Cased-control study	Dietary factors in PD	103 sporadic PD, 103 control cases	Food frequency questionnaire		No association between consumption of vitamin A-containing foods and PD.	[58]
Cased-control study	Serum levels of retinoids in a PD case-control study	61 sporadic PD, 61 control cases	Whole blood collected		No association between carotenoids / retinoids serum levels and PD.	[62]
Cased-control study	Dietary factors in PD	144 sporadic PD, 432 control cases	Food frequency questionnaire		No association between consumption of vitamin A-containing foods and PD.	[63]
Cased-control study	Serum levels of retinoids in a PD case-control study	104 sporadic PD, 52 control cases	Venous blood collected		Serum level of β -carotene (but not retinol) is lower for PD patients compared to control, and is also lower in early PD group than in advanced PD group.	[67]
Cased-control study	Reduction of ALDH1A1 expression in DA neurons	13 sporadic PD, 14 control cases	Brain SNc sections	ALDH1A1 \searrow	ALDH1 mRNA is specifically and markedly down-regulated in DA neurons of SNc in PD.	[71]
Cased-control study	ALDH1A1 as a peripheral biomarker for diagnosing sporadic PD	22 sporadic PD, 33 control cases	Whole blood collected		ALDH1A1 is a biomarker with high sensitivity and specificity to identify the risk of developing PD.	[72]
Cased-control study	Reduction of ALDH1A1 expression in DA neurons	10 sporadic PD, 9 control cases	Brain SNc sections	ALDH1A1 \searrow DA neurons \searrow	The clustering of ALDH1A1-positive and -negative DA neurons in human SNc is different between PD and control cases. A reduction of ALDH1A1 may render DA neurons more prone to degenerate.	[73]
Cased-control study	Serum levels of retinoids in a PD case-control study	41 sporadic PD, 41 control cases	Whole blood collected		No significant changes of retinoids and carotenoids serum levels in PD patients.	[207]
Prospective study	Risk factors for PD	13,979 people at start; 395 incident PD and 2,320 control cases, 17 years later	Food frequency questionnaire		No significant association between vitamin A intake and PD.	[64]
Prospective study	Link between dietary antioxidant vitamins and risk of PD	371 incident PD in 124,221 population cohort	Food frequency questionnaire		High intakes of carotenoids do not appear to reduce the risk of PD.	[66]
Prospective study	Link between dietary antioxidant vitamins and risk of PD	1,329 incident PD in 84,774 population cohort	Food frequency questionnaire		Dietary intake of β -carotene is associated with a lower risk of PD.	[68]

*compared to control cases.

Table 2

Summary of the main studies relating the impact of vitamin A pathway alterations (dietary intake, receptors or ALDH1A1) in Parkinson's disease rodent models. VAD, vitamin A deficient; D1R, dopamine D1 receptor; D2R, dopamine D2 receptor; TH, tyrosine hydroxylase; DA, dopamine; DOPAC, 3,4-dihydroxyphenylacetic acid; 5HT, serotonin; HVA, homovanillic acid; A53T mice, transgenic mice that overexpress human α -synuclein with a PD-associated mutation (A53T)

Model	Behavioral tests	Motor behavior*	Striatal changes*	Survival of SNc neurons*	α -synuclein aggregation*	Study
A53T/aldh1a1 KO mice	Rotarod, open field	↘	↗ α -synuclein	↘	↗	[73]
RAR/RXR KO mice	Rotarod, open field	↘↘	D1R; D2R ↘↘			[76]
VAD rats	Rotarod, motor activity	↘↘	D1R, D2R, TH, DA/DOPAC = ChAT ↘↘			[84]
VAD rats	EEG recording, motor activity, open field	↘	DA and 5HT = DOPAC ↘			[85]
RAR β antagonist in mice	EEG recording	↘ wake periods	DOPAC and HVA = TH, D1R ↘ ratio HVA/DA ↘			[113]
aldh1a1 KO mice			baseline DA ↗ evoked DA ↘ DOPAC ↘	More TH ⁺ neurons		[133]
aldh1a1/aldh2 KO mice	Rotarod, open field, actimetry, gait, Y-maze	↘↘	DA, DOPAC, DOPAC/DA ↘ DOPAL ↗	↘		[134]

*compared to control group.

tors in the striatum is under the control of retinoid receptors [74, 76]. Furthermore, Nurr1 is an orphan nuclear receptor expressed in dopaminergic neurons that dimerize with RXR and has been linked to PD [75]. In mutant mice lacking RAR/RXR or Nurr1, dopaminergic neuron development was altered and associated with motor impairments in adult age [8, 36, 76]. These early observations indicate that retinoid receptors are involved in the development of the nigro-striatal pathway, with consequences on motor function, and it has been postulated that retinoids could be involved in the pathogenesis of PD (Table 2).

Later, important results have been obtained with the ligand of these receptors, namely RA or synthetic derivatives (Table 3). Injection of nanoparticles loaded with RA powerfully reduces dopamine neuron degeneration in the SNc in a mouse model of PD [77]. Similarly, pre-treatment with 9-cis RA reduces midbrain dopamine neuron loss in a rat model of PD [78], as well as neurotoxicity induced by methamphetamine [79]. Furthermore, protection of midbrain dopamine neurons is afforded by exogenous ligand activation of retinoid-related receptors (Nurr1-RXR heterodimers) [80, 81]. In particular, chronic *in vivo* injections of Nurr1:RXR α heterodimer activators in mouse models of PD reduce motor impairments,

increase TH⁺ neurons in the SNc and TH⁺ terminals in the striatum [80]. Vitamin A has also been shown, *in vitro*, to destabilize aggregates of α -synuclein [82], although a contrary result has been reported [83].

Beyond the use of retinoid derivatives, some studies have directly investigated the role of dietary vitamin A in the pathogenesis and pathophysiology of PD (Tables 2 and 3). Vitamin A deficient rats have motor impairments similar to those observed in rat models of PD or in mice lacking retinoid receptors [84]. Another study reported that vitamin A deficiency impaired dopaminergic transmission [85]. Furthermore, the only study that has investigated the impact of dietary vitamin A supplementation showed that vitamin A administration for 4 weeks before the 6-OHDA lesion in the SNc improved movement deficits in 6-OHDA rats [86]. While the behavioural improvement was small, the effect on neuronal survival was not clear. Therefore, more studies are needed with vitamin A supplementation in animal models.

From these preclinical studies, it is clear that vitamin A derivatives and their receptors are involved in the homeostasis of SNc dopaminergic neurons. However, direct involvement of vitamin A metabolism in the pathogenesis of PD has not been established.

Table 3

Summary of the main studies relating the impact of vitamin A pathway enhancement (dietary intake, vitamin A derivatives, or receptors ligands) in Parkinson's disease rodent models

PD model	Supplementation	<i>in vitro/in vivo</i>	Behavioral tests	Motor behavior*	Survival of SNc neurons*	SNc proteins changes*	Striatal proteins changes*	α -synuclein*	Neuroinfla.*	Ox. stress*	Study
Mice MPTP; 6-OHDA; AAV α -synuclein	NURR1:RXR α activator	Both	Rotarod	↗↗	↗↗	TH, DA ↗↗	TH ↗↗				[75]
MPTP mice	RA in striatum	<i>in vivo</i>			↗↗	TH, Pitx3, NURR1 ↗↗	TH, NURR1 (old mice) ↗↗				[77]
6-OHDA rats	9-cis RA (icv)	Both	Rotational test	↗	↗↗	TH ↗↗	DA release & clearance ↗↗				[78]
Metamphet. in rats	9-cis RA (icv)	Both	Motor activity	↗	↗↗		TH ↗				[79]
6-OHDA on mouse primary cultures	RXR agonists	<i>in vitro</i>			↗						[81]
α -synuclein	Vitamin A and β -carotene	<i>in vitro</i>						↘ α -synuclein fibrils			[82]
A53T mice	DHA; RXR over expression; 9-CIS RA	<i>in vitro</i>						↗ oligomers			[83]
6-OHDA rats	vit. A	<i>in vivo</i>	Rotarod	No effect	No effect	TH =	TH =		Iba-1 ↘ GFAP ↗	TNF- α , Il-1 β ↘	[86]

*compared to PD model without supplementation.

VITAMIN A AND DERIVATIVES FOR THE PROTECTION OF DOPAMINERGIC NEURONS IN THE BRAIN

Antioxidant power of vitamin A

Oxidative stress is defined as 'an imbalance between the oxidants and antioxidants in favor of the oxidants, leading to a disruption of redox signaling and control and/or molecular damage' [87, 88] and is a factor for neuronal degeneration in PD [89]. In this context, oxidative stress refers mainly to the formation of free radicals, reactive oxygen species (ROS) that are overproduced by complex I and II of mitochondria in dopaminergic neurons [90]. These free radicals are toxic for cells due to their reactivity and ability to damage biochemical elements. In physiological conditions, the balance between oxidants and antioxidants is maintained by enzymatic antioxidants and non-enzymatic oxidants, and retinoids are part of the antioxidant system [88]. In parkinsonism, accumulation of excessive free radicals in dopaminergic neurons underlies dysfunction of mitochondrial complex I [90, 91]. The origin of this dysfunction is not known and likely to be multiple. Yet, some features of midbrain dopaminergic neurons are thought to favor oxidative stress: (i) their highly branched unmyelinated axons have high bioenergetic demands [92], (ii) pacemaker activity of these neurons (2–10 Hz) produces a constant relatively high level of intracellular calcium level and thus causes a continuous bioenergetic burden for calcium buffering [93, 93], and (iii) metabolism of dopamine creates ROS metabolites that interfere with the function of mitochondrial complex I [90, 93]. Of note, dopamine-3,4-dihydroxyphenylacetaldehyde (DOPAL) is a highly reactive metabolite of dopamine that exacerbates α -synuclein aggregation [94]. These features are not exhaustive and other pro-oxidant factors in dopaminergic neurons, such as metals, do exist (for review, see [89]). Together, multiple sources of free radical production are proposed to explain the vulnerability of midbrain dopaminergic neurons to degenerate in PD.

Retinoids and carotenoids are antioxidants due to their ability to inhibit peroxidation, scavenge free radicals and maintain the balance between oxidants and antioxidants [91, 95–98]. However, studies indicate that high doses of vitamin A supplementation can conversely induce an imbalance between oxidants and pro-oxidants, in favor of oxidants [99–101]. Most

of the data concerning the antioxidant role of vitamin A and associated retinoids have been collected from the heart or liver, but data have rarely explored the antioxidant (or pro-oxidant) role of vitamin A in the neurodegeneration of dopaminergic neurons. Indeed, the role of antioxidants in the progression of PD has been investigated for other vitamins, such as vitamin E or vitamin C [91], but not directly for vitamin A or other retinoids. Therefore, the role of dietary vitamin A on oxidative stress in PD remains to be explicitly studied.

The neuroinflammation action of vitamin A

Neuroinflammation is a common feature of neurodegenerative diseases. Even though the role of neuroinflammation in the pathophysiology of these diseases is established [102], involvement of neuroinflammation in their pathogenesis is not yet demonstrated. However, reducing neuroinflammation is generally a way to curtail disease progression and several pharmacological agents have been tested to address this gap in understanding.

In this context, it is logical to investigate the potential anti-inflammatory role of vitamin A in PD. Indeed, vitamin A is involved in the development, maturation and differentiation of immune organs and cells [103]. Moreover, vitamin A has the propensity to reduce pro-inflammatory factors and enhance anti-inflammatory factors, in the periphery as well as in the brain [104, 105]. Specific to neuroinflammation, *in vitro* studies demonstrate that application of RA to astrocytes or microglial cells reduces inflammatory responses induced by lipopolysaccharide (LPS), a bacterial endotoxin [106, 107]. Related to PD, a study clearly demonstrated that pharmacological stimulation of RAR *in vitro* and *in vivo* reduces neurodegeneration of midbrain dopaminergic neurons induced by an immune challenge. A more recent study observed a protective effect of dietary vitamin A in a rat model of PD by reducing pro-inflammatory factors (TNF- α , IL-1 β , Iba-1), which was accompanied with an increase in GFAP staining, suggesting increased astrocytic reactivity [86].

The role of retinoids in dampening neuroinflammation appears an important mechanism to impede neurodegeneration in PD. However, given the limited number of preclinical studies, more research is needed to better understand the underlying mechanisms.

In PD, the classical scheme of neuroinflammation indicates that pro-inflammatory factors are produced

by activated microglia and astrocytes. In addition, pro-inflammatory factors released by peripheral macrophages can penetrate brain tissue because the blood-brain barrier becomes leaky [108]. It is likely that these pro-inflammatory factors contribute to oxidative stress in dopaminergic neurons, which further activate neuroinflammatory processes [60, 102, 108, 109].

Related to neuroinflammation and oxidative stress, studies exploring the role of vitamin A metabolism in the control of circadian rhythms and sleep deserve attention. A main prodromal symptom observed in PD is sleep disorder [110] and it is also observed in rat models of PD [111]. These sleep disorders are mainly a lack of REM sleep, but can also exhibit other forms of sleep deficits. Retinoid signaling is involved in sleep wave rhythms, as demonstrated in mice with RAR KO mice [112], vitamin A deficiency [85] or pharmacological inhibitors [113]. Importantly, these studies also highlight that reduced vitamin A function induced changes in sleep wave rhythms that were associated with alterations in striato-nigral dopaminergic transmission [85, 113].

In addition, circadian rhythms are also altered in PD, which could contribute (even initiate) to the vicious cycle between neuroinflammation and oxidative stress [114]. Multiple evidence shows that vitamin A, through RA and RAR/RXR receptors, controls the gene of the circadian oscillator [115].

Vitamin A and neurogenesis

Under physiological conditions, neurogenesis happens in restricted regions of the adult mammalian brain such as the subgranular zone of hippocampal dentate gyrus and the subventricular zone (SVZ) of lateral ventricles, two areas that neural stem cells develop [116]. Neurogenic niches are endogenous sources for new neurons that can be used for brain repair strategies [117]. It has been demonstrated that RA is necessary for adult hippocampal neurogenesis [3, 118]. Indeed, nutritional vitamin A deficiency induced spatial memory deficits and adult hippocampal neurogenesis alterations, that are corrected by RA treatment or vitamin A supplementation in rodents [119, 120].

In rodent models of PD, due to the proximity of the SVZ with the striatum (and thus dopaminergic afferences), stem cells from this area can be recruited following a lesion. Some studies have established the potential therapeutic benefit of neurotrophic factors and proposed them as suitable

candidates for stimulating neurogenesis after induction of neurodegeneration [121]. Among these neurotrophic factors, CDNF (cerebral dopamine neurotrophic factor) is the most promising treatment for PD due to its selective effects on dopaminergic neurons. Thus, it has been shown that intra-SVZ administration of CDNF enhances the proliferation and migration of neural stem cells toward the 6-hydroxydopamine (6-OHDA)-lesioned striatum accompanied by improvement of motor dysfunctions in parkinsonian rats [122].

In this context, vitamin A and RA are also promising molecules to enhance the generation and long-term survival of SVZ-derived neurons after PD lesions. Indeed, RA is a potent mitogen for SVZ neuroblasts, and is required for their migration to the olfactory bulb [123]. More recently, the manipulation of endogenous stem cell populations from the SVZ created an opportunity to induce neurogenesis and influence brain regenerative capacities in the adult brain. Herein, new approaches demonstrated the ability of RA loaded-nanoparticles to induce neurogenesis exclusively after being internalized by SVZ stem cells both *in vivo* and *in vitro* [124–126]. Similarly, combined RA treatment with environmental enrichment enhanced the generation and long-term survival of SVZ-derived striatal neurons after stroke [127].

These data indicate that recruiting endogenous SVZ neural stem cells toward striatal regions exhibiting retrograde degeneration of dopaminergic afferents in PD brains by stimulating retinoid signaling could be a way to slow down the neurodegeneration of nigrostriatal dopaminergic neurons, and to allow functional recovery. Retinoids could also be used as a molecular factor to improve the durability of cell grafts that are currently developed to promote cell repair [4, 128]. In this later case, retinoid signaling could be envisaged as a treatment strategy.

Pivotal role for ALDH1A1

Beside the potential role for vitamin A to control neuroinflammation and oxidative stress, vitamin A's action may also be specific to dopaminergic neurons by controlling the retinaldehyde dehydrogenase 1 (ALDH1A1) enzyme.

As mentioned above, RA in the adult brain controls homeostasis of dopaminergic neurons and dopaminergic transmission by controlling the expression of tyrosine hydroxylase, D2-like receptors, and the RA synthetic enzyme, ALDH1A1 [3]. ALDH1A1 is a

retinaldehyde dehydrogenase enzyme that is part of the aldehyde dehydrogenase superfamily of enzymes whose main function is to synthesize RA from retinal [129] (Figs. 1 and 2). These enzymes form a vast family that has multiple functions, such as acetaldehyde detoxification [129]. It exists as 3 isotypes of RALDH: RALDH1 (ALDH1A1), RALDH2 and RALDH3. ALDH1A1 is coded by the gene *Aldh1a1* but the enzyme has multiple names in the literature: Ald1a1, RALDH1, ALDH1, Ahd2. Excluding dopaminergic neurons, RALDH2 is the main isotype expressed in the brain [130]. Dopaminergic neurons are unique in the brain because they are the only ones to express the ALDH1A1 isotype in the adult brain [21]. Specifically, only a subset of midbrain dopaminergic neurons in the SNc and ventral tegmental area express this enzyme [131], making ALDH1A1 a marker for this subpopulation. This dopaminergic subpopulation is interconnected with striatal neurons and mice lacking these neurons display motor impairments [132]. Mice lacking the ALDH1A1 enzyme also display motor impairments and alterations in dopamine release [132–134], consistent with associated motor deficits and reduced levels of ALDH1A1 reported from clinical studies [71–73] (Table 1). Remarkably, ALDH1A1^{-/-} mice also display a lack of mu opioid receptors MOR1 expression in striosomes compartments [24, 25].

In addition to these preclinical data brought by ALDH1A1^{-/-} mice, it appears that ALDH1A1 labelled neurons correspond to the dopaminergic subpopulation that is the most vulnerable to degeneration in PD [135, 136]. Specifically, data obtained from humans and rodent models of PD indicate that neurons that degenerate early are those that express the gene *Aldh1a1* coding for ALDH1A1 [73, 131, 137].

This raises the question of why ALDH1A1 expressing neurons are more vulnerable in PD? Is ALDH1A1 a marker for neuroprotection or neurodegeneration? ALDH1A1 is neuroprotective, because these dopaminergic neurons stop expressing ALDH1A1 before degenerating [71, 73, 137]. Importantly, these observations are not only based on animal studies, since a reduction of ALDH1A1⁺ neurons has also been observed in the SNc of patients with PD [137]. Therefore, the scenario envisaged is that ALDH1A1⁺ neurons stop expressing ALDH1A1 (for unknown reasons, discussed later), which triggers their degeneration. In support with this, accumulating data show that ALDH1A1, beyond synthesizing RA, plays an essential detoxifying role. Indeed, ALDH1A1 has the ability to degrade

DOPAL, a highly oxidizing dopamine metabolite that induces oxidative stress and aggregation of α -synuclein proteins [94]. Thus, DOPAL could be a major contributor to dopaminergic neuron loss in PD, whereas ALDH1A1 would preserve dopamine neurons by degrading DOPAL [73, 94, 137, 138]. This potential mechanism is consistent with the 'catecholaldehyde hypothesis' [139–141].

In support of the catecholaldehyde hypothesis, unintentional over-inhibition of ALDH1A in humans has produced patients with parkinsonism symptoms, following disulfiram (used as Antabuse® for the treatment of alcoholism) intoxication [142, 143]. Furthermore, the toxicity of some pesticides, such as benomyl or rotenone, is due to the inhibition of aldehyde dehydrogenase enzymes [138, 144–146]. However, in these articles, the ALDHs that are studied are mostly mitochondrial ones and ALDH1A1 is not investigated. Still, pesticide exposure is a known cause of PD and these studies highlight the crucial role of ALDH enzymes in the survival of dopaminergic neurons.

Beyond this pharmacological/exogenous inhibition of ALDH1A1, we do not know why and how dopaminergic neurons stop expressing ALDH1A1 in parkinsonian patients. Nevertheless, ALDH1A1 expression is controlled by retinoid signaling. Knowing that ALDH1A1 expression may be an important prodromal biomarker for PD [73, 137], we propose the novel hypothesis that vitamin A metabolism is involved in PD pathogenesis. Indeed, ALDH1A1 expression is modulated by RA levels. More precisely, RA stimulates the expression of PITX3, a transcription factor that controls expression of ALDH1A1 [77, 147, 148]. Mice deficient for PITX3 exhibit strong dopamine loss in the SNc [149–151] and exhibit motor symptoms that are similar to animals with neurotoxic lesions [152, 153], which can be reversed by RA administration [147].

In support of our hypothesis that vitamin A metabolism is involved in PD pathogenesis, we recently showed that aging in rats is accompanied by a decreased bioavailability of RA in the brain [18]. Since aging is the greatest risk factor for PD, the collapse of retinoid signaling in the brain could be a contributing factor in a vicious cycle leading to degeneration of ALDH1A1 + dopaminergic neurons in the SNc. This proposed mechanism is presented in Fig. 3. Multiple, disparate evidence supports our hypothesis. However, the real association between vitamin A status, RA bioavailability and dopaminergic neuron loss in PD remains to be demonstrated.




Fig. 3. Model of the dual role played by ALDH1A1 in the nigro-striatal pathway. ALDH1A1 is involved in the metabolic pathway of RA because it synthesizes RA from retinal. In parallel, ALDH1A1 is involved in catabolic pathway of dopamine because it degrades DOPAL to DOPAC. Considering that RA controls the expression of ALDH1A1 through PITX3, the model proposes that ALDH1A1 expression is controlled by vitamin A bioavailability. SNc, substantia nigra pars compacta; SNr, substantia nigra pars reticulata.

This proposed mechanism also raises further questions about the subpopulation of dopaminergic neurons that do not express ALDH1A1. Do dopamine neurons without ALDH1A1 produce DOPAL? If yes, what are the enzymes responsible for DOPAL degradation in these neurons? What enzyme isoforms are expressed? Furthermore, data indicate that ALDH1A1 determines the GABAergic nature of neurons from the SNc [31, 154], which is in contrast (but not necessarily in contradiction) with data indicating that RA is involved in the development of dopaminergic pathways [8, 34, 36, 74, 75].

In conclusion, data support the theory that ALDH1A1 is a pivotal enzyme for understanding the role of vitamin A in the neurodegeneration of dopaminergic neurons, but further investigations are needed to confirm and strengthen these data.

PERIPHERAL ACTIONS OF VITAMIN A IN PD

The enteric nervous system (ENS)

The ENS is the autonomous nervous system of the gastrointestinal (GI) tract and it is connected to autonomous and central nervous systems by the vagus nerve. ENS neurons are remarkable in their diversity, and they are highly organized [155]. In particular, the ENS is rich in dopaminergic neurons.

GI dysfunctions, such as constipation or slowed gastric emptying, are symptoms frequently observed in PD, which can appear years before motor symptoms [156, 157]. These prodromal symptoms are thought to be related to altered ENS function. Since Lewy bodies and alpha-synuclein aggregates have

been found in the ENS and dorsal motor nucleus of the vagus nerve in PD patients, the disease may originate from enteric alpha-synucleopathy and reach midbrain dopamine neurons through the dorsal motor nucleus of the vagus nerve [158]. These mechanisms may involve dopaminergic neurons and/or epithelial cells that produce dopamine, for the same reasons outlined earlier for CNS dopaminergic neurons. Since proposing this hypothesis in 2003, numerous preclinical and clinical studies have been conducted [159–164]. In rodents, alpha-synuclein transport from the GI to the midbrain and associated motor impairments have recently been demonstrated [157, 164, 165]. In humans, some evidence highlights the link between GI dysfunction, synucleopathy in the ENS and severity of PD symptoms [157, 164]. However, this evidence has also been rebutted by other studies [166–169]. Therefore, the precise role of ENS synucleopathy in the pathophysiology of PD is still under debate.

In this context, we wonder whether vitamin A can protect ENS neurons via similar mechanisms proposed for midbrain dopaminergic neurons. RA plays a critical role in the development and differentiation of ENS neurons [170]. As for central dopaminergic neurons, we could extrapolate that factors involved in the development of ENS neurons are also critical for their survival at adult age. In addition, vitamin A deficiency in rats reduces the number of cholinergic and nitrergic neurons in the ENS, which is associated with altered colon motility [171]. Interestingly, reduced colon contractility has also been found in RALDH KO mice [172]. However, the impact of vitamin A deficiency on ENS dopaminergic neurons does not appear to have been investigated. From these data, we can *postulate* that vitamin A metabolism is important for survival of ENS dopaminergic neurons, and may protect from their degeneration, but more research is needed.

The gut microbiome

Gut microbiota composition has recently become a focus for PD researchers [173–176]. A key study in 2016 demonstrated that human gut microbiota from Parkinson's patients worsens motor symptoms in a mouse model of PD [177]. Furthermore, a two-year follow-up study revealed that low counts of specific bacteria such as *Bifidobacterium* (Actinobacterium phylum) and *B. fragilis* (Bacteroidetes phylum) at the beginning of the study was associated with much worse symptoms of the disease two years later [178].

However, there is still a lack of preclinical and clinical data explaining the links between impoverished gut microbiome and progression of PD [179, 180].

Vitamin A deficiency may alter gut microbiota. In mice, vitamin A deficiency reduces bacteria from the Bacteroidetes phylum (to whom *B. fragilis* belongs to), which altered energy homeostasis of the animal overall, and resulted in glucose and insulin intolerance [181, 182]. Young rats from vitamin A deficient mothers also displayed a dysbiosis of colonic mucosal microbiota, in particular with reduced members of the Bacteroidetes phylum [183]. Another study revealed that retinol, but not other retinoids such as RA or beta-carotene, inhibits growth of *B. vulgatus* [184]. As a consequence, vitamin A deficiency in mice increases the growth of *B. vulgatus*, but the consequence of this imbalance was not evaluated. Conversely, RA is needed for *Bifidobacterium* growth [185].

Thus, two observations may converge; vitamin A likely modulates gut microbiota composition, and PD is associated with specific microbiota modifications. However, the lack of studies about microbiota related to PD, as well as related to vitamin A is too significant to draw any hypothesis about the mechanisms linking vitamin A dysfunction to microbiota changes in PD.

Link between vitamin A and other hormones through RXR dimerization

Finally, vitamin A may influence the development of PD pathophysiology indirectly via hormonal systems. Indeed, RXRs have a particular role because they are dimer partners (sometimes obligatory) for multiple nuclear receptors, such as thyroid receptor (TR), Nurr1, Nur77 or vitamin D receptor (VDR). Thus, retinoids (9CDHRA in particular) through their binding to RXRs are at a crossroad between multiple hormonal pathways, and vitamin A is considered a hormone [186]. Furthermore, expression of RXRs themselves are controlled by retinoids, therefore, vitamin A status can regulate the function of RXRs [187, 188]. In the context of PD, the bioavailability of vitamin A can thus have an impact on hormonal systems, which affect symptom progression, health and wellbeing, but this interaction could be easily overlooked.

The case of thyroid hormones

Thyroid receptors (TRs) are nuclear hormone receptors for which heterodimerization with RXR is obligatory. More precisely, RXR modulates the

action of TR with the outcome dependent on the ligands of both receptors [17]. Vitamin A deficiency is associated with altered thyroid function, while hypothyroidism is associated to decreased RAR in humans [189]. RA modulates the expression of proteins involved in thyrocyte function and can stimulate their differentiation [189]. Vitamin A and thyroid metabolisms are thus intricately linked even though underlying mechanisms are not fully understood. With aging, levels of both RARs and TRs are decreased as shown in rats and humans [53, 190–193]. However, studies diverge on RA's ability to restore TR mRNA levels in aged rats [53, 190–192]. Furthermore, thyroid hormones stimulate metabolism and can increase the production of ROS by mitochondria [194]. Thus, thyroid dysfunction may be involved in idiopathic PD pathophysiology through oxidative stress. In a preclinical study, increased levels of T3 have been found in 6-OHDA lesioned rats suggesting that central dopaminergic denervation can act on thyroid levels at the periphery, possibly through dysfunction

of hypothalamic-pituitary axis [195]. In patients, thyroid hormone function has been poorly documented so far and results are mitigated. One study indicated that subclinical hypothyroidism was more frequent in PD patients [196]. Another study indicated an association between the severity of motor symptoms and level of free circulating thyroid hormones (fT3) [197]. However, hypothyroidism did not differ significantly between parkinsonian patients and controls in a third study [198]. Therefore, due to the lack of studies the interaction between vitamin A and thyroid function in the pathophysiology of PD has not been established. Further investigations should be conducted, particularly in patients with idiopathic origin, who are over 60 years of age at the time of diagnosis, an age where the thyroid function can be impaired by aging processes.

The case of vitamin D

Vitamin D is renowned for its roles in immune function and maintaining bone density (through the control of calcium serum levels). As for TRs, expres-




Fig. 4. Proposed roles and mechanisms of vitamin A metabolism in the pathophysiology of Parkinson's disease. Inset with α-synuclein aggregates from [110].

sion of vitamin D nuclear receptors (VDRs) is controlled by RA, and RXR is an obligatory heterodimer for VDRs [199]. A correlation between vitamin D levels and PD has been observed, but vitamin D supplementation has not significantly reduced disease status [200–202]. The SNc and the striatum are two structures highly enriched for VDR and 1 α -hydroxylase, the enzyme that synthesizes the active form of vitamin D [203]. Pre-clinical data show that vitamin D protects dopaminergic neurons from oxidative stress [199, 203–206]. *In vitro*, low (but not high) doses of the active form of vitamin D protects dopaminergic neurons (primary mesencephalic cultures) from toxins [204]. *In vivo*, treatment with the active form of vitamin D 1 week before 6-OHDA lesion in rats increases motor activity, dopamine content in the striatum and number of TH⁺ neurons [205, 206]. These studies suggest that the positive effect of vitamin D is mediated by its antioxidant effect. Thus, beyond the interactions that may exist between vitamins A and D in the immune system, these two vitamins may act synergistically in the brain. However, the combined action of vitamins D and A on the survival of dopaminergic neurons in PD, notably through their partner nuclear receptors, has not been investigated. Yet, this constitutes an interesting avenue with easily translatable therapeutic outcome.

CONCLUSION

This review explored multiple mechanisms that vitamin A metabolism may contribute to PD pathogenesis and PD pathophysiology. Indeed, altered vitamin A metabolism and bioavailability is likely to contribute to oxidative stress, neuroinflammation, dopaminergic cell death, disturbance in biological rhythms and endocrine homeostasis (Fig. 4). Thus, vitamin A, as a nutritional factor may be at the crossroad of multiple environmental and genetic factors of PD. Still, the central underlying role that decreased vitamin A metabolism may have in PD has been largely ignored. This may be due to the lack of adapted methods for quantifying vitamin A metabolism and bioavailability in humans.

From our overview, we propose that vitamin A metabolism may be involved in the pathogenesis and pathophysiology of PD in multiple ways (Fig. 4). Of note, other possible ways have not been discussed here, such as the potential role of retinoids in autophagy. At this stage, the most promising way

by which vitamin A metabolism may influence PD pathogenesis and treatment is through the impact of vitamin A on ALDH1A1 expression and neuroinflammation. A first step is to understand the role of ALDH1A1 in controlling dopaminergic cell survival, within the schema of the catecholaldehyde hypothesis. However, other mechanisms of vitamin A metabolism are likely relevant including oxidative stress, neurogenesis in the SVZ, ENS function and microbiota, thyroid hormone and vitamin D function, but more data are needed to fully understand the role of individual or combined mechanisms.

Finally, it is important to keep in mind that pharmacological treatments with retinoids have been studied for a long time and these drugs have to be used in precise conditions to preclude side effects [7]. In this framework, targeting vitamin A, and not RA and its receptors, may be a more conservative strategy since vitamin A supplementation increases bioavailability for the brain, and is better managed by organisms than derivatives, with fewer side effects. Together, a better understanding of the role played by vitamin A metabolism in PD could open the way for new approaches to dampen symptoms and improve health and wellbeing for PD patients.

ACKNOWLEDGMENTS

This work was supported by funding from France Parkinson, Health Research Council (17–284) and Brain Research New Zealand

CONFLICT OF INTEREST

The authors have no conflict of interest to report.

REFERENCES

- [1] Maden M (2007) Retinoic acid in the development, regeneration and maintenance of the nervous system. *Nat Rev Neurosci* **8**, 755-765.
- [2] Goodman AB (2006) Retinoid receptors, transporters, and metabolizers as therapeutic targets in late onset Alzheimer disease. *J Cell Physiol* **209**, 598-603.
- [3] Lane MA, Bailey SJ (2005) Role of retinoid signalling in the adult brain. *Prog Neurobiol* **75**, 275-293.
- [4] Puttagunta R, Di Giovanni S (2011) Retinoic acid signaling in axonal regeneration. *Front Mol Neurosci* **4**, 59.
- [5] Shearer KD, Stoney PN, Morgan PJ, McCaffery PJ (2012) A vitamin for the brain. *Trends Neurosci* **35**, 733-741.
- [6] Sodhi RK, Singh N (2014) Retinoids as potential targets for Alzheimer's disease. *Pharmacol Biochem Behav* **120**, 117-123.
- [7] Clark JN, Whiting A, McCaffery P (2020) Retinoic acid receptor-targeted drugs in neurodegenerative disease. *Expert Opin Drug Metab Toxicol* **16**, 1097-1108.

- [8] Eichele G (1997) Retinoids: From hindbrain patterning to Parkinson disease. *Trends Genet* **13**, 343-345.
- [9] Asson-Batres MA, Rochette-Egly C, eds. (2016) *The Biochemistry of Retinoid Signaling II: The Physiology of Vitamin A - Uptake, Transport, Metabolism and Signaling*, Springer Netherlands, Dordrecht.
- [10] Sauvant P, Feart C, Sauvant J, Pallet V (2015) Vitamin A and cognitive impairment. In *Diet and Nutrition in Dementia and Cognitive Decline*, Elsevier, pp. 623-635.
- [11] Krężel W, Kastner P, Chambon P (1999) Differential expression of retinoid receptors in the adult mouse central nervous system. *Neuroscience* **89**, 1291-1300.
- [12] Krężel W, Rühl R, de Lera AR (2019) Alternative retinoid X receptor (RXR) ligands. *Mol Cell Endocrinol* **491**, 110436.
- [13] Rühl R, Krężel W, de Lera AR (2018) 9-Cis-13,14-dihydroretinoic acid, a new endogenous mammalian ligand of retinoid X receptor and the active ligand of a potential new vitamin A category: Vitamin A5. *Nutr Rev* **76**, 929-941.
- [14] Kojetin DJ, Matta-Camacho E, Hughes TS, Srinivasan S, Nwachukwu JC, Cavett V, Nowak J, Chalmers MJ, Marciano DP, Kamenecka TM, Shulman AI, Rance M, Griffin PR, Bruning JB, Nettles KW (2015) Structural mechanism for signal transduction in RXR nuclear receptor heterodimers. *Nat Commun* **6**, 8013.
- [15] Niewiadomska-Cimicka A, Krzyżosiak A, Ye T, Podleśny-Drabiniok A, Dembélé D, Dollé P, Krężel W (2017) Genome-wide analysis of RAR β transcriptional targets in mouse striatum links retinoic acid signaling with Huntington's disease and other neurodegenerative disorders. *Mol Neurobiol* **54**, 3859-3878.
- [16] Blomhoff R, Blomhoff HK (2006) Overview of retinoid metabolism and function. *J Neurobiol* **66**, 606-630.
- [17] Li D, Yamada T, Wang F, Vulin AI, Samuels HH (2004) Novel roles of retinoid X receptor (RXR) and RXR ligand in dynamically modulating the activity of the thyroid hormone receptor/RXR heterodimer. *J Biol Chem* **279**, 7427-7437.
- [18] Dumetz F, Buré C, Alfons S, Bonneu M, Richard E, Touyarot K, Marie A, Schmitter J-M, Bosch-Bouju C, Pallet V (2020) Normalization of hippocampal retinoic acid level corrects age-related memory deficits in rats. *Neurobiol Aging* **85**, 1-10.
- [19] Noy N (2016) Vitamin A transport and cell signaling by the retinol-binding protein receptor STRA6. In *The Biochemistry of Retinoid Signaling II: The Physiology of Vitamin A - Uptake, Transport, Metabolism and Signaling*, Asson-Batres MA, Rochette-Egly C, eds. Springer Netherlands, Dordrecht, pp. 77-93.
- [20] McCaffery P, Zhang J, Crandall JE (2006) Retinoic acid signaling and function in the adult hippocampus. *J Neurobiol* **66**, 780-791.
- [21] McCaffery P, Drager UC (1994) High levels of a retinoic acid-generating dehydrogenase in the meso-telencephalic dopamine system. *Proc Natl Acad Sci U S A* **91**, 7772-7776.
- [22] Zetterström RH, Lindqvist E, Mata de Urquiza A, Tomac A, Eriksson U, Perlmann T, Olson L (1999) Role of retinoids in the CNS: Differential expression of retinoid binding proteins and receptors and evidence for presence of retinoic acid. *Eur J Neurosci* **11**, 407-416.
- [23] Ferreira R, Napoli J, Enver T, Bernardino L, Ferreira L (2020) Advances and challenges in retinoid delivery systems in regenerative and therapeutic medicine. *Nat Commun* **11**, 4265.
- [24] Pan J, Yu J, Sun L, Xie C, Chang L, Wu J, Hawes S, Saez-Atienzar S, Zheng W, Kung J, Ding J, Le W, Chen S, Cai H (2019) ALDH1A1 regulates postsynaptic μ -opioid receptor expression in dorsal striatal projection neurons and mitigates dyskinesia through transsynaptic retinoic acid signaling. *Sci Rep* **9**, 3602.
- [25] Sgobio C, Wu J, Zheng W, Chen X, Pan J, Salinas AG, Davis MI, Lovinger DM, Cai H (2017) Aldehyde dehydrogenase 1-positive nigrostriatal dopaminergic fibers exhibit distinct projection pattern and dopamine release dynamics at mouse dorsal striatum. *Sci Rep* **7**, 5283.
- [26] Davidi D, Schechter M, Elhadi SA, Matatov A, Nathanson L, Sharon R (2020) α -Synuclein Translocates to the Nucleus to Activate Retinoic-Acid-Dependent Gene Transcription. *iScience* **23**, 100910.
- [27] Bozelli JC, Kamski-Hennekam E, Melacini G, Epand RM (2021) α -Synuclein and neuronal membranes: Conformational flexibilities in health and disease. *Chem Phys Lipids* **235**, 105034.
- [28] Colom-Cadena M, Pegueroles J, Herrmann AG, Henstridge CM, Muñoz L, Querol-Vilaseca M, Martín-Paniello CS, Luque-Cabecerans J, Clarimon J, Belbin O, Núñez-Llaves R, Blesa R, Smith C, McKenzie C-A, Frosch MP, Roe A, Fortea J, Andilla J, Loza-Alvarez P, Gelpi E, Hyman BT, Spires-Jones TL, Lleó A (2017) Synaptic phosphorylated α -synuclein in dementia with Lewy bodies. *Brain* **140**, 3204-3214.
- [29] Rataj-Baniowska M, Niewiadomska-Cimicka A, Paschaki M, Szyszka-Niagolov M, Carramolino L, Torres M, Dollé P, Krężel W (2015) Retinoic acid receptor β controls development of striatonigral projection neurons through FGF-dependent and Meis1-dependent mechanisms. *J Neurosci* **35**, 14467-14475.
- [30] Pedrini S, Bogush A, Ehrlich ME (2008) Phosphatidylinositol 3-kinase and protein kinase C zeta mediate retinoic acid induction of DARPP-32 in medium size spiny neurons *in vitro*. *J Neurochem* **106**, 917-924.
- [31] Chatzi C, Brade T, Duester G (2011) Retinoic acid functions as a key GABAergic differentiation signal in the basal ganglia. *PLoS Biol* **9**, e1000609.
- [32] Liao W-L, Tsai H-C, Wang H-F, Chang J, Lu K-M, Wu H-L, Lee Y-C, Tsai T-F, Takahashi H, Wagner M, Ghyselinck NB, Chambon P, Liu F-C (2008) Modular patterning of structure and function of the striatum by retinoid receptor signaling. *Proc Natl Acad Sci U S A* **105**, 6765-6770.
- [33] Cardoso T, Lévesque M (2020) Toward generating subtype-specific mesencephalic dopaminergic neurons *in vitro*. *Front Cell Dev Biol* **8**, 443.
- [34] Hegarty SV, Sullivan AM, O'Keefe GW (2013) Midbrain dopaminergic neurons: A review of the molecular circuitry that regulates their development. *Dev Biol* **379**, 123-138.
- [35] Wallén Å, Zetterström RH, Solomin L, Arvidsson M, Olson L, Perlmann T (1999) Fate of mesencephalic AHD2-expressing dopamine progenitor cells in Nurr1 mutant mice. *Exp Cell Res* **253**, 737-746.
- [36] Zetterström RH, Solomin L, Jansson L, Hoffer BJ, Olson L, Perlmann T (1997) Dopamine neuron agenesis in Nurr1-deficient mice. *Science* **276**, 248-250.
- [37] Podleśny-Drabiniok A, Sobska J, de Lera AR, Golembiowska K, Kamińska K, Dollé P, Cebrat M, Krężel W (2017) Distinct retinoic acid receptor (RAR)

- isotypes control differentiation of embryonal carcinoma cells to dopaminergic or striatopallidal medium spiny neurons. *Sci Rep* **7**, 13671.
- [38] Chen L, Lau AG, Sarti F (2014) Synaptic retinoic acid signaling and homeostatic synaptic plasticity. *Neuropharmacology* **78**, 3-12.
- [39] Chiang MY, Misner D, Kempermann G, Schikorski T, Giguère V, Sucov HM, Gage FH, Stevens CF, Evans RM (1998) An essential role for retinoid receptors RARbeta and RXRgamma in long-term potentiation and depression. *Neuron* **21**, 1353-1361.
- [40] Misner DL, Jacobs S, Shimizu Y, de Urquiza AM, Solomin L, Perlmann T, De Luca LM, Stevens CF, Evans RM (2001) Vitamin A deprivation results in reversible loss of hippocampal long-term synaptic plasticity. *Proc Natl Acad Sci U S A* **98**, 11714-11719.
- [41] Nomoto M, Takeda Y, Uchida S, Mitsuda K, Enomoto H, Saito K, Choi T, Watabe AM, Kobayashi S, Masushige S, Manabe T, Kida S (2012) Dysfunction of the RAR/RXR signaling pathway in the forebrain impairs hippocampal memory and synaptic plasticity. *Mol Brain* **5**, 8.
- [42] Dumetz F, Giniès R, Buré C, Marie A, Alfos S, Pallet V, Bosch-Bouju C (2020) Neuronal morphology and synaptic plasticity in the hippocampus of vitamin A deficient rats. *Nutr Neurosci*, doi: 10.1080/1028415X.2020.1809877
- [43] Aoto J, Nam CI, Poon MM, Ting P, Chen L (2008) Synaptic signaling by all-trans retinoic acid in homeostatic synaptic plasticity. *Neuron* **60**, 308-320.
- [44] Wang H-L, Zhang Z, Hintze M, Chen L (2011) Decrease in calcium concentration triggers neuronal retinoic acid synthesis during homeostatic synaptic plasticity. *J Neurosci* **31**, 17764-17771.
- [45] Sarti F, Zhang Z, Schroeder J, Chen L (2013) Rapid suppression of inhibitory synaptic transmission by retinoic acid. *J Neurosci* **33**, 11440-11450.
- [46] Arendt KL, Zhang Y, Jurado S, Malenka RC, Südhof TC, Chen L (2015) Retinoic acid and LTP recruit postsynaptic AMPA receptors using distinct SNARE-dependent mechanisms. *Neuron* **86**, 442-456.
- [47] Arendt KL, Zhang Z, Ganesan S, Hintze M, Shin MM, Tang Y, Cho A, Graef IA, Chen L (2015) Calcineurin mediates homeostatic synaptic plasticity by regulating retinoic acid synthesis. *Proc Natl Acad Sci U S A* **112**, E5744-E5752.
- [48] Hsu Y-T, Li J, Wu D, Südhof TC, Chen L (2019) Synaptic retinoic acid receptor signaling mediates mTOR-dependent metaplasticity that controls hippocampal learning. *Proc Natl Acad Sci U S A* **116**, 7113-7122.
- [49] Yee AX, Chen L (2016) Differential regulation of spontaneous and evoked inhibitory synaptic transmission in somatosensory cortex by retinoic acid. *Synapse* **70**, 445-452.
- [50] Maghsoodi B, Poon MM, Nam CI, Aoto J, Ting P, Chen L (2008) Retinoic acid regulates RARalpha-mediated control of translation in dendritic RNA granules during homeostatic synaptic plasticity. *Proc Natl Acad Sci U S A* **105**, 16015-16020.
- [51] Loweth JA, Reimers JM, Caccamise A, Stefanik MT, Woo KKY, Chauhan NM, Werner CT, Wolf ME (2019) mGlu1 tonically regulates levels of calcium-permeable AMPA receptors in cultured nucleus accumbens neurons through retinoic acid signaling and protein translation. *Eur J Neurosci* **50**, 2590-2601.
- [52] Féart C, Siewe A, Samieri C, Peuchant E, Helmer C, Alfos S, Pallet V, Barberger-Gateau P (2010) Plasma retinol and association with socio-demographic and dietary characteristics of free-living older persons: The Bordeaux sample of the three-city study. *Int J Vitam Nutr Res* **80**, 32-44.
- [53] Féart C, Mingaud F, Enderlin V, Husson M, Alfos S, Higuieret P, Pallet V (2005) Differential effect of retinoic acid and triiodothyronine on the age-related hypoxpression of neurogranin in rat. *Neurobiol Aging* **26**, 729-738.
- [54] Bonhomme D, Pallet V, Dominguez G, Servant L, Henkous N, Lafenêtre P, Higuieret P, Béracochéa D, Touyarot K (2014) Retinoic acid modulates intrahippocampal levels of corticosterone in middle-aged mice: Consequences on hippocampal plasticity and contextual memory. *Front Aging Neurosci* **6**, 6.
- [55] Goodman DS, Huang HS, Shiratori T (1965) Tissue distribution and metabolism of newly absorbed vitamin A in the rat. *J Lipid Res* **6**, 390-396.
- [56] Harrison EH (2005) Mechanisms of digestion and absorption of dietary vitamin A. *Annu Rev Nutr* **25**, 87-103.
- [57] Hodges JK, Tan L, Green MH, Ross AC (2016) Vitamin A supplementation increases the uptake of chylomicron retinyl esters into the brain of neonatal rats raised under vitamin A-marginal conditions. *J Nutr* **146**, 1677-1683.
- [58] Anderson C, Checkoway H, Franklin GM, Beresford S, Smith-Weller T, Swanson PD (1999) Dietary factors in Parkinson's disease: The role of food groups and specific foods. *Mov Disord* **14**, 21-27.
- [59] Boulos C, Yaghi N, El Hayeck R, Heraoui GN, Fakhoury-Sayegh N (2019) Nutritional risk factors, microbiota and Parkinson's disease: What is the current evidence? *Nutrients* **11**, 1896.
- [60] Ciulla M, Marinelli L, Cacciato I, Stefano AD (2019) Role of dietary supplements in the management of Parkinson's disease. *Biomolecules* **9**, 271.
- [61] Erro R, Brigo F, Tamburin S, Zamboni M, Antonini A, Tinazzi M (2018) Nutritional habits, risk, and progression of Parkinson disease. *J Neurol* **265**, 12-23.
- [62] Jiménez-Jiménez FJ, Molina JA, Fernández-Calle P, Vázquez A, Cabrera-Valdivia F, Catalán MJ, García-Albea E, Bermejo F, Codoceo R (1993) Serum levels of beta-carotene and other carotenoids in Parkinson's disease. *Neurosci Lett* **157**, 103-106.
- [63] Johnson CC, Gorell JM, Rybicki BA, Sanders K, Peterson EL (1999) Adult nutrient intake as a risk factor for Parkinson's disease. *Int J Epidemiol* **28**, 1102-1109.
- [64] Paganini-Hill A (2001) Risk factors for parkinson's disease: The leisure world cohort study. *Neuroepidemiology* **20**, 118-124.
- [65] Takeda A, Nyssen OP, Syed A, Jansen E, Bueno-de-Mesquita B, Gallo V (2014) Vitamin A and carotenoids and the risk of Parkinson's disease: A systematic review and meta-analysis. *Neuroepidemiology* **42**, 25-38.
- [66] Zhang SM, Hernán MA, Chen H, Spiegelman D, Willett WC, Ascherio A (2002) Intakes of vitamins E and C, carotenoids, vitamin supplements, and PD risk. *Neurology* **59**, 1161-1169.
- [67] Kim JH, Hwang J, Shim E, Chung E-J, Jang SH, Koh S-B (2017) Association of serum carotenoid, retinol, and tocopherol concentrations with the progression of Parkinson's Disease. *Nutr Res Pract* **11**, 114.
- [68] Yang F, Wolk A, Håkansson N, Pedersen NL, Wirdefeldt K (2017) Dietary antioxidants and risk of Parkinson's disease in two population-based cohorts. *Mov Disord* **32**, 1631-1636.

- [69] Royal W, Gartner S, Gajewski CD (2002) Retinol measurements and retinoid receptor gene expression in patients with multiple sclerosis. *Mult Scler J* **8**, 452-458.
- [70] Kane MA, Napoli JL (2010) Quantification of endogenous retinoids. *Methods Mol Biol* **652**, 1-54.
- [71] Galter D, Buervenich S, Carmine A, Anvret M, Olson L (2003) ALDH1 mRNA: Presence in human dopamine neurons and decreases in substantia nigra in Parkinson's disease and in the ventral tegmental area in schizophrenia. *Neurobiol Dis* **14**, 637-647.
- [72] Grünblatt E, Zehetmayer S, Jacob CP, Müller T, Jost WH, Riederer P (2010) Pilot study: Peripheral biomarkers for diagnosing sporadic Parkinson's disease. *J Neural Transm* **117**, 1387-1393.
- [73] Liu G, Yu J, Ding J, Xie C, Sun L, Rudenko I, Zheng W, Sastry N, Luo J, Rudow G, Troncoso JC, Cai H (2014) Aldehyde dehydrogenase 1 defines and protects a nigrostriatal dopaminergic neuron subpopulation. *J Clin Invest* **124**, 3032-3046.
- [74] Samad TA, Krezel W, Chambon P, Borrelli E (1997) Regulation of dopaminergic pathways by retinoids: Activation of the D2 receptor promoter by members of the retinoic acid receptor-retinoid X receptor family. *Proc Natl Acad Sci U S A* **94**, 14349-14354.
- [75] Spathis AD, Asvos X, Ziavra D, Karampelas T, Topouzis S, Cournia Z, Qing X, Alexakos P, Smits LM, Dalla C, Rideout HJ, Schwamborn JC, Tamvakopoulos C, Fokas D, Vassilatis DK (2017) Nurr1:RXR α heterodimer activation as monotherapy for Parkinson's disease. *Proc Natl Acad Sci U S A* **114**, 3999-4004.
- [76] Krezel W, Ghyselinck N, Samad TA, Dupé V, Kastner P, Borrelli E, Chambon P (1998) Impaired locomotion and dopamine signaling in retinoid receptor mutant mice. *Science* **279**, 863-867.
- [77] Esteves M, Cristóvão AC, Saraiva T, Rocha SM, Baltazar G, Ferreira L, Bernardino L (2015) Retinoic acid-loaded polymeric nanoparticles induce neuroprotection in a mouse model for Parkinson's disease. *Front Aging Neurosci* **7**, 20.
- [78] Yin L-H, Shen H, Diaz-Ruiz O, Bäckman CM, Bae E, Yu S-J, Wang Y (2012) Early post-treatment with 9-cis retinoic acid reduces neurodegeneration of dopaminergic neurons in a rat model of Parkinson's disease. *BMC Neurosci* **13**, 120.
- [79] Reiner DJ, Yu S-J, Shen H, He Y, Bae E, Wang Y (2014) 9-Cis retinoic acid protects against methamphetamine-induced neurotoxicity in nigrostriatal dopamine neurons. *Neurotox Res* **25**, 248-261.
- [80] Spathis AD, Asvos X, Ziavra D, Karampelas T, Topouzis S, Cournia Z, Qing X, Alexakos P, Smits LM, Dalla C, Rideout HJ, Schwamborn JC, Tamvakopoulos C, Fokas D, Vassilatis DK (2017) Nurr1:RXR α heterodimer activation as monotherapy for Parkinson's disease. *Proc Natl Acad Sci U S A* **114**, 3999-4004.
- [81] Friling S, Bergsland M, Kjellander S (2009) Activation of Retinoid X Receptor increases dopamine cell survival in models for Parkinson's disease. *BMC Neurosci* **10**, 146.
- [82] Ono K, Yamada M (2007) Vitamin A potently destabilizes preformed alpha-synuclein fibrils *in vitro*: Implications for Lewy body diseases. *Neurobiol Dis* **25**, 446-454.
- [83] Yakunin E, Loeb V, Kisos H, Biala Y, Yehuda S, Yaari Y, Selkoe DJ, Sharon R (2012) α -Synuclein neuropathology is controlled by nuclear hormone receptors and enhanced by docosahexaenoic acid in a mouse model for Parkinson's disease: NHR controls α -Syn neuropathology. *Brain Pathol* **22**, 280-294.
- [84] Carta M, Stancampiano R, Tronci E, Collu M, Usiello A, Morelli M, Fadda F (2006) Vitamin A deficiency induces motor impairments and striatal cholinergic dysfunction in rats. *Neuroscience* **139**, 1163-1172.
- [85] Kitaoka K, Hattori A, Chikahisa S, Miyamoto K, Nakaya Y, Sei H (2007) Vitamin A deficiency induces a decrease in EEG delta power during sleep in mice. *Brain Res* **1150**, 121-130.
- [86] Kunzler A, Ribeiro CT, Gasparotto J, Petiz LL, da Rosa Silva HT, da Silva JD, Bortolin R, de Souza PO, Barreto F, Espitia-Perez P, Schnorr CE, Somensi N, Moreira JCF, Gelain DP (2019) The effects of retinol oral supplementation in 6-hydroxydopamine dopaminergic denervation model in Wistar rats. *Neurochem Int* **125**, 25-34.
- [87] Sies H (2015) Oxidative stress: A concept in redox biology and medicine. *Redox Biol* **4**, 180-183.
- [88] Zhang L, Wang X, Cueto R, Effi C, Zhang Y, Tan H, Qin X, Ji Y, Yang X, Wang H (2019) Biochemical basis and metabolic interplay of redox regulation. *Redox Biol* **26**, 101284.
- [89] Chang K-H, Chen C-M (2020) The role of oxidative stress in Parkinson's disease. *Antioxidants (Basel)* **9**, 597.
- [90] Michel PP, Hirsch EC, Hunot S (2016) Understanding dopaminergic cell death pathways in Parkinson disease. *Neuron* **90**, 675-691.
- [91] Sutachan JJ, Casas Z, Albarracin SL, Stab BR, Samudio I, Gonzalez J, Morales L, Barreto GE (2012) Cellular and molecular mechanisms of antioxidants in Parkinson's disease. *Nutr Neurosci* **15**, 120-126.
- [92] Surmeier DJ, Schumacker PT, Guzman JD, Ilijic E, Yang B, Zampese E (2017) Calcium and Parkinson's disease. *Biochem Biophys Res Commun* **483**, 1013-1019.
- [93] Guzman JN, Sanchez-Padilla J, Wokosin D, Kondapalli J, Ilijic E, Schumacker PT, Surmeier DJ (2010) Oxidant stress evoked by pacemaking in dopaminergic neurons is attenuated by DJ-1. *Nature* **468**, 696-700.
- [94] Burke WJ, Kumar VB, Pandey N, Panneton WM, Gan Q, Franko MW, O'Dell M, Li SW, Pan Y, Chung HD, Galvin JE (2008) Aggregation of alpha-synuclein by DOPAL, the monoamine oxidase metabolite of dopamine. *Acta Neuropathol (Berl)* **115**, 193-203.
- [95] Cha J-H, Yu Q-M, Seo J-S (2016) Vitamin A supplementation modifies the antioxidant system in rats. *Nutr Res Pract* **10**, 26-32.
- [96] Chang K-H, Cheng M-L, Chiang M-C, Chen C-M (2018) Lipophilic antioxidants in neurodegenerative diseases. *Clin Chim Acta* **485**, 79-87.
- [97] Palace VP, Khaper N, Qin Q, Singal PK (1999) Antioxidant potentials of vitamin A and carotenoids and their relevance to heart disease. *Free Radic Biol Med* **26**, 746-761.
- [98] Tao L, Huang K, Wang J, Xue Y, Zhou Y, He F, Shen Y, Wang J, Gu X, Ji K, Qian L, Guo X (2019) Retinol palmitate protects against myocardial ischemia/reperfusion injury via reducing oxidative stress and inhibiting apoptosis. *Am J Transl Res* **11**, 1510-1520.
- [99] de Oliveira MR (2015) Vitamin A and retinoids as mitochondrial toxicants. *Oxid Med Cell Longev* **2015**, 140267.
- [100] de Oliveira MR, Oliveira MWS, Behr GA, Hoff MLM, da Rocha RF, Moreira JCF (2009) Evaluation of the effects of vitamin A supplementation on adult rat substantia nigra and striatum redox and bioenergetic states: Mitochondrial impairment, increased 3-nitrotyrosine

- and alpha-synuclein, but decreased D2 receptor contents. *Prog Neuropsychopharmacol Biol Psychiatry* **33**, 353-362.
- [101] Pasquali MAB, Gelain DP, Oliveira MR, Behr GA, Motta LL, Rocha RF, Klamt F, Moreira JCF (2009) Vitamin A supplementation induces oxidative stress and decreases the immunoccontent of catalase and superoxide dismutase in rat lungs. *Exp Lung Res* **35**, 427-438.
- [102] Nolan YM, Sullivan AM, Toulouse A (2013) Parkinson's disease in the nuclear age of neuroinflammation. *Trends Mol Med* **19**, 187-196.
- [103] Huang Z, Liu Y, Qi G, Brand D, Zheng SG (2018) Role of vitamin A in the immune system. *J Clin Med* **7**, 258.
- [104] Katsuki H, Kurimoto E, Takemori S, Kurauchi Y, Hisatsune A, Isohama Y, Izumi Y, Kume T, Shudo K, Akaike A (2009) Retinoic acid receptor stimulation protects midbrain dopaminergic neurons from inflammatory degeneration via BDNF-mediated signaling. *J Neurochem* **110**, 707-718.
- [105] Mohammadzadeh Honarvar N, Saedisomeolia A, Abdolahi M, Shayeganrad A, Taheri Sangsari G, Hassanzadeh Rad B, Muench G (2017) Molecular anti-inflammatory mechanisms of retinoids and carotenoids in Alzheimer's disease: A review of current evidence. *J Mol Neurosci* **61**, 289-304.
- [106] Choi W-H, Ji K-A, Jeon S-B, Yang M-S, Kim H, Min K-J, Shong M, Jou I, Joe E-H (2005) Anti-inflammatory roles of retinoic acid in rat brain astrocytes: Suppression of interferon-gamma-induced JAK/STAT phosphorylation. *Biochem Biophys Res Commun* **329**, 125-131.
- [107] van Neerven S, Nemes A, Imholz P, Regen T, Denecke B, Johann S, Beyer C, Hanisch U-K, Mey J (2010) Inflammatory cytokine release of astrocytes *in vitro* is reduced by all-trans retinoic acid. *J Neuroimmunol* **229**, 169-179.
- [108] Fuzzati-Armentero MT, Cerri S, Blandini F (2019) Peripheral-central neuroimmune crosstalk in Parkinson's disease: What do patients and animal models tell us? *Front Neurol* **10**, 232.
- [109] Wang Q, Liu Y, Zhou J (2015) Neuroinflammation in Parkinson's disease and its potential as therapeutic target. *Transl Neurodegener* **4**, 19.
- [110] Obeso JA, Stamelou M, Goetz CG, Poewe W, Lang AE, Weintraub D, Burn D, Halliday GM, Bezard E, Przedborski S, Lehericy S, Brooks DJ, Rothwell JC, Hallett M, DeLong MR, Marras C, Tanner CM, Ross GW, Langston JW, Klein C, Bonifati V, Jankovic J, Lozano AM, Deuschl G, Bergman H, Tolosa E, Rodriguez-Violante M, Fahn S, Postuma RB, Berg D, Marek K, Standaert DG, Surmeier DJ, Olanow CW, Kordower JH, Calabresi P, Schapira AHV, Stoessl AJ (2017) Past, present, and future of Parkinson's disease: A special essay on the 200th Anniversary of the Shaking Palsy: The Shaking Palsy: Past, Present and Future. *Mov Disord* **32**, 1264-1310.
- [111] Requejo C, López-de-Ipiña K, Ruiz-Ortega JÁ, Fernández E, Calvo PM, Morera-Herreras T, Miguez C, Cardona-Grifoll L, Cepeda H, Ugedo L, Lafuente JV (2020) Changes in day/night activity in the 6-OHDA-induced experimental model of Parkinson's disease: Exploring prodromal biomarkers. *Front Neurosci* **14**, 590029.
- [112] Maret S (2005) Retinoic acid signaling affects cortical synchrony during sleep. *Science* **310**, 111-113.
- [113] Kitaoka K, Shimizu M, Shimizu N, Chikahisa S, Nakagomi M, Shudo K, Yoshizaki K, Séi H (2011) Retinoic acid receptor antagonist LE540 attenuates wakefulness via the dopamine D1 receptor in mice. *Brain Res* **1423**, 10-16.
- [114] Vallée A, Lecarpentier Y, Guillevin R, Vallée J-N (2020) Circadian rhythms, neuroinflammation and oxidative stress in the story of Parkinson's disease. *Cells* **9**, 314.
- [115] Ransom J, Morgan PJ, McCaffery PJ, Stoney PN (2014) The rhythm of retinoids in the brain. *J Neurochem* **129**, 366-376.
- [116] Ming G-L, Song H (2011) Adult neurogenesis in the mammalian brain: Significant answers and significant questions. *Neuron* **70**, 687-702.
- [117] Fon D, Zhou K, Ercole F, Fehr F, Marchesan S, Minter MR, Crack PJ, Finkelstein DI, Forsythe JS (2014) Nanofibrous scaffolds releasing a small molecule BDNF-mimetic for the re-direction of endogenous neuroblast migration in the brain. *Biomaterials* **35**, 2692-2712.
- [118] Jacobs S, Lie DC, DeCicco KL, Shi Y, DeLuca LM, Gage FH, Evans RM (2006) Retinoic acid is required early during adult neurogenesis in the dentate gyrus. *Proc Natl Acad Sci U S A* **103**, 3902-3907.
- [119] Bonhomme D, Minni AM, Alfos S, Roux P, Richard E, Higuere P, Moisan M-P, Pallet V, Touyarot K (2014) Vitamin A status regulates glucocorticoid availability in Wistar rats: Consequences on cognitive functions and hippocampal neurogenesis? *Front Behav Neurosci* **8**, 20.
- [120] Bonnet E, Touyarot K, Alfos S, Pallet V, Higuere P, Abrous DN (2008) Retinoic acid restores adult hippocampal neurogenesis and reverses spatial memory deficit in vitamin A deprived rats. *PLoS One* **3**, e3487.
- [121] Nasrolahi A, Mahmoudi J, Akbarzadeh A, Karimipour M, Sadigh-Eteghad S, Salehi R, Farhoudi M (2018) Neurotrophic factors hold promise for the future of Parkinson's disease treatment: Is there a light at the end of the tunnel? *Rev Neurosci* **29**, 475-489.
- [122] Nasrolahi A, Mahmoudi J, Karimipour M, Akbarzadeh A, Sadigh-Eteghad S, Salehi R, Farajdokht F, Farhoudi M (2019) Effect of cerebral dopamine neurotrophic factor on endogenous neural progenitor cell migration in a rat model of Parkinson's disease. *EXCLI J* **18**, 139-153.
- [123] Wang T-W, Zhang H, Parent JM (2005) Retinoic acid regulates postnatal neurogenesis in the murine subventricular zone-olfactory bulb pathway. *Development* **132**, 2721-2732.
- [124] Maia J, Santos T, Aday S, Agasse F, Cortes L, Malva JO, Bernardino L, Ferreira L (2011) Controlling the neuronal differentiation of stem cells by the intracellular delivery of retinoic acid-loaded nanoparticles. *ACS Nano* **5**, 97-106.
- [125] Santos T, Ferreira R, Quartin E, Boto C, Saraiva C, Bragança J, Peça J, Rodrigues C, Ferreira L, Bernardino L (2017) Blue light potentiates neurogenesis induced by retinoic acid-loaded responsive nanoparticles. *Acta Biomater* **59**, 293-302.
- [126] Santos T, Ferreira R, Maia J, Agasse F, Xapelli S, Cortes L, Bragança J, Malva JO, Ferreira L, Bernardino L (2012) Polymeric nanoparticles to control the differentiation of neural stem cells in the subventricular zone of the brain. *ACS Nano* **6**, 10463-10474.
- [127] Plane JM, Whitney JT, Schallert T, Parent JM (2008) Retinoic acid and environmental enrichment alter subventricular zone and striatal neurogenesis after stroke. *Exp Neurol* **214**, 125-134.
- [128] Gaillard A, Jaber M (2011) Rewiring the brain with cell transplantation in Parkinson's disease. *Trends Neurosci* **34**, 124-133.

- [129] Koppaka V, Thompson DC, Chen Y, Ellermann M, Nicolaou KC, Juvonen RO, Petersen D, Deitrich RA, Hurley TD, Vasilioi V (2012) Aldehyde dehydrogenase inhibitors: A comprehensive review of the pharmacology, mechanism of action, substrate specificity, and clinical application. *Pharmacol Rev* **64**, 520-539.
- [130] Goodman T, Crandall JE, Nanesco SE, Quadro L, Shearer K, Ross A, McCaffery P (2012) Patterning of retinoic acid signaling and cell proliferation in the hippocampus. *Hippocampus* **22**, 2171-2183.
- [131] Poulin J-F, Caronia G, Hofer C, Cui Q, Helm B, Ramakrishnan C, Chan CS, Dombek DA, Deisseroth K, Awatramani R (2018) Mapping projections of molecularly defined dopamine neuron subtypes using intersectional genetic approaches. *Nat Neurosci* **21**, 1260-1271.
- [132] Wu J, Kung J, Dong J, Chang L, Xie C, Habib A, Hawes S, Yang N, Chen V, Liu Z, Evans R, Liang B, Sun L, Ding J, Yu J, Saez-Atienzar S, Tang B, Khaliq Z, Lin D-T, Le W, Cai H (2019) Distinct connectivity and functionality of aldehyde dehydrogenase 1a1-positive nigrostriatal dopaminergic neurons in motor learning. *Cell Rep* **28**, 1167-1181.e7.
- [133] Anderson DW, Schray RC, Duester G, Schneider JS (2011) Functional significance of aldehyde dehydrogenase ALDH1A1 to the nigrostriatal dopamine system. *Brain Res* **1408**, 81-87.
- [134] Wey MC-Y, Fernandez E, Martinez PA, Sullivan P, Goldstein DS, Strong R (2012) Neurodegeneration and motor dysfunction in mice lacking cytosolic and mitochondrial aldehyde dehydrogenases: Implications for Parkinson's disease. *PLoS One* **7**, e31522.
- [135] Damier P, Hirsch EC, Agid Y, Graybiel AM (1999) The substantia nigra of the human brain. II. Patterns of loss of dopamine-containing neurons in Parkinson's disease. *Brain* **122** (Pt 8), 1437-1448.
- [136] Fearnley JM, Lees AJ (1991) Ageing and Parkinson's disease: Substantia nigra regional selectivity. *Brain* **114** (Pt 5), 2283-2301.
- [137] Cai H, Liu G, Sun L, Ding J (2014) Aldehyde Dehydrogenase 1 making molecular inroads into the differential vulnerability of nigrostriatal dopaminergic neuron subtypes in Parkinson's disease. *Transl Neurodegener* **3**, 27.
- [138] Fitzmaurice AG, Rhodes SL, Lulla A, Murphy NP, Lam HA, O'Donnell KC, Barnhill L, Casida JE, Cockburn M, Sagasti A, Stahl MC, Maidment NT, Ritz B, Bronstein JM (2013) Aldehyde dehydrogenase inhibition as a pathogenic mechanism in Parkinson disease. *Proc Natl Acad Sci U S A* **110**, 636-641.
- [139] Goldstein DS (2020) The catecholaldehyde hypothesis: Where MAO fits in. *J Neural Transm* **127**, 169-177.
- [140] Goldstein DS, Kopin IJ, Sharabi Y (2014) Catecholamine autotoxicity. Implications for pharmacology and therapeutics of Parkinson disease and related disorders. *Pharmacol Ther* **144**, 268-282.
- [141] Panneton WM, Kumar VB, Gan Q, Burke WJ, Galvin JE (2010) The neurotoxicity of DOPAL: Behavioral and stereological evidence for its role in Parkinson disease pathogenesis. *PLoS One* **5**, e15251.
- [142] Charles V, Boulvin S, Haven F, Gille M (2006) [Wernicke's encephalopathy associated with chronic disulfiram intoxication]. *Rev Neurol (Paris)* **162**, 1252-1256.
- [143] Laplane D, Attal N, Sauron B, de Billy A, Dubois B (1992) Lesions of basal ganglia due to disulfiram neurotoxicity. *J Neurol Neurosurg Psychiatry* **55**, 925-929.
- [144] Casida JE, Ford B, Jinsmaa Y, Sullivan P, Cooney A, Goldstein DS (2014) Benomyl, aldehyde dehydrogenase, DOPAL, and the catecholaldehyde hypothesis for the pathogenesis of Parkinson's disease. *Chem Res Toxicol* **27**, 1359-1361.
- [145] Fitzmaurice AG, Rhodes SL, Cockburn M, Ritz B, Bronstein JM (2014) Aldehyde dehydrogenase variation enhances effect of pesticides associated with Parkinson disease. *Neurology* **82**, 419-426.
- [146] Leiphon LJ, Picklo MJ (2007) Inhibition of aldehyde detoxification in CNS mitochondria by fungicides. *Neurotoxicology* **28**, 143-149.
- [147] Jacobs FMJ, Smits SM, Noorlander CW, von Oertel L, van der Linden AJA, Burbach JPH, Smid MP (2007) Retinoic acid counteracts developmental defects in the substantia nigra caused by Pitx3 deficiency. *Development* **134**, 2673-2684.
- [148] Smid MP (2017) Molecular programming of mesodiencephalic dopaminergic neuronal subsets. *Front Neuroanat* **11**, 59.
- [149] Hwang D (2003) Selective loss of dopaminergic neurons in the substantia nigra of Pitx3-deficient aphakia mice. *Mol Brain Res* **114**, 123-131.
- [150] Suarez LM, Alberquilla S, García-Montes JR, Moratalla R (2018) Differential synaptic remodeling by dopamine in direct and indirect striatal projection neurons in Pitx3^{-/-} mice, a genetic model of Parkinson's disease. *J Neurosci* **38**, 3619-3630.
- [151] van den Munkhof P (2003) Pitx3 is required for motor activity and for survival of a subset of midbrain dopaminergic neurons. *Development* **130**, 2535-2542.
- [152] Le W, Zhang L, Xie W, Li S, Dani JA (2015) Pitx3 deficiency produces decreased dopamine signaling and induces motor deficits in Pitx3(-/-) mice. *Neurobiol Aging* **36**, 3314-3320.
- [153] Li J, Dani JA, Le W (2009) The role of transcription factor Pitx3 in dopamine neuron development and Parkinson's disease. *Curr Top Med Chem* **9**, 855-859.
- [154] Kim J-I, Ganesan S, Luo SX, Wu Y-W, Park E, Huang EJ, Chen L, Ding JB (2015) Aldehyde dehydrogenase 1a1 mediates a GABA synthesis pathway in midbrain dopaminergic neurons. *Science* **350**, 102-106.
- [155] Furness JB (2006) The organisation of the autonomic nervous system: Peripheral connections. *Auton Neurosci* **130**, 1-5.
- [156] Cersosimo MG, Benarroch EE (2012) Pathological correlates of gastrointestinal dysfunction in Parkinson's disease. *Neurobiol Dis* **46**, 559-564.
- [157] Smith LM, Parr-Brownlie LC (2019) A neuroscience perspective of the gut theory of Parkinson's disease. *Eur J Neurosci* **49**, 817-823.
- [158] Braak H, Rüb U, Gai WP, Del Tredici K (2003) Idiopathic Parkinson's disease: Possible routes by which vulnerable neuronal types may be subject to neuroinvasion by an unknown pathogen. *J Neural Transm* **110**, 517-536.
- [159] Braak H, de Vos RAI, Bohl J, Del Tredici K (2006) Gastric alpha-synuclein immunoreactive inclusions in Meissner's and Auerbach's plexuses in cases staged for Parkinson's disease-related brain pathology. *Neurosci Lett* **396**, 67-72.
- [160] Holmqvist S, Chutna O, Bousset L, Aldrin-Kirk P, Li W, Björklund T, Wang Z-Y, Roybon L, Melki R, Li J-Y (2014) Direct evidence of Parkinson pathology spread from the gastrointestinal tract to the brain in rats. *Acta Neuropathol (Berl)* **128**, 805-820.

- [161] Fasano A, Visanji NP, Liu LWC, Lang AE, Pfeiffer RF (2015) Gastrointestinal dysfunction in Parkinson's disease. *Lancet Neurol* **14**, 625-639.
- [162] Shannon KM, Keshavarzian A, Mutlu E, Dodiya HB, Daian D, Jaglin JA, Kordower JH (2012) Alpha-synuclein in colonic submucosa in early untreated Parkinson's disease. *Mov Disord* **27**, 709-715.
- [163] Derkinderen P, Rouaud T, Lebouvier T, Bruley des Varannes S, Neunlist M, De Giorgio R (2011) Parkinson disease: The enteric nervous system spills its guts. *Neurology* **77**, 1761-1767.
- [164] Lionnet A, Leclair-Visonneau L, Neunlist M, Murayama S, Takao M, Adler CH, Derkinderen P, Beach TG (2018) Does Parkinson's disease start in the gut? *Acta Neuropathol (Berl)* **135**, 1-12.
- [165] Perez-Pardo P, Kliet T, Dodiya HB, Broersen LM, Garssen J, Keshavarzian A, Kraneveld AD (2017) The gut-brain axis in Parkinson's disease: Possibilities for food-based therapies. *Eur J Pharmacol* **817**, 86-95.
- [166] Borghammer P, Van Den Berge N (2019) Brain-first versus gut-first Parkinson's disease: A hypothesis. *J Parkinsons Dis* **9**, S281-S295.
- [167] Desmet A-S, Cirillo C, Tack J, Vandenberghe W, Vanden Berghe P (2017) Live calcium and mitochondrial imaging in the enteric nervous system of Parkinson patients and controls. *eLife* **6**, e26850.
- [168] Visanji NP, Marras C, Hazrati L-N, Liu LWC, Lang AE (2014) Alimentary, my dear Watson? The challenges of enteric α -synuclein as a Parkinson's disease biomarker. *Mov Disord* **29**, 444-450.
- [169] Ruffmann C, Parkkinen L (2016) Gut feelings about α -synuclein in gastrointestinal biopsies: Biomarker in the making? *Mov Disord* **31**, 193-202.
- [170] Sato Y, Heuckeroth RO (2008) Retinoic acid regulates murine enteric nervous system precursor proliferation, enhances neuronal precursor differentiation, and reduces neurite growth *in vitro*. *Dev Biol* **320**, 185-198.
- [171] Tan M, Yang T, Liu H, Xiao L, Li C, Zhu J, Chen J, Li T (2021) Maternal vitamin A deficiency impairs cholinergic and nitrergic neurons, leading to gastrointestinal dysfunction in rat offspring via RAR β . *Life Sci* **264**, 118688.
- [172] Wright-Jin EC, Grider JR, Duester G, Heuckeroth RO (2013) Retinaldehyde dehydrogenase enzymes regulate colon enteric nervous system structure and function. *Dev Biol* **381**, 28-37.
- [173] Hasegawa S, Goto S, Tsuji H, Okuno T, Asahara T, Nomoto K, Shibata A, Fujisawa Y, Minato T, Okamoto A, Ohno K, Hirayama M (2015) Intestinal dysbiosis and lowered serum lipopolysaccharide-binding protein in Parkinson's disease. *PLoS One* **10**, e0142164.
- [174] Keshavarzian A, Green SJ, Engen PA, Voigt RM, Naqib A, Forsyth CB, Mutlu E, Shannon KM (2015) Colonic bacterial composition in Parkinson's disease. *Mov Disord* **30**, 1351-1360.
- [175] Scheperjans F, Aho V, Pereira PAB, Koskinen K, Paulin L, Pekkonen E, Haapaniemi E, Kaakkola S, Eerola-Rautio J, Pohja M, Kinnunen E, Murros K, Auvinen P (2015) Gut microbiota are related to Parkinson's disease and clinical phenotype. *Mov Disord* **30**, 350-358.
- [176] Unger MM, Spiegel J, Dillmann K-U, Grundmann D, Philippeit H, Bürmann J, Faßbender K, Schwierz A, Schäfer K-H (2016) Short chain fatty acids and gut microbiota differ between patients with Parkinson's disease and age-matched controls. *Parkinsonism Relat Disord* **32**, 66-72.
- [177] Sampson TR, Debelius JW, Thron T, Janssen S, Shastri GG, Ilhan ZE, Challis C, Schretter CE, Rocha S, Gradinaru V, Chesselet M-F, Keshavarzian A, Shannon KM, Krajmalnik-Brown R, Wittung-Stafshede P, Knight R, Mazmanian SK (2016) Gut microbiota regulate motor deficits and neuroinflammation in a model of Parkinson's disease. *Cell* **167**, 1469-1480.e12.
- [178] Minato T, Maeda T, Fujisawa Y, Tsuji H, Nomoto K, Ohno K, Hirayama M (2017) Progression of Parkinson's disease is associated with gut dysbiosis: Two-year follow-up study. *PLoS One* **12**, e0187307.
- [179] Yang D, Zhao D, Ali Shah SZ, Wu W, Lai M, Zhang X, Li J, Guan Z, Zhao H, Li W, Gao H, Zhou X, Yang L (2019) The role of the gut microbiota in the pathogenesis of Parkinson's disease. *Front Neurol* **10**, 1155.
- [180] Jackson A, Forsyth CB, Shaikh M, Voigt RM, Engen PA, Ramirez V, Keshavarzian A (2019) Diet in Parkinson's disease: Critical role for the microbiome. *Front Neurol* **10**, 1245.
- [181] Zhou Y, Zhou J, Zhang Y, Tang J, Sun B, Xu W, Wang X, Chen Y, Sun Z (2020) Changes in intestinal microbiota are associated with islet function in a mouse model of dietary vitamin A deficiency. *J Diabetes Res* **2020**, 2354108.
- [182] Tian Y, Nichols RG, Cai J, Patterson AD, Cantorna MT (2018) Vitamin A deficiency in mice alters host and gut microbial metabolism leading to altered energy homeostasis. *J Nutr Biochem* **54**, 28-34.
- [183] Sun W, Chen J (2020) Vitamin A deficiency in the early-life periods alters a diversity of the colonic mucosal microbiota in rats. *Front Nutr* **7**, 14.
- [184] Hibberd MC, Wu M, Rodionov DA, Li X, Cheng J, Griffin NW, Barratt MJ, Giannone RJ, Hettich RL, Osterman AL, Gordon JI (2017) The effects of micronutrient deficiencies on bacterial species from the human gut microbiota. *Sci Transl Med* **9**, eaal4069.
- [185] Konieczna P, Ferstl R, Ziegler M, Frei R, Nehrbass D, Lauener RP, Akdis CA, O'Mahony L (2013) Immunomodulation by *Bifidobacterium infantis* 35624 in the murine lamina propria requires retinoic acid-dependent and independent mechanisms. *PLoS One* **8**, e62617.
- [186] Reichrath J, Lehmann B, Carlberg C, Varani J, Zouboulis C (2007) Vitamins as hormones. *Horm Metab Res* **39**, 71-84.
- [187] Enderlin V, Higuieret D, Alfos S, Husson M, Jaffard R, Higuieret P, Pallet V (2000) Vitamin A deficiency decreases the expression of RAR β and RXR β/γ in adult mouse brain: Effect of RA administration. *Nutr Neurosci* **3**, 173-181.
- [188] Pallet V, Azais-Braesco V, Enderlin V, Grolier P, Noël-Suberville C, Garcin H, Higuieret P (1997) Aging decreases retinoic acid and triiodothyronine nuclear expression in rat liver: Exogenous retinol and retinoic acid differentially modulate this decreased expression. *Mech Ageing Dev* **99**, 123-136.
- [189] Brossaud J, Pallet V, Corcuff J-B (2017) Vitamin A, endocrine tissues and hormones: Interplay and interactions. *Endocr Connect* **6**, R121-R130.
- [190] Enderlin V, Pallet V, Alfos S, Dargelos E, Jaffard R, Garcin H, Higuieret P (1997) Age-related decreases in mRNA for brain nuclear receptors and target genes are reversed by retinoic acid treatment. *Neurosci Lett* **229**, 125-129.
- [191] Husson M, Enderlin V, Alfos S, Boucheron C, Pallet V, Higuieret P (2004) Expression of neurogranin and neuro-modulin is affected in the striatum of vitamin A-deprived rats. *Mol Brain Res* **123**, 7-17.

- [192] Enderlin V, Alfos S, Pallet V, Garcin H, Azais-Braesco V, Jaffard R, Higuieret P (1997) Aging decreases the abundance of retinoic acid (RAR) and triiodothyronine (TR) nuclear receptor mRNA in rat brain: Effect of the administration of retinoids. *FEBS Lett* **412**, 629-632.
- [193] Feart C, Pallet V, Boucheron C, Higuieret D, Alfos S, Letenneur L, Dartigues JF, Higuieret P (2005) Aging affects the retinoic acid and the triiodothyronine nuclear receptor mRNA expression in human peripheral blood mononuclear cells. *Eur J Endocrinol* **152**, 449-458.
- [194] Villanueva I, Alva-Sánchez C, Pacheco-Rosado J (2013) The role of thyroid hormones as inducers of oxidative stress and neurodegeneration. *Oxid Med Cell Longev* **2013**, 15.
- [195] Vairetti M, Ferrigno A, Rizzo V, Ambrosi G, Bianchi A, Richelmi P, Blandini F, Armentero M-T (2012) Impaired hepatic function and central dopaminergic denervation in a rodent model of Parkinson's disease: A self-perpetuating crosstalk? *Biochim Biophys Acta* **1822**, 176-184.
- [196] Tandeter H, Levy A, Gutman G, Shvartzman P (2001) Subclinical thyroid disease in patients with Parkinson's disease. *Arch Gerontol Geriatr* **33**, 295-300.
- [197] Umehara T, Matsuno H, Toyoda C, Oka H (2015) Thyroid hormone level is associated with motor symptoms in de novo Parkinson's disease. *J Neurol* **262**, 1762-1768.
- [198] Munhoz RP, Teive HAG, Troiano AR, Hauck PR, Herdoiza Leiva MHB, Graff H, Werneck LC (2004) Parkinson's disease and thyroid dysfunction. *Parkinsonism Relat Disord* **10**, 381-383.
- [199] Fernandes de Abreu DA, Eyles D, Féron F (2009) Vitamin D, a neuro-immunomodulator: Implications for neurodegenerative and autoimmune diseases. *Psychoneuroendocrinology* **34 Suppl 1**, S265-277.
- [200] Zhou Z, Zhou R, Zhang Z, Li K (2019) The association between vitamin D status, vitamin D supplementation, sunlight exposure, and Parkinson's disease: A systematic review and meta-analysis. *Med Sci Monit Int Med J Exp Clin Res* **25**, 666-674.
- [201] Rimmelzwaan LM, van Schoor NM, Lips P, Berendse HW, Eekhoff EMW (2016) Systematic review of the relationship between vitamin D and Parkinson's disease. *J Parkinsons Dis* **6**, 29-37.
- [202] Sleeman I, Aspray T, Lawson R, Coleman S, Duncan G, Khoo TK, Schoenmakers I, Rochester L, Burn D, Yarnall A (2017) The role of vitamin D in disease progression in early Parkinson's disease. *J Parkinsons Dis* **7**, 669-675.
- [203] Eyles DW, Smith S, Kinobe R, Hewison M, McGrath JJ (2005) Distribution of the vitamin D receptor and 1 α -hydroxylase in human brain. *J Chem Neuroanat* **29**, 21-30.
- [204] Shinpo K, Kikuchi S, Sasaki H, Moriwaka F, Tashiro K (2000) Effect of 1,25-dihydroxyvitamin D(3) on cultured mesencephalic dopaminergic neurons to the combined toxicity caused by L-buthionine sulfoximine and 1-methyl-4-phenylpyridine. *J Neurosci Res* **62**, 374-382.
- [205] Smith MP, Fletcher-Turner A, Yurek DM, Cass WA (2006) Calcitriol protection against dopamine loss induced by intracerebroventricular administration of 6-hydroxydopamine. *Neurochem Res* **31**, 533-539.
- [206] Wang JY, Wu JN, Cherng TL, Hoffer BJ, Chen HH, Borlongan CV, Wang Y (2001) Vitamin D(3) attenuates 6-hydroxydopamine-induced neurotoxicity in rats. *Brain Res* **904**, 67-75.
- [207] Foy CJ, Passmore AP, Vahidassr MD, Young IS, Lawson JT (1999) Plasma chain-breaking antioxidants in Alzheimer's disease, vascular dementia and Parkinson's disease. *QJM* **92**, 39-45.