

## Erratum to: DNA damage in normal-weight obese syndrome measured by Comet assay

Barbara Tomasello · Giuseppe Malfa ·  
Fabio Galvano · Marcella Renis

Published online: 9 January 2011  
© Springer-Verlag 2011

**Erratum to: Mediterr J Nutr Metab**  
**DOI 10.1007/s12349-010-0035-6**

The original publication of this article contained some incorrect data in Table 2. The corrected data are given below:

**Table 2** Determination of Fpg-sensible sites assessed in lymphocytes by FPG-Comet assay at pH 12.1

	NW	NWO	OB
Buffer pH 12.1	18% ± 0.9	27.3% ± 1.08*	28.12% ± 1.40*
Breaks/10 <sup>9</sup> Da	0.5	0.83	0.91
FPG pH 12.1	21.8% ± 1.10	37.8% ± 1.61* <sup>#</sup>	35.6% ± 1.04* <sup>#</sup>
Breaks/10 <sup>9</sup> Da	0.61	1.25	1.16
Fpg-sites/10 <sup>9</sup> Da	0.11	0.36	0.25
8-Oxo-dGua/10 <sup>6</sup> dG	0.067	0.22	0.15

Values expressed as %TDNA (mean ± SEM). Each data is averaged from three different experiments in which each sample was processed in duplicate. %TDNA are also converted in: (a) breaks/10<sup>9</sup> Da using for calculation of DNA break frequencies the calibration curve reported by Collins et al. [31]; (b) 8-oxo-dGua/10<sup>6</sup> dG using as conversion factor 1 8-oxo-Gua per 10<sup>9</sup> Da is equivalent to 1 8-oxo-Gua per 0.61 × 10<sup>6</sup> Gua as reported by Griffiths et al. [29]

\*  $P < 0.001$  significant versus correspondent NW values at pH 12.1

#  $P < 0.001$  significant versus correspondent buffer values

The online version of the original article can be found under  
doi:10.1007/s12349-010-0035-6.

B. Tomasello · G. Malfa · F. Galvano · M. Renis (✉)  
Department of Biological Chemistry,  
Medical Chemistry and Molecular Biology,  
University of Catania, Viale A. Doria, 6,  
95125 Catania, Italy  
e-mail: renis@unict.it