

Erratum

Erratum to: Screening for possible hypopituitarism following mild traumatic brain injury: The first all-female study. Who do we need to evaluate further?

Lára Ósk Eggertsdóttir Claessen, Hafrún Kristjánsdóttir, María K. Jónsdóttir, Sigrún Helga Lund, Ingunn S.U. Kristensen and Helga Ágústa Sigurjónsdóttir

[NeuroRehabilitation, 52(2), 2023, 259–271, DOI 10.3233/NRE-220194]
<https://content.iospress.com/articles/neurorehabilitation/nre220194>

On page 264 there was an error in the layout of Table 2; some data was misplaced in the wrong column. The correct Table 2 is as follows:

		Total (n = 133)	Evaluation by an endocrinologist		OR [95%CI]	P value
			No (n = 45)	Yes (n = 88)		
Still playing	Yes	72 (54.1%)	28 (62.2%)	44 (50.9%)	---	0.182
	No	61 (45.9%)	17 (37.8%)	44 (50.0%)		
Age**		29.2 (17; 46)	31.2 (20.0; 46.0)	28.2 (17.0; 46.0)	0.94 [0.95 – 0.99]	0.034*
BMI, mean (SD)		26.2 (4.7)	25.8 (3.7)	26.4 (5.2)	---	0.442
SBP, mean (SD)		124 (12.0)	123 (12.5)	125 (11.8)	---	0.361
DBP, mean (SD)		78.3 (9.1)	77 (7.7)	78.9 (9.7)	---	0.268
Previous medical history	Yes	56 (42.1%)	18 (40.0%)	38 (43.2%)	---	0.725
	No	77 (57.9%)	27 (60.0%)	50 (56.8%)		
Previous hormonal disease	Yes	13 (9.8%)	4 (8.9%)	9 (10.2%)	---	0.806
	No	120 (90.2%)	41 (91.1%)	79 (89.8%)		
HoC	No	54 (40.6%)	18 (40.0%)	36 (40.9%)	---	0.442
	Before mTBI	21 (15.8%)	9 (20.0%)	12 (13.6%)		
	After mTBI	22 (16.5%)	10 (22.2%)	12 (13.6%)		
Menstrual changes	Before and after mTBI	36 (27.1%)	8 (17.8%)	28 (31.8%)	---	0.257
	After mTBI	17 (12.8%)	5 (11.1%)	12 (13.6%)		
Menstrual changes	No changes	116 (87.2%)	40 (88.9%)	76 (86.4%)	---	0.680
Time from mTBI (years) **		5.0 (0.04; 35.2)	5.8 (0.04; 35.2)	4.7 (0.04; 25.6)	---	0.308
Number of mTBI **		2.2 (1;4)	2.2 (1;4)	2.1 (1;4)	---	0.568
Number of mTBI symptoms **		3.9 (0; 13)	2.9 (0;12)	4.3 (0;13)	1.15 [1.01 – 1.33]	0.045*